

Estimating Greenhouse Gas Emissions in Agriculture

A Manual to Address Data Requirements for Developing Countries



Cover photo:

©FAO/Daniel Hayduk

©FAO/Rocío Dánica Cóndor Golec

©FAO/Sia Kambou

Estimating Greenhouse Gas Emissions In Agriculture

A Manual to Address Data Requirements for Developing Countries

Francesco N. Tubiello, Rocío D. Cóndor-Golec, Mirella Salvatore, Angela Piersante, Sandro Federici, Alessandro Ferrara, Simone Rossi, Alessandro Flammini, Paola Cardenas, Riccardo Biancalani, Heather Jacobs, Paulina Prasula, and Paolo Prosperi.

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

ISBN 978-92-5-108674-2

© FAO, 2015

FAO encourages the use, reproduction and dissemination of material in this information product. Except where otherwise indicated, material may be copied, downloaded and printed for private study, research and teaching purposes, or for use in non-commercial products or services, provided that appropriate acknowledgement of FAO as the source and copyright holder is given and that FAO's endorsement of users' views, products or services is not implied in any way.

All requests for translation and adaptation rights, and for resale and other commercial use rights should be made via www.fao.org/contact-us/licence-request or addressed to copyright@fao.org.

FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org.

Contents

Acknov Acrony	wledgments yms	vii viii
1. Intro	oduction	1
2. The	e Institutional Framework	3
2.1	The UN Framework Convention on Climate Change	3
2.2	•	
	forestry and other land use sector under the UNFCCC	4
2.3	·	6
3. The	e IPCC Guidelines for National GHG Inventories	9
3.1	The Structure of IPCC Guidelines	9
3.2	The quality of the Emission Inventory	10
3.3	The AFOLU sector and its structure	12
3.4	Reporting methods	16
3.5	Estimation process	17
	3.5.1 The identification of key categories	18
	3.5.2 The selection of methods and measurement	19
	3.5.3 The selection and collection of activity data	20
	3.5.4 The selection of emission factors or carbon stock change factors	21
4. Add	dressing data and methodological requirements	23
4.1	Activity data sources	24
4.2	Emission factors	29
4.3	Metadata and emissions categories for agriculture	29
4.4	Metadata and emissions categories for forestry and land use	33
5. Min	nimum data availability for developing countries:	
Stru	ucture and data organization using FAOSTAT as guide	35
5.1	3 ** ** *	45
	5.1.1 Enteric fermentation	45
	5.1.2 Manure management	51
	5.1.3 Rice cultivation	58
	5.1.4 Synthetic fertilizers	61
	5.1.5 Manure applied to soils	67
	5.1.6 Manure left on pasture	73
	5.1.7 Crop residues	78
	5.1.8 Cultivation of organic soils	85
	5.1.9 Burning crop residues	88
	5.1.10 Burning-Savanna	93
	5.1.11 Energy use	96
5.2	•	101
	5.2.1 Forest land	101
	5.2.2 Cropland	110
	5.2.3 Grassland	113
	5.2.4 Burning-Biomass	115

Annexe		
1.	Glossary: concepts, definitions and symbols	120
2.	Data activities: example of national questionnaire on data collection	129
3.	Parameters for the different emission categories	131
4.	Overview of existing reporting requirements for mitigation in the agriculture,	
	land use sector under the UNFCCC	174
Referen	ices	179
FIGURE	S	
1.	Processes linked to national GHG reporting	6
	Main source and sink categories of 1996 and 2006 IPCC Guidelines	9
	Evolution of IPCC Guidelines for National GHG Emission Inventories	10
	Main steps in compiling a GHG Inventory	11
	GHG emission sources/removals and processes in managed ecosystems	12
6.	Mapping of AFOLU categories across IPCC Guidelines	15
7.	Categorization of Sources and Sinks of IPCC Guidelines reporting tables, 1996 - 2006	16
8.	Equation to calculate GHG emissions/removals	16
9.	Decision tree for the livestock category	20
10.	Map of savanna burned areas in 2012	26
11.	Savanna dry biomass burned in 2012	27
12.	Map of N ₂ O emissions from savanna burned in 2012	28
13.	FAOSTAT database homepage	36
14.	FAOSTAT homepage	37
15.	The FAOSTAT Emissions database	38
16.	Metadata in the FAOSTAT Emissions database	39
	The "Browse data" option	40
	The "Download data" option	40
	Example "Browse data"	41
	Example "Download data"	43
	"Number of heads of sheep" from the FAOSTAT database	46
	"Number of heads of dairy cattle" from FAOSTAT database	47
	"Number of heads of cattle" from the FAOSTAT database	47
	Livestock sub-categories of the FAOSTAT Emissions database	48
	"Stock of laying animals" from the FAOSTAT database	53
	"Stock of chickens" from the FAOSTAT database	54
	Manure management items from the FAOSTAT Emissions database	54
	Paddy rice area harvested, from the FAOSTAT database	59
	Fertilizers archive from the FAOSTAT database	63
	Fertilizers dataset from the FAOSTAT database	63
	Synthetic fertilizers items from the FAOSTAT Emissions database	64
	Manure applied to soils items from the FAOSTAT Emissions database	69
	Manure left on pasture items from the FAOSTAT Emissions database	75
	Area harvested and yield data	80
	Crops residues items	80
	Organic soils data from the FAOSTAT database	87
	Area harvested data from the FAOSTAT database	90
38.	Burning crops items	91

40. Data on the area equipped for power irrigation from the AQUASTAT database $\,$

41. The "Global Forest Resources Assessment 2010" homepage

94

98

103

39. Area burned data

42.	Forestry area from the FRA database	104
43.	Total forest area in FAOSTAT	105
44.	Carbon stock dataset from the FRA database	106
45.	Cropland items from the FAOSTAT Emissions database	111
46.	Grassland items from the FAOSTAT Emissions database	114
47.	Burning-Biomass items from the FAOSTAT Emissions database	118
48.	Example of a FAOSTAT Questionnaire	129
TABLES		
1.	Categorization of sources and sinks of IPCC Guidelines reporting tables, 1996-2006	16
2.	Tier 1 CH ₄ emission factors from enteric fermentation for cattle	22
3.	FAOSTAT emissions database sources of activity data	24
4.	Data structure in the FAOSTAT "Emissions – Agriculture" domain	30
5.	Data structure of the "Emissions – Land Use" domain	34
6.	Tier 1 enteric fermentation emissions factors for cattle	50
	Results of livestock enteric fermentation (cattle) in Morocco in the year 2010	50
	Emission factor for methane (kg CH ₄ head ⁻¹), by animal category and IPCC area	131
	Emission factor for methane for each animal category, kg CH ₄ head ⁻¹ , by country	132
	Nitrogen excreted in manure for animal category, kg N head-1 by IPCC area	139
4A.	Nitrogen excretion rate (kg N (1000 kg animal mass) ⁻¹ day ⁻¹), by animal category and IPCC area	139
	Share of nitrogen manure treated in MS lagoon, by animal category and IPCC area	140
6A.	Share of nitrogen manure treated in MS slurry, by animal category and IPCC area	140
	Share of nitrogen manure treated in MS solid storage, by animal category and IPCC area	141
	Share of nitrogen manure treated in MS dry lot, by animal category and IPCC area	141
	Share of nitrogen manure treated in MS daily spread, by animal category and IPCC area	142
	Share of nitrogen manure treated in MS digester, by animal category and IPCC area	142
	Share of nitrogen manure treated in MS pit below one month, by animal category and IPCC area	143
	Share of nitrogen manure treated in MS pit above one month, by animal category and IPCC area	143
	Share of nitrogen manure treated in other MS, by animal category and IPCC area	144
	Fraction of nitrogen manure from MS slurry applied to soil, by animal category and IPCC area	144
	Fraction of nitrogen manure from MS lagoon applied to soil, by animal category and IPCC area	145
	Fraction of applied organic N fertilizer in MS solid storage or each animal category, by IPCC area	145
	Fraction of nitrogen manure from MS dry lot applied to soil, by animal category and IPCC area	146
	Fraction of nitrogen manure from MS daily spread applied to soil, by animal category and IPCC area	146
	Fraction of nitrogen manure from MS digester applied to soil, by animal category and IPCC area	147
20A.	Fraction of nitrogen manure from MS pit below and above one month applied to soil,	4.47
01.4	by animal category and IPCC area	147
	Fraction of nitrogen manure from other MS applied to soil, by animal category and IPCC area	148
	Emission factor for direct N ₂ O emissions (kg N ₂ O-N (kg N) ⁻¹), by MS and IPCC area	148
	Percent of nitrogen losses due to runoff and leaching in MS, by IPCC area	149
24A.	Emission factor for indirect N_2O emissions from atmospheric deposition (kg N_2O -N (kg N_3 -N + NO_x -N) ⁻¹), by IPCC area	140
25.4		149
25A.	Emission factor for indirect N ₂ O emissions from leaching and runoff	140
264	(kg N ₂ O–N (kg N) ⁻¹), by IPCC area	149
∠bA.	Emission factor for direct N2O emissions from nitrogen inputs (kg N ₂ O–N (kg N input) ⁻¹), by IPCC area	150
27^		150
ZIA	Percent of nitrogen losses due to runoff and leaching in managed soils and pasture (kg N (kg N additions or deposition, by grazing animals) ⁻¹), by IPCC area	150
201	Fraction of synthetic nitrogen fertilizer that volatilises	100
20A.	(kg N volatilised (kg of N applied) ⁻¹), by IPCC area	150
	ing it relatinged ing of it applied/ if, by it be also	100

29A.	Water regimes (ha), by country	151
30A.	Seasonal emission factor of methane(g m ⁻²), by country	152
31A.	Amount of nitrogen from bedding (kg N animal ⁻¹), by animal category and IPCC area	154
32A.	Fraction of nitrogen manure lost in MS lagoon, by animal category and IPCC area	154
33A.	Fraction of nitrogen manure lost in MS slurry, by animal category and IPCC area	155
34A.	Fraction of nitrogen manure lost in MS solid storage, by animal category and IPCC area	155
35A.	Fraction of nitrogen manure lost in MS dry lot, by animal category and IPCC area	156
36A.	Fraction of nitrogen manure lost in MS daily spread, by animal category and IPCC area	156
37A.	Fraction of nitrogen manure lost in MS pit below and above one month,	
	by animal category and IPCC area	157
38A.	Fraction of nitrogen manure, urine and dung deposited on pastures that volatilises	
	(kg N volatilised (kg of N applied or deposited) ⁻¹), by animal category and IPCC area	157
39A.	Fraction of managed manure used for feed, by animal category and IPCC area	158
40A.	Fraction of managed manure used for fuel, by animal category and IPCC area	158
41A.	Fraction of managed manure used for construction, by animal category and IPCC area	159
42A.	Fraction of nitrogen manure deposited on pastures, ranges and paddocks,	
	by animal category and IPCC area	159
43A.	Share of manure of dung and urine excreted on fields and burned for fuel,	
	by animal category and IPCC area	160
44A.	Emission factor for N ₂ O emissions from urine and dung deposited on pastures	
	(kg N ₂ O–N (kg N input) ⁻¹), by animal category and IPCC area	160
45A.	Dry matter fraction of harvested product, by crop and IPCC area	161
46A.	Slope element, by crop and IPCC area	161
47A.	Intercept element, by crop and IPCC area	162
48A.	Ratio of below-ground residues to above-ground biomass, by crop and IPCC area	162
49A.	Combustion factor, by crop and IPCC area	163
50A.	Nitrogen content of above-ground residues, by crop and IPCC area	163
51A.	Ratio of below-ground residues to harvested yield, by crop and IPCC area	164
52A.	Fuel biomass consumption value(t d.m ha ⁻¹), by crop	164
53A.	Emission factors for burning agricultural residues (g kg ⁻¹ d.m. burnt)	164
54A.	Emission factors and net calorific value, by energy carrier	165
55A.	Average carbon stock in forest living biomass (t ha ⁻¹), by region and year	165
56A.	Total carbon stock in forest living biomass (M t), by country	166
57A.	Emission factor for N ₂ O emissions from cropland and grassland organic soils, by climatic class	170
58A.	Emission factor for C emissions from cropland organic soils, by climatic class	171
59A.	Emission factor for C emissions from grassland organic soils, by climatic class	171
60A.	Correspondence between IPCC and FAOSTAT Nomenclatures of Animals categories	172

Acknowledgments

This document is a collaboration between the Climate, Energy and Tenure Division and the Statistics Division of FAO. The authors are members of the FAO *Monitoring and Assessment of GHG Emissions in Agriculture Project* (MAGHG): Francesco N Tubiello, Rocío D. Cóndor-Golec, Mirella Salvatore, Angela Piersante, Sandro Federici, Alessandro Ferrara, Simone Rossi, Alessandro Flammini, Paola Cardenas, Riccardo Biancalani, Heather Jacobs, Paulina Prasula, and Paolo Prosperi.

The work leading to this document benefited from many interactions with stakeholders in member countries, who provided insight into their technical and institutional needs towards strengthening national capacity for rural statistics and greenhouse gas emission estimations. It would be impossible to name them all here: contributors include representatives from over seventy countries who participated in a series of Regional Capacity Development Workshops on Improving Statistics for Greenhouse Gases, held between 2012 and 2014, including the Inception Workshop on GHG Statistics (Da Lat, Vietnam, Oct 2012); The Second Workshop on Statistics for GHG Emissions (Port of Spain, Trinidad and Tobago, Jun 2013); the Third FAO Workshop on Statistics for GHG Emissions (Casablanca, Morocco, Dec 2013) and the Meso-American Workshop on National Emission Inventories and Mitigation Plans in Agriculture, Land Use, Land Use Change and Forestry (San Jose, Costa Rica, Jun 2014).

Additionally, several colleagues at FAO and at other institutions made many useful suggestions and comments during peer review that greatly improved this work. In particular, we wish to thank Mr Simon Eggleston (Global Forest Observations Initiative), Ms Kimberly Todd (United Nations Development Programme), Ms Eleonora di Cristofaro (Italian Institute for Environmental Protection and Research) and Ms Laura Meza (FAO Regional Representative Latin America and the Caribbean).

This document was made possible by generous funding by the Governments of Germany and Norway to the *Monitoring and Assessment of GHG Emissions in Agriculture Project* (MAGHG), Trust Funds GCP/GLO/286/ GER and GCP/GLO/325/NOR, and by the *Global Strategy to Improve Agricultural and Rural Statistics* for the development of common standards and technical guidelines.

Acronyms

AFOLU Agriculture, Forestry and Other Land Use

AD Activity Data

BUR Biennial Update Report

CGRN Coalition for Rainforest Nations
COP Conference of the Parties
CRF Common Report Format

CS Country Specific
DOM Dead Organic Matter

ES Economic and Social Development Department of FAO

EF Emission Factors
EIT Economies in Transition

ESS Economic and Social Statistics Division of FAO

FAO Food and Agriculture Organization of the United Nations FRA Global Forest Resource Assessment database of FAO

FO Forestry Department of FAO
FOLU Forestry and Other Land Use
GAEZ Global Agro-Ecological Zones
GFED4 Global Fire Emission Database 4
GHGI Greenhouse Gas Inventory

GHG Greenhouse Gas

GPG Good Practice Guidance

GS Global Strategy

GWP Global Warming Potential

HWSD Harmonized World Soil Database
IEA International Energy Agency
IEF Implied Emission Factor

IPCC Intergovernmental Panel on Climate Change

LDCs Least Developed Countries

LECB Low Emissions Capacity Building

LULUCF Land Use, Land-Use Change and Forestry

MAGHG Monitoring and Assessment of GHG Emissions and Mitigation Potential in Agriculture

MDG Millennium Development Goals
MMS Manure Management Systems

MOP Meeting of the Parties

MRV Monitoring Reporting and Verification
NAMAs Nationally Appropriate Mitigation Actions

NCs National Communications

NGHGI National Greenhouse Gas Inventory NGOs Non-governmental organizations NIR National Inventory Report

NRC Climate, Energy and Tenure Division of FAO

OECD Organization for Economic Co-operation and Development

REDD+ Reduced Emissions from Deforestation and Forest Degradation in developing countries;

and the role of conservation, sustainable management of forests and enhancement

of forest carbon stocks in developing countries

SIDS Small Island Developing States

SBI Subsidiary Body Implementation

SBSTA Subsidiary Body for Scientific and Technological Advice

SC Steering Committee SOM Soil Organic Matter

TFI Task Force on National Greenhouse Gas Inventories

UNCED United Nations Conference on Environment and Development

UNDP United Nations Development Program

UNFCCC United Nations Framework Convention on Climate Change

UNREDD United Nations Program on Reduction of Emissions from Deforestation and Forest Degradation

WG Working Group

Introduction

Countries report their greenhouse gas (GHG) emissions and removals from all sectors via national GHG Inventories, submitted to the United Nations Framework Convention on Climate Change (UNFCCC) in accordance with international climate policy agreements and technical guidelines developed by the Intergovernmental Panel on Climate Change (IPCC).

The agriculture sector represents a unique challenge for national inventory compilers, especially in developing countries, due to significant difficulties in compiling and regularly updating national statistics for agriculture, forestry and land use —the first necessary step in preparing national GHG estimates.

The limited capacity to identify and collect reliable activity data and to quantify emissions by sources and removals by sinks, including in countries where agriculture and land use activities are a key component of the national economy and a driver of employment, could furthermore lead to limited access to international climate finance of importance to rural development, such as for instance REDD+ activities and Nationally Appropriate Mitigation Actions (NAMAs).

FAO supports its Member Countries with data, guidelines and technical expertise towards an enhanced global knowledge base on GHG emissions and mitigation potentials. The *Monitoring and Assessment of GHG Emissions and Mitigation Potentials in Agriculture – MAGHG* Project of the Climate, Energy and Land Tenure Division's Mitigation Programme (MICCA), in close collaboration with the FAO Statistics Division and the FAO Forestry Division UN REDD Programme, has developed and made available relevant activity data, GHG emission estimates databases and analysis tools through the FAOSTAT database. These products are used in regional and country-level capacity development activities that support practitioners in assessing and reporting GHG emissions from agriculture and land use categories, with a view to strengthening their national processes, with a focus on preparation and submission of GHG Inventories, Biennial Update Reports (BURs) and NAMAs.

This Manual provides Member Countries with a tool and methodology to help identify, build and access the minimum set of activity data needed for GHG estimation. Required data is largely drawn from country's official national agricultural and forestry statistics, as disseminated in FAO's corporate database FAOSTAT, and integrated by geo-spatial data obtained from recognized international sources. Users are provided with step-by-step guidance on how to use this minimum set to build a default, yet complete national GHG emission dataset for agriculture and land use, which follows the default, Tier 1 approach of the Intergovernmental Panel on Climate Change (IPCC) Guidelines on National GHG Inventories.

This Manual therefore contributes to FAO and the Global Strategy support to national processes towards improved agricultural and rural statistics. It can be used as a guide by staff of national statistical offices, environmental ministries and other relevant national agencies, to understand the international context of international climate policy (Ch. 2) and international guidelines (Ch. 3), identify needs for improved agricultural and rural data as well as emission estimates towards improving GHG Inventories (Ch. 4), while supplying practical information and examples based on accessing and using the FAOSTAT Emissions database for agriculture and land use (Ch. 5).

Improving statistical processes for GHG estimation has wider implications beyond climate change mitigation. Improved statistics on agricultural and land use activities enable Member Countries better identify climate responses that are consistent with their rural development and food security objectives, including preserving natural resources, increasing resilience of production systems and creating new employment opportunities.

The Institutional Framework

2.1 THE UN FRAMEWORK CONVENTION ON CLIMATE CHANGE

UN Member States meeting at the United Nations Conference on Environment and Development (UNCED, Rio de Janeiro, 1992), agreed to sign a Declaration on sustainable development as an instrument to ensure a healthy and productive life for human beings, in harmony with nature, for current and future generations. Sustainable Development commitments of UN Member States include the need to ensure that activities within their jurisdiction or control do not cause damage to the environment, both within their territories as well as of other States or areas beyond the limits of their national jurisdiction.

For these reasons, UNCED signatories also signed the United Nations Framework Convention on Climate Change (UNFCCC), with the commitment to *stabilize greenhouse gas* (*GHG*) concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system, within a time-frame sufficient to allow ecosystems to adapt naturally to climate change, to ensure that food production is not threatened, and to enable economic development to proceed in a sustainable manner (UNFCCC, Art. 2) To achieve these global sustainable development goals in the Agriculture, Forestry and Other Land Use sector (AFOLU), Parties to the UNFCCC shall promote sustainable management, and promote and cooperate in the conservation and enhancement, as appropriate, of sinks and reservoirs, including biomass, forests, as well as other terrestrial and coastal ecosystems.

In 1997, to strengthen the global response to climate change, the countries that were party to this Convention also adopted the Kyoto Protocol to the UNFCCC. The Kyoto Protocol legally binds developed countries to meet emission reduction targets. The Protocol's first commitment period began in 2008 and ended in 2012. The second commitment period began on 1 January 2013, and will end in 2020.

Today, there are 195 Parties to the Convention and 192 Parties to the Kyoto Protocol. The UNFCCC Secretariat supports all bodies involved in international climate change negotiations, especially the Conference of the Parties (COP), the Conference of the Parties serving as the Meeting of the Parties (CMP), the Subsidiary Bodies (SBSTA and SBI) which advise the COP and CMP, and the Bureau of the COP and CMP (which deals mainly with procedural and organizational issues arising from the COP and CMP, and also enjoys certain technical functions).

A brief description of how these UNFCCC bodies are interrelated is available at http://unfccc.int/bodies/items/6241.php.

At the same time, sustainable rural development and eradication of hunger, as specified in both UNCED and UNFCCC core principles, are at the basis of FAO mission. In particular, FAO generates knowledge and provides assistance to its member countries towards appropriate adaptation and mitigation responses in the agriculture, fisheries and forestry sectors, to ensure sustainability of production systems and maintain food security under climate change.

2.2 EXISTING REPORTING REQUIREMENTS FOR MITIGATION IN THE AGRICULTURE, FORESTRY AND OTHER LAND USE SECTOR UNDER THE UNFCCC

Parties to the UNFCCC have committed to implement specific actions and programs to mitigate climate change. In particular, UNFCCC Art. 4 states:

- "All Parties, taking into account their common but differentiated responsibilities and their specific national and regional development priorities, objectives and circumstances, shall:
- (a) (a) Develop, periodically update, publish and make available to the Conference of the Parties [...]national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol, using comparable methodologies to be agreed upon by the Conference of the Parties;
- (b) Formulate, implement, publish and regularly update national and, where appropriate, regional programmes containing measures to mitigate climate change by addressing anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol [...]
- (c) Communicate to the Conference of the Parties information related to implementation [...]."

From Art. 4 of the Convention, different reporting requirements have emerged for developed and developing countries (UNFCCC Annex I and non-Annex I Parties, respectively).

This note provides information on UNFCCC reporting requirements associated to mitigation of climate change in the AFOLU sector for developed and developing countries. Within the AFOLU sector, mitigation consists of actions taken to reduce GHG emissions and enhance carbon sinks and reservoirs, compared to a business-as-usual scenario/reference level. Parties need to report these actions periodically to the UNFCCC, as specified in a series of decisions agreed upon at the annual Conference of Parties (COP) and summarized in Tab. 1 in the Appendix. Specifically:

Annex I Parties report:

- National Communications (NC), containing information, from the last submitted GHG Inventory, on national GHG emissions/removals, climate-related policies and measures, GHG projections, vulnerability and adaptation to climate change, financial assistance and technology transfer to non-Annex I Parties, and actions to raise public awareness on climate change;
- National GHG Inventories (NGHGI), containing information on GHG emissions and removals, such as activity data, emission factors, and methodologies used to estimate these emissions. A NGHGI is composed of two distinct documents: the Common Reporting Format (CRF) tables, which contain a time-series of GHG emission estimates (from 1990 till the year x-2; where x is the submission year of the NGHGI); and the National Inventory Report (NIR), which includes all information on background data and methods used, and the data analysis and institutional arrangements underlying the preparation of the NGHGI.
- Biennial Reports (BRs), which outline progress in achieving net emissions reductions and provision of financial, technological, and capacity-building support to non-Annex I Parties for dealing with climate change.

National communications are to be submitted by Annex I Parties every 4 years (decisions 8/CP.1, 11/CP.4). They are prepared and reported periodically by Annex I Parties based on agreed reporting guidelines (decision 4/CP.5), and, for the GHG estimates, on methodology developed by the Intergovernmental Panel on Climate Change (IPCC), as adopted for the NGHGI by the COP. Submissions by Annex I Parties can be found here: http://unfccc.int/national_reports/annex_i_natcom/submitted_natcom/items/4903.php

National GHG inventories are submitted (3/CP.5) by Annex I Parties annually. They are prepared on the basis of reporting guidelines agreed by the COP (24/CP.19) and on methodologies developed by IPCC. National inventory arrangements should be in place to ensure that a NGHGI is fully compliant with reporting requirements and is

submitted on time. Submissions by Annex I Parties can be found here: http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/8108.php

Biennial Reports are to be submitted (2/CP.17) by developed country Parties every 2 years; the first biennial report (BR1) is due by 1 January 2014. They are prepared on the basis of agreed reporting guidelines (2/CP.17 Annex I) and on methodologies developed by IPCC, as per NGHGI. Submissions by Parties can be found here: http://unfccc.int/national_reports/biennial_reports_and_iar/submitted_biennial_reports/items/7550.php

Each report is subject to a review process assisted by the UNFCCC Secretariat and implemented by experts taken from the UNFCCC Roaster of Experts (RoE).

Non-Annex I Parties report:

- National Communications (NC), containing information on national circumstances, national GHG emissions/ removals, steps taken or envisaged to implement the Convention, and any other information considered relevant to the achievement of the objective of the Convention including, if feasible, material relevant to calculations of global emissions and emission trends;
- Biennial Update Reports (BURs), containing updated information on national circumstances and institutional arrangements for reporting on a continuous basis, national GHG emissions/removals information, including a national inventory report, and information on mitigation actions, effects, needs, and support received.

National communications may be submitted (decision 10/CP.2) by non-Annex I Parties every 4 years following decisions for each submission taken by the Conference of the Parties (COP). They are prepared and reported periodically by non-Annex I Parties based on agreed reporting guidelines (decision 17/CP.8) based on methodologies developed by the IPCC and adopted by the COP. Submissions by non-Annex I can be found here: http://unfccc.int/national_reports/non-annex_i_natcom/items/2979.php

Biennial Update Reports are to be submitted (2/CP.17) by non-Annex I Parties every 2 years, and are prepared on the basis of agreed reporting guidelines (decision 2/CP.17) based on methodologies developed by the IPCC and adopted by the COP. Least developed country Parties and small island developing States may submit biennial update reports at their discretion. The first biennial report (BUR1) is due by December 2014 and it is expected to contain information on current levels and trends of GHG emissions and removals within their territories.


The Biennial Update Reports will be subject to a technical assessment as part of the International Consultation and Analysis process, which is aimed at increasing the transparency of mitigation actions and their effects.

For activities (commonly referred ad REDD+) listed in Decision 1/CP.16, paragraph 70:

- Information, on the Forest Reference Emissions Levels and/or Forest Reference Levels, is prepared on the basis of agreed reporting guidelines (Decision 12/CP.17) and methodologies developed by the IPCC. The information is subject to assessment;
- Information on safeguard is to be reported;
- For receiving payments for results-based actions, information on forest-related emissions by sources and removals by sinks resulting from the implementation of the activities is prepared on the basis of agreed reporting guidelines (Decision 14/CP.19) and methodologies developed by the IPCC and is to be reported, in an annex to the BUR. The information is assessed, as part of the ICA process.

Detailed information on all references made above is available in Annex 4.

Figure 1 below illustrates the processes linked to national GHG reporting.



2.3 NATIONAL GHG INVENTORY SYSTEMS AND THEIR ROLE IN INTERNATIONAL PROCESSES

The core elements of information communicated by both Annex I and non-Annex I countries party to the Convention, pursuant to the UNFCCC, are those on GHG emissions and removals of greenhouse gases (GHG inventories), and the activities that Parties have undertaken to implement their contributions to the Convention's ultimate goals, i.e. their Policies and Measures for Mitigation and for Adaptation.

To collect, analyze and report this information, each country must have a national data system in place. This system can be defined as a set of institutional arrangements among all bodies involved in ensuring the collection of adequate data within established time-schedules, their analysis (including quality assurance), the compilation of data into estimates of GHG emissions and removals (including quality checks) and the quality assurance of those estimates.

All institutional arrangements should identify the relevant responsible body for the task at hand (e.g. the forestry service), and contain information on the task (e.g. collecting data on forests' carbon stocks) including data quality requirements and timing, on the resources available, and finally on the entity or entities to which the body must report (e.g. the GHG inventory unit entrusted with compiling GHG estimates for forest land).

Further information is available from the Handbook on National Greenhouse Gas Inventory Management Systems, authored by the Consultative Group of Experts on National Communications from Parties not included in Annex I to the Convention.

The Intergovernmental Panel on Climate Change

The Intergovernmental Panel on Climate Change (IPCC) is the leading international body for the assessment of climate change. It was established in 1988 by the United Nations Environment Programme (UNEP) and the World Meteorological Organization (WMO), to provide the world with a clear scientific view on the current state of knowledge on climate change, and its potential environmental and socio-economic impacts. In the same year, the UN General Assembly endorsed the WMO's and UNEP's joint establishment of the IPCC.

One of the IPCC's main activities is the preparation of comprehensive Assessment Reports on the state of scientific, technical and socio-economic knowledge on climate change, its causes, potential impacts and response strategies.

Since 1992, the IPCC has prepared methodologies and guidelines (IPCC National Greenhouse Gas Inventories Programme) to assist the Parties to the United Nations Framework Convention on Climate Change (UNFCCC) and its Kyoto Protocol in preparing national inventories of GHG emissions by source and removals by sinks. The latest major publication is the 2006 IPCC Guidelines for National Greenhouse Gas Inventories.

The IPCC's Task Force on National Greenhouse Gas Inventories (TFI), based in Japan, is the executive body for the implementation of the IPCC National Greenhouse Gas Inventories Programme.

The IPCC Guidelines for National GHG Inventories

3.1 THE STRUCTURE OF IPCC GUIDELINES

The 2006 IPCC Guidelines for National Greenhouse Gas Inventories (2006 IPCC Guidelines) were prepared upon the invitation of the UNFCCC, to provide good practice methodologies for countries to use in preparing GHG inventories when reporting to the UNFCCC. As shown in the picture below, the GHG inventory sectors are: Energy, Industrial Processes & Product Uses (IPPU), Agriculture, Forestry & Other Land Use (AFOLU) and Waste.


The 2006 IPCC Guidelines are an evolutionary development with respect to the 1996 IPCC Guidelines, the GPG 2000 and the GPG-LULUCF 2003. The 2006 approach ensures continuity and enables experiences with the existing guidelines, new scientific information, and the results of the UNFCCC review process to be incorporated.

One of the most significant changes between the 1996 and 2006 versions of the Guidelines is the assembly of Land Use, Land Use Change and Forestry (LULUCF) and the Agriculture sectors into a single AFOLU sector.

The categories from both sectors (LULUCF and Agriculture) were integrated into this new framework, to resolve inconsistencies and avoid double counting. In addition, this enabled:

- Removal of the somewhat arbitrary distinction between these categories in the previous versions of the guidance, and promotion of consistent use of data between them, especially for more detailed methods;
- Consistency in the treatment of gases in the Agriculture and LULUCF Sectors, thus enabling a more consistent treatment of land conversions.





This integration recognizes that the processes underlying GHG emissions and removals, as well as the different forms of terrestrial carbon stocks, can occur across all types of land, and is intended to improve consistency and completeness in estimating and reporting GHG emissions and removals.

Figure 3 below (IPCC 2006) shows the evolution from the 1996 to the 2006 versions of the IPCC guidelines, having regard to the Agriculture/LUCF, Agriculture/LULUCF and AFOLU sectors.

Evolution of IPCC Guidelines for National GHG Emission Inventories 1996 IPCC GLs GPG & GPG-LULUCF 2006 IPCC Guidelines Agriculture and Land Use Agriculture and Land Use, Agriculture and Land Use, Change and Forestry Land-Use Change and Land-use Change and (LUCF) separate sectors. Forestry (LULUCF) separate Forestry (LUCF) combined into a single sector sectors. Agriculture, Forestry and Only the most important Other Land Use (AFOLU). activities resulting in GHG Provides good practice and emissions/removals. uncertainty management Same approach as GPGguidance. LULUCF. Retained use of Managed Implicit assumption about estimating emissions and Includes all land use Land. removals only over lands emissions/removals split subject to human into six land-use categories Inclusion and consolidation of intervention. from all pools. several previously optional categories. Only accounted for Explicit Use of managed above-ground biomass land as a proxy for Refinement of methods and and soil C pools anthropogenic emissions/ improved defaults removals

3.2 THE QUALITY OF THE EMISSION INVENTORY

FIGURE 3


The IPCC Guidelines reflect the international data quality requirements defined by the UNI ISO 14064 (http://www.iso.org/iso/catalogue_detail?csnumber=38381), in order to build inventories that are consistent, comparable, complete, accurate and transparent; this, in turn, enables inventory quality to be improved over time (IPCC-2006, Chapter 1.4).

The 2006 Guidelines encourage the continuous improvement of emission inventory compilation, by adopting the principles required by international standards during all steps of the process and by recommending regular communication and consultation with data providers throughout all the inventory activities (from data collection to final reporting).

The inventory brings benefits in terms of both efficiency and quality, through the relationship between data suppliers and inventory compilers. This ongoing communication and consultation will also enable the inventory compilers to be promptly informed of any new datasets being developed, and also provide opportunities to influence the planning and specifications of data providers' data collection activities.

In each phase of the GHGI (shown below, Figure 4), the inventory compiler should verify the quality of the data (absence of bias and magnitude of uncertainties) the appropriateness of the data for the methodology applied (consistency of definitions and timeliness), the suitability of the methodology selected for the data available, the improvement in estimate accuracy and data collection quality (accuracy in sampling design and measurements, increase of time frequency).

As mentioned, Figure 4 below summarises the GHGI's main steps:



- **Data collection.** This activity covers the evaluation of existing sources of data, and the planning of new emission measurements and surveys; extensive reference is made to the information provided by other organisations. The IPCC guidelines provide information on initiating and maintaining a data collection program, on the data collection process and on other general issues.
- Uncertainty analysis. The IPCC guidelines provide information on estimating and combining uncertainties, along with a discussion of the conceptual underpinnings of inventory uncertainty, and the uncertainty issues related to specific categories of emissions and removals. The uncertainty analysis characterises the range and likelihood of possible values for the national inventory, as a whole and for its individual components. An awareness of the uncertainty of parameters and results provides inventory compilers with insight when evaluating suitable data for the inventory, during the data collection and compilation phases.
- Methodological choice and key category analysis. IPCC guidelines provide good practice guidance on how to
 identify key categories of emissions and removals. This phase includes the concept of the key category together
 with decision trees, to guide users in choosing the methodology for each category.
- Time-series consistency. This activity provides methods for ensuring time-series consistency, in cases where it is not possible to use the same method and/or data over the entire period. The IPCC guidelines provide good practice guidance on when to recalculate estimates for previous years, and methods for accounting for changes in emissions and removals over time.


- Quality Assurance (QA) and Quality Control (QC) and Verification. The IPCC Guidelines describe the
 general QA/QC aspects to be considered when compiling an inventory of emissions and removals, including
 good practice guidance on sector-specific quality control checks. Techniques for verifying inventories using
 external data are also outlined.
- Reporting. The IPCC Guidelines contain reporting guidance and tables, and specifically address issues related
 to reporting, including the following: reporting coverage, in terms of sectors, categories, national territory, gases,
 pools; its time-frame; the use of notation keys, units and digits. This aims to ensure completeness, promote
 transparency and establish comparability.

3.3 THE AFOLU SECTOR AND ITS STRUCTURE

The Agriculture, Forestry and Other Land Use (AFOLU) sector deals with anthropogenic GHG emissions and removals, defined as all emissions and removals occurring on 'managed land' and that are associated with the use of land, including agriculture and husbandry. Managed land is land where human interventions and practices have been applied to perform production, ecological or social functions (IPCC, 2006).

Figure 5 below provides a synthetic image on how land use and management can influence a variety of ecosystem processes, which in turn can affect greenhouse gas fluxes such as photosynthesis, respiration, decomposition, nitrification/denitrification, enteric fermentation, and combustion.

These processes involve transformations of carbon and nitrogen, driven by biological (activity of microorganisms, plants, and animals) and physical processes (combustion, leaching, and run-off).



Source: (IPCC, 2006).

The estimates of GHG emissions and removals deriving from AFOLU include:

- CO₂ emissions and removals resulting from C stock changes in biomass, dead organic matter (DOM), soil organic matter (SOM) of organic and mineral soils, and harvested woody products (HWP) for all managed lands;
- CO₂ from cultivated organic soils;
- non-CO₂ emissions from fire on all managed land;
- CH₄ emissions from rice cultivation;
- N₂O emissions from all managed soils;
- CO₂ emissions associated with liming and urea application to managed soils;
- CH₄ emissions from livestock enteric fermentation;
- CH₄ and N₂O emissions from manure management systems.

Greenhouse gas fluxes can be estimated in two ways:

- as net changes in C stocks over time (used for most CO₂ fluxes) and
- directly, as gas flux rates to and from the atmosphere (used for estimating non-CO₂ emissions and some CO₂ emissions).

The use of C stock changes to estimate CO_2 emissions and removals is based on the fact that changes in ecosystem C stocks occur predominantly (but not exclusively) through CO_2 exchange between the land surface and the atmosphere (i.e. other C transfer process such as leaching are assumed to be negligible). Hence, increases in total C stocks over time are equated with a net removal of CO_2 from the atmosphere, and decreases in total C stocks (less transfers to other pools such as harvested wood products) are equated with a net emission of CO_2 . Non- CO_2 emissions are largely a product of microbiological processes (i.e. within soils, animal digestive tracts and manure) and of the combustion of organic materials.

The emission and removal processes are organized by the ecosystem components:

- **Biomass**: plant biomass, including above-ground and below-ground parts, is the main conduit for CO₂ removal from the atmosphere.
- **Dead organic matter:** the bulk of biomass production contained in living plant material, which is eventually transferred to dead organic matter (DOM) pools (i.e. dead wood and litter).
- **Soils:** as dead organic matter is fragmented and decomposed, it is transformed into soil organic matter. Additionally, soils can be fertilized and amended.
- **Livestock**: animal production systems, particularly those featuring ruminant animals, can be significant sources of greenhouse gas emissions.

AFOLU emission sources/sinks are categorized by:

1. Land use. Each land use is subdivided into land remaining in the same category (e.g. Forest land Remaining forest land) and land converted from one category to another (e.g. Forest land converted to cropland). Countries may choose to further stratify land in each category by climatic or other ecological regions, depending on the choice of the method and its requirements. Greenhouse gas emissions and CO₂ removals determined for each specific land use include CO₂ (as carbon stock changes) from biomass, dead organic matter, soils and HWP, if any, as well as non-CO₂ emissions from burning.


Land use comprises the following categories:

- **Forest land**. This category includes all land with woody vegetation consistent with the thresholds used to define forest land in the national greenhouse gas inventory. It also includes systems with a vegetation structure that currently fall below, but that *in situ* could potentially reach, the threshold values used by a country to define the forest land category.
- **Cropland**. This category includes cropped land, including rice fields, and agro-forestry systems where the vegetation structure falls below the thresholds used for the forest land category.

- **Grassland**. This category includes rangelands and pasture land that are not considered Cropland. It also includes systems with woody vegetation and other non-grass vegetation, such as herbs and brushes that fall below the threshold values used in the forest land category. The category also includes all grassland, from wild lands to recreational areas as well as agricultural and silvo-pastoral systems, consistent with national definitions.
- Wetlands. This category includes areas of peat extraction and land that is covered or saturated by water for all
 or part of the year (e.g. peatlands), and that do not fall within the forest land, cropland, grassland or settlements
 categories. It includes reservoirs as a managed sub-division, and natural rivers and lakes as unmanaged subdivisions.
- **Settlements**. This category includes all developed land, including transportation infrastructure and human settlements of any size, unless they are already included under other categories. The elements falling within this category should be consistent with national definitions.
- Other land. This category includes bare soil, rock, ice, and all land areas that do not fall into any of the other five categories. It enables the total of identified land areas to match the national area, where data are available. If data are available, countries are encouraged to classify unmanaged lands by the above land-use categories (e.g. Unmanaged forest land, unmanaged grassland, and unmanaged wetlands).
- 2. Agricultural practices include the following categories:
- Enteric fermentation. This is the fermentation process that produces methane as a by-product of the normal livestock digestive process, in which microbes resident in the animal's digestive system ferment the feed consumed by the animal.
- Manure management. Livestock manure is primarily composed of organic material and water. Under
 anaerobic conditions, the organic material is decomposed by anaerobic bacteria. The end products of anaerobic
 decomposition are methane, carbon dioxide, and stabilized organic material. The methane production potential
 of manure depends on the manure's specific composition, which in turn depends on the composition and
 digestibility of the animal diet.
- **Biomass burning.** It includes only living biomass. Although, the fuel components other than live biomass are often very significant, especially in peatlands (i.e. DOM).
- Managed soils. This category includes all agricultural soils.
- This category includes direct and indirect nitrous oxide emissions that are usually estimated from data on nitrogen supplied to soils, including nitrogen fertiliser usage or sales, crops residue management, organic amendments, cultivation of organic soils (i.e. drainage of peatlands in agricultural land) and land-use conversions that enhance mineralisation of nitrogen in soil organic matter.
- Liming is used to reduce soil acidity and improve plant growth in managed systems, particularly agricultural lands and managed forests. Adding carbonates to soils in the form of lime (e.g., calcic limestone (CaCO₃), or dolomite (CaMg(CO₃)₂) leads to CO₂ emissions as the carbonate limes dissolve and release bicarbonate (2HCO₃), which evolves into CO₂ and water (H₂O). These are methods for soil fertilization. Adding urea to soils during fertilisation leads to a loss of the CO₂ that was fixed in the industrial production process. Urea (CO(NH₂)₂) is converted into ammonium (NH₄⁺), a hydroxyl ion (OH⁻), and bicarbonate (HCO₃⁻), in the presence of water and urease enzymes. Similar to the soil reaction following addition of lime, the bicarbonate that is formed evolves into CO₂ and water. This source category is included because the CO₂ removal from the atmosphere during urea manufacturing is estimated in the Industrial Processes and Product Use Sector.
- **Rice cultivation.** This category refers to the anaerobic decomposition of organic material in flooded rice fields that produces methane, which escapes to the atmosphere primarily through air-bubbles and by being transported through the rice plants. The amount emitted is a function of the rice species, the number and duration of harvests, the soil type and temperature, the irrigation method, and fertilizer use.

This structure has undergone some changes over time to meet user needs, as reflected in the IPCC Guidelines of 1996 – 2006.

Figure 6 below shows the mapping of the emission category structures of AFOLU (from IPCC-2006), LULUCF from the GPG for LULUCF 2003 and 1996, and LUCF from the Revised Guidelines 1996.



Different sectoral mapping

- "Enteric fermentation: Poultry" is now reported under "other";
- "Manure management" is now reported by animal type only;
- "Agricultural soils" and "CO₂ emissions and removals from soils" have been moved from managed soils to 3G liming, 3H urea application, 3D direct N₂O emissions.

Fires, previously under "Forest and grassland conversion", "Field burning of agricultural residues" and "Prescribed burning of savannas", are re-classified under biomass burning.

New categories

The following new categories have been introduced:

- 3D indirect N₂O emissions from the atmospheric deposition of nitrogen in NO_x and NH₃;
- 3H urea application;
- 3D Harvested wood products.


Table 1 shows the differences in Categorization of Sources and Sinks of UNFCCC reporting tables related to the Revised 1996 IPCC Guidelines, the Good Practice Guidance and the 2006 IPCC Guidelines.

TABLE 1
Categorization of sources and sinks of IPCC Guidelines reporting tables, 1996-2006

Revised 1996 IPCC Guidelines			IPCC GPG 2000	2006 IPCC Guidelines (AFOLU	
4	AGRICULTURE	4	AGRICULTURE	3	AGRICULTURE
4.A	Enteric fermentation	4.A	Enteric fermentation	3.A	Enteric fermentation
4.B	Manure management	4.B	Manure management	3.B	Manure management
4.C	Rice cultivations	4.C	Rice cultivations	3.C	Rice cultivations
4.D	Agricultural soils	4.D	Agricultural soils	3.D	Agricultural soils
4.E	Prescribed burning of savannas	4.E	Prescribed burning of savannas	3.E	Prescribed burning of savannas
4.F	Field burning of agricultural residues	4.F	Field burning of agricultural residues	3.F	Field burning of agricultural residues
				3.G	Liming
				3.H	Urea application
				3.1	Other carbon-containing fertilizers
4.G	Other	4.G	Other	3.J	Other
			IPCC GPG 2003		
5	LAND-USE CHANGE & FORESTRY	5	LULUCF	4	LULUCF
5.A	Change in forest and other woody biomass stocks	5.A	Forest land	4.A	Forest land
5.B	Manure management	5.B	Cropland	4.B	Cropland
5.C	Rice cultivations	5.C	Grassland	4.C	Grassland
5.D	Agricultural soils	5.D	Wetlands	4.D	Wetlands
		5.E	Settlements	4.E	Settlements
		5.F	Other land	4.F	Other land
	Other (harvested wood products)	5.G	Other (harvested wood products)	4.G	Other (harvested wood products)

3.4 REPORTING METHODS

The fundamental formula for estimating the quantity of GHG emissions can always be expressed as the multiplication of the activity data (AD) by the emission factor (EF), as follows:



Where:

• EF = Emission factor

Emission factors are coefficients that quantify the emissions or removals of a gas per unit activity data. Emission factors are based on samples of measurements, averaged at various levels of detail depending upon the Tier methodology used, to develop a representative rate of emission for a given activity level under a given set of operating conditions.

AD = Activity data

Activity data describe the magnitude of a human activity resulting in emissions or removals of greenhouse gases, taking place during a given period of time and over a specified area.


Data on livestock type and numbers, the area extent of managed agriculture, pastures, forests and associated changes or the amount of synthetic or organic fertilizer applied, are all examples of AD relevant to the computation of emissions for the AFOLU sector.

In addition to the AD and the EF, this basic equation can also incorporate other estimation parameters to reflect actual emissions or removals.

The quantification of GHG emissions in an inventory is a multi-step process for each category (for Non-Annex I, a detailed step-by-step training is available at http://unfccc.int/resource/ cd roms/na1/ghg inventories/index.htm).

3.5 ESTIMATION PROCESS

The IPCC 2006 Guidelines provide comprehensive documentation on how to proceed for estimating the emission and removals, as shown in the Inventory development cycle below (IPCC, 2006):



Below the initial steps of the process are described; these steps are also analysed and reported in the guidelines for each gas category.

3.5.1 The identification of key categories

A key category is one that is prioritised within the national inventory system, because its estimate has a significant influence on a country's total inventory of greenhouse gases in terms of the absolute level, the trend, or the uncertainty in emissions and removals.

Whenever the term "key category" is used, it includes both source and sinks categories. Key categories should be the priority for countries when allocating inventory resources for data collection, compilation, quality assurance/ quality control and reporting. The main purposes are:

- To focus available resources for improving data and methods upon categories identified as key;
- To analyse what level of methodological complexity can be applied (e.g. Tier 2 and 3 methods);
- To give further attention to key categories, in terms of quality assurance and quality control.

Any inventory compiler who has experience in preparing a national GHG inventory will be able to identify key categories, in terms of their contribution to the absolute level of national emissions and removals. For those inventory compilers who have prepared a time series, the quantitative determination of key categories should include an evaluation of both the absolute level and the trend of emissions and removals. Some key categories may be identified only when their influence on the trend of the national inventory is taken into account.

Two approaches for performing the key category analysis have been developed. Both approaches identify key categories in terms of their contribution to the absolute level of national emissions and removals and to the trend of emissions and removals.

- In Approach 1, key categories are identified using a pre-determined cumulative emissions threshold. Key
 categories are those that, when summed together in descending order of magnitude, add up to 95 percent of the
 total level.
- Approach 2 can be used by inventory compilers, if category uncertainties or parameter uncertainties exist. Under Approach 2, categories are sorted according to their contribution to this uncertainty.

The results of Approach 2 are additional to those that may be obtained from Approach 1.

For each key category, where relevant, the inventory compiler should determine whether certain subcategories are particularly significant. Usually, for this purpose, the subcategories should be ranked according to their contribution to the aggregate key category. Those subcategories that, considered together, contribute over 60 percent to the key category should be treated as being particularly significant. It may be appropriate to focus efforts towards making methodological improvements with regard to these most significant subcategories.

Having regard to the key categories for AFOLU, it is necessary to identify the following elements:

- Which CO₂ net emissions or removals, from which land-use categories and management activities, are significant;
- Which CH₄ or N₂O emissions, from which land-use and management activities, are significant;
- Within categories, which (sub)category and/or which C pool is significant;
- In accordance with the key category analysis, the IPCC determines which tier level methodology should be applied when preparing the estimate.

3.5.2 The selection of methods and measurement

The IPCC-2006 Guidelines report three tiered approaches related to methods used in the AFOLU Sector:

- Tier 1 is the basic method;
- Tier 2 is the intermediate method; and
- Tier 3 is the most demanding, in terms of complexity and data requirements.

Generally, moving to higher tiers improves the inventory's accuracy and reduces uncertainty, but the complexity and resources required for conducting inventories also increase. Tiers 2 and 3 are sometimes referred to as higher-tier methods, and are generally considered to be more accurate.

If necessary, a combination of tiers can be used, e.g. Tier 2 can be used for biomass and Tier 1 for soil carbon. The methods and data presented focus on Tier 1 inventories. The methods will be generally applicable to Tier 2 inventories, but the default data presented for Tier 1 will be partly or wholly replaced with national data as part of the Tier 2 estimation. Tier 3 methods are not described in detail, but good practices in their application are outlined.

The framework of the Tier structure for AFOLU methods is as follows:

- **Tier 1** methods are designed to be the simplest to use; equations and default parameter values (e.g., emission and stock change factors) for these methods are provided by IPCC guidelines.

Country-specific activity data are necessary, but for Tier 1 sources of activity data estimates are often globally available (e.g. deforestation rates, agricultural production statistics, global land cover maps, fertilizer use, livestock population data, etc.), although these data are usually spatially coarse. It is good practice to use data from official international sources when national data are lacking.

Tier 2 can use the same methodological approach as Tier 1, but applies emission and stock change factors that are based on country- or region-specific data, for the most important land-use or livestock categories.

Country-defined emission factors are more appropriate for the climatic regions, land-use systems and livestock categories in that country. Higher temporal and spatial resolution and more disaggregated activity data are typically used in Tier 2 to correspond with country-defined coefficients for specific regions and specialized land-use or livestock categories.


In **Tier 3**, higher order methods are used, including models and inventory measurement systems tailored to national circumstances, repeated over time, and driven by high-resolution activity data and disaggregated at the sub-national level.

These higher order methods provide estimates of greater certainty than lower tiers. Higher order systems may include comprehensive field sampling repeated at regular time intervals and/or GIS-based systems of age, class/production data, soils data, and land-use and management activity data, integrating several types of monitoring. Pieces of land where a land-use change occurs can usually be tracked over time, at least statistically. In most cases, these systems present a climate dependency, and thus provide source estimates with inter-annual variability. Detailed dis-aggregation of livestock population according to animal type, age, body weight etc., can be used. Models should undergo quality checks, audits, and validations, and be thoroughly documented.

To facilitate analysis of the methods, the Guidelines propose using a "decision tree" model, which assists the inventory compiler to navigate the guidance and select the appropriate tiered methodology for their circumstances, based on their assessment of the key categories.

Figure 9 below shows an example of methodological tiers decision tree, for "livestock categories" (Source IPCC, 2006).

FIGURE 9
Decision tree for the livestock category



This should be repeated for each livestock species and gas. It is important to note, however, that many countries often do not have access to Tier 2 data even for key categories. This manual guides them, even in these cases, on how to build a Tier 1 inventory, which can then be used as a solid, sustainable base for further refinement once human and financial resources become available.

3.5.3 The selection and collection of activity data

Data collection is an integral part of developing and updating a GHG inventory. Formalised data collection activities should be established, adapted to countries' national circumstances, and reviewed periodically as a part of implementing good practices.

Data sources:

• National and International Literature

The IPCC Guidelines recommend collecting data from bodies such as National Statistical Agencies, and national

regulatory authorities responsible for permitting of industrial and other processes subject to pollution emission legislation.

There will be occasions, however, when other sources of specialised literature provide activity data, e.g. UN statistics (including the FAOSTAT database: http://faostat3.fao.org/faostat-gateway/go/to/home/E), the US Geological Survey (USGS) reports on commodities, and technical reports, guidelines, standards, sectoral surveys issued by industrial trade associations.

Surveys & Census information

Survey and census information provide the best agricultural, production and energy statistics that can be used for greenhouse gas inventories. Generally, these data are compiled by national statistical agencies (NSAs) or relevant ministries, for national policy purposes or to comply with international demand for data, or other activities that are beyond the direct control of the inventory compiler. However, the needs of the inventory may sometimes trigger or influence surveys or censuses.

More detailed information is available at the section on Data Collection of the IPCC guidelines (IPCC 2006, Vol.1 – CH. 2.2.5).

3.5.4 The selection of emission factors or carbon stock change factors

The IPCC Guidelines define the emission factor as the average emission rate of a given GHG for a given source, relative to units of activity.

The 2006 IPCC Guidelines, adopting the Tier 1 methods for all categories, enable the use of readily available national or international statistics, in combination with provided default emission factors and additional parameters provided, thus making it feasible to prepare estimates for all countries.

A practical example could be the calculation of methane emissions from enteric fermentation.

Methane is produced by the fermentation of feed within an animal's digestive system. Generally, a higher feed intake produces higher methane emission. Although the extent of methane production is affected by the composition of the animal's diet, feed intake is positively related to animal size, growth rate and production (e.g. milk production, wool growth, or pregnancy). All these variables assume a range of values in accordance with the husbandry characteristics, which in turn change according to the region.

Selecting the Tier 1 method, the 2006 IPCC provides the default emission factors that are more appropriate for the country's livestock characteristics. These have been drawn from previous studies, and are organized by region for ease of use.

The Table 2 below shows the emission factors of enteric fermentation by the cattle sub-category, and by regional area (Table 10.11 of the 2006 IPCC, Vol4, Ch10).

TABLE 2 Tier 1 CH_4 emission factors from enteric fermentation for cattle

Regional characteristics	Cattle category	Emission factor (kg CH ₄ ead ⁻¹ y ^{r-1})	Comments
North America: Highly productive commercialized dairy sector feeding high quality forage and grain.	Dairy	128	Average milk production of 8,400 kg head ⁻¹ yr ⁻¹ .
Separate beef cow herd, primarily grazing with feed supplements seasonally. Fast-growing beef steers/ heifers finished in feedlots on grain. Dairy cows are a small part of the population.	Other cattle	53	Includes beef cows, bulls, calves, growing steers/heifers, and feedlot cattle.
Western Europe: Highly productive commercialised dairy sector feeding high quality forage and grain. Dairy cows also used for beef calf production. Very	Dairy	117	Average milk production of $6,000 \text{ kg head}^{-1} \text{ yr}^{-1}$.
small dedicated beef cow herd. Minor amount of feedlot feeding with grains.	Other cattle	57	Includes bulls, calves, and growing steers/heifers.
Eastern Europe: Commercialised dairy sector feeding	Dairy	99	Average milk production of 2,550 kg head ⁻¹ yr ⁻¹ .
mostly forages. Separate beef cow herd, primarily grazing. Minor amount of feedlot feeding with grains.	Other cattle	58	Includes beef cows, bulls, and young.
Oceania: Commercialised dairy sector based on grazing. Separate beef cow herd, primarily grazing rangelands	Dairy	90	Average milk production of 2,200 kg head ⁻¹ yr ⁻¹ .
of widely varying quality. Growing amount of feedlot feeding with grains. Dairy cows are a small part of the population.	Other cattle	60	Includes beef cows, bulls, and young.
Latin America: Commercialised dairy sector based on grazing. Separate beef cow herd grazing pastures	Dairy	72	Average milk production of 800 kg head ⁻¹ yr ⁻¹ .
and rangelands. Minor amount of feedlot feeding with grains. Growing non-dairy cattle comprise a large portion of the population.	Other cattle	56	Includes beef cows, bulls, and young.
Asia: Small commercialised dairy sector. Most cattle are multi-purpose, providing draft power and some milk	Dairy	68	Average milk production of 1,650 kg head ⁻¹ yr ¹ .
within farming regions. Small grazing population. Cattle of all types are smaller than those found in most other regions.	Other cattle	47	Includes multi-purpose cows, bulls, and young
Africa and Middle East: Commercialised dairy sector based on grazing with low production per cow.	Dairy	46	Average milk production of 475 kg head ⁻¹ yr ⁻¹ .
Most cattle are multi-purpose, providing draft power and some milk within farming regions. Some cattle graze over very large areas. Cattle are smaller than those found in most other regions.	Other Cattle	31	Includes multi-purpose cows, bulls, and young
Indian Subcontinent: Commercialised dairy sector based on crop by-product feeding with low	Dairy	58	Average milk production of 900 kg head ⁻¹ yr ⁻¹ .
production per cow. Most bullocks provide draft power and cows provide some milk in farming regions. Small grazing population. Cattle in this region are the smallest compared to cattle found in all other regions.	Other Cattle	27	Includes cows, bulls, and young. Young comprise a large portion of the population

Addressing data and methodological requirements

The key gaps and needs for improving statistical data and apply methodologies for estimation of national GHG inventories, which are routinely communicated by practitioners in member countries, are institutional gaps and a lack of basic data. Institutional gaps refer to the need to improve coordination across and within relevant national agencies involved in statistical data compilation and analysis on the one hand, and GHG data estimation on the other, with the objective of ensuring efficient production, inclusion in national statistical systems, and long-term sustainability of databases. Lack of basic data is a well-recognized gap especially for agriculture, forestry and land use data, being the focus of several ongoing international capacity development programs, including the *Global Strategy to Improve Agricultural and Rural Statistics*, funded by international and national agencies alike.

While the two gaps mentioned above are typical of many statistical domains, those linked to GHG reporting processes present a unique feature, one that opens opportunities for progress through practical steps and useful cross-linkages. The unique feature lies in the 'legal' nature of GHG inventories for developed countries, so-called Annex I parties, who submit annually their GHG inventories for reporting as per international climate policy agreements under the UNFCCC. For those Annex I parties that are also signatories to the Kyoto Protocol, national GHG inventories are the *de facto* Monitoring, Verification and Reporting (MRV) international instrument used to assess national performance against country-specific GHG reduction targets.

Even for developing countries with no GHG emission reduction targets, a robust and complete GHG Inventory system—allowing for analysis of time-series and extrapolation of trends, as well as identification of hotspots and of mitigation options— allows all member countries to communicate transparently their intended national mitigation actions to the international community, thus representing a mean for accessing international funding mechanisms such as REDD+ and NAMAs.

As discussed in previous chapters, each member country builds its own GHG inventory following the relevant IPCC guidelines, in line with its national capacity, including the possibility to utilize sub-national activity data and refined emission factors where possible. It is nonetheless useful to also build, at the outset, a simplified, reference GHG Inventory based on official national data and IPCC Tier 1 methodologies, to be used by national practitioners as: i) a starting point whenever national capacity for higher Tiers is lacking; ii) to determine national data gaps and perform QA/QC analyses; and iii) to evaluate specific opportunities and costs for inventory refinement at higher Tiers. A simplified reference inventory is furthermore useful to compare data from different countries within a coherent data framework.

And yet, many countries to date lack a reference, Tier 1 GHG inventory, while lack of more advanced national data and methods continues to hinder completion of GHG inventories at higher Tiers, especially for agriculture, forestry and land use.

This Manual shows which data, emission factors and computation procedures are necessary and sufficient to build a simplified yet robust GHG national Inventory at Tier 1. In order to provide practical steps to the reader, reference is regularly made to the FAOSTAT Emissions database, which provides a complete set of activity data and emission estimates using the IPCC 2006 Guidelines, covering all member countries over the period 1961-present, and for all relevant agriculture, forestry and land use sectors.

4.1 ACTIVITY DATA SOURCES

Official national activity data (AD) needed to build a Tier 1 inventory include those typically communicated by member countries to FAO and collected in the FAOSTAT (agriculture activities and land resources) and FRA (forest activities) databases. The FAO process for food and agricultural data received annually by countries includes collection, harmonization, gap filling in case of no communication, and statistical data validation by member countries and FAO according to international standards of the United Nations system. For more information on relevant FAO statistical processes, please see http://www.fao.org/statistics/en/ and http://www.fao.org/forestry/fra/en/.

Member countries do not report to FAO a complete set of national data needed to calculate all required GHG emissions, however. Additional data elements need to be computed with the help of geospatial databases and satellite remote-sensing information. Table 3 below summarises all the activity secondary data sources that can be used by national practitioners to calculate national GHG emissions for AFOLU, as available at FAO through FAOSTAT and including additional relevant databases.

TABLE 3 FAOSTAT emissions database sources of activity data

		CATEGORY	GAS	Activity Data Source	
	Enteric Fermentation CH ₄ Manure Management CH4,N ₂ O		7		
			CH4,N₂O		
	Rice Cultivation		CH₄	FAOSTAT Agriculture:	
		Synthetic Fertilizers	N₂O	National questionnaires	
	ils	Manure applied to soils	N₂O	Other International sources FAOSTAT Imputation methods	
TURE	ral so	Manure left on pasture	N₂O		
	重	Crop residues	N₂O		
AGRICULTURE	Agricultural soils	Cultivated organic soils	N₂O	Harmonized World Soil (HSWD) Global Land Cover 2000 (GLC 2000) Gridded livestock of the world (GLW)	
	Burning - Savanna		CH₄, N₂O	Global Fire Emission Database 4 (GFED 4) Climatic map (JRC) Global Ecological Zones (GEZ)	
	Bur	ning - Crop residues	N₂O	FAOSTAT	
	For	est land	CO ₂	Global Forest Resources Assessment (FRA)	
共	$ \begin{array}{c} \text{Cropland} & \text{CO}_2 \\ \\ \text{Grassland} & \text{CO}_2 \end{array} $				Harmonized World Soil (HSWD) Global Land Cover 2000 (GLC)
LULUCF			CO ₂	2000)	
	Bur	ning Biomass	CH₄, N₂O, CO₂	Global Fire Emission Database 4Climatic map (JRC)Global Ecological Zones (GEZ)	

These activity data sources are discussed in further detail below:

1) **FAOSTAT database.** As described in the previous chapter, the data derive from national questionnaires and pertain to various agricultural sectors (samples of these questionnaires may be viewed at http://www.fao.org/economic/ess/ess-home/questionnaires/en/). In particular, the questionnaires on production, fertilizers, land

use and irrigation collect basic data for the calculation of the GHG emissions. These are compiled by National Statistics Offices (NSOs) or Ministers of Agriculture, and sent to FAO Statistics Division on an annual basis. The questionnaire data are checked with countries and validated, so that coherent and consistent time series may be obtained (more information is available in Annex 2 to this publication). In case of gaps or missing official data, time series are completed with other international data sources and calculated through estimation methods. The data are disseminated online after a thorough evaluation process.

Detailed information on FAO concepts and the definitions and classifications adopted by FAOSTAT is available at the "Methods & Standards" section of the FAOSTAT website: http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E. Annex 3 to this Manual includes a Glossary that explains the concepts and definitions of selected indicators, and of items representing the GHG emissions' data activity. The correspondence between the IPCC classification and the FAOSTAT Commodity list of livestock items is reported in Table 60A.

2) Global Forestry Resources Assessments. These are the most comprehensive assessments of forests and forestry performed to date. For its design and implementation, FAO worked closely with countries and specialists, through regular contact and expert consultations, and organizing ten regional and sub-regional workshops and other training sessions for national correspondents.

Data is disseminated through periodic Forestry Resources Assessment (FRA) reports. Detailed information is available at http://www.fao.org/forestry/fra/fra2010/en/.

To date, FRA includes information collated from 233 countries and territories for four points in time: 1990, 2000, 2005 and 2010. It covers the extent, condition, uses and values of forests and other wooded land, with the aim of assessing all benefits deriving from forest resources.

3) **Geospatial data.** Data of emissions from burning activities and from cultivated organic soils need to be calculated using Geographic Information System (GIS) analysis and satellite data processing. Further details are provided in the metadata and calculation procedure sections.

Source data include, for instance:

- Harmonized World Soil (HWSD): http://www.fao.org/nr/lman/abst/lman_080701_en.htm
- Global Land Cover 2000 (GLC 2000), prepared by the European Commission Joint Research Centre (JRC): http://bioval.jrc.ec.europa.eu/products/glc2000/glc2000.php
- Global Fire Emission Database 4 (GFED4): http://www.globalfiredata.org/
- Climatic map (JRC) based on IPCC methodology: http://eusoils.jrc.ec.europa.eu/projects/Renewable Energy/
- Global Ecological Zones (GEZ): http://www.fao.org/docrep/017/ap861e/ap861e00.pdf

Figures 10-12 below represent, as a notable example, three fundamental steps that could be used in calculating GHG emissions from biomass burning.

Figure 10 shows an example of the total burned area for savannas in Africa in 2013. This is the first input dataset for retrieving final emissions.

FIGURE 10 Map of savanna burned areas in 2012




Figure 11 displays the biomass burned for savannas in 2012, calculated with parameters from the 2006 IPCC Guidelines.

FIGURE 11 Savanna dry biomass burned in 2012





Figure 12 is the final result of data processing, with N_2O gas emissions from savannas burned in Africa in 2012.

FIGURE 12 Map of $N_2\text{O}$ emissions from savanna burned in 2012



4.2 EMISSION FACTORS

The IPCC Guidelines provide a complete set of deafult emission factors (EFs) for agriculture, forestry and land use activities, often specified at regional and/or subcategory/subdivision level, as derived from a thorough literature review. As discussed in the Introduction, Ch. 5 provides a complete guide on how to assign specific EFs to each relevant category or sub-category of interest, with the provision of explanations and detailed tables by country. Each member country chooses the IPCC guidelines most appropriate to national circumstances, taking into consideration COP/MOP decisions taken under the UNFCCC processes and discussed in Ch. 2 above. For simplicity, this Manual only addresses the process of determining emission factors and emission estimations related to the application of the IPCC 2006 Guidelines. This is a useful exercise regardless of specific national choices of guidelines, considering that the IPCC 2006 guidelines, as discussed in Ch.3, are consistent with and in fact incorporate most of the Revised 1996 IPCC Guidelines, including GPG 2000 and GPG 2003. Below we provide a general view of the kind of variables that need to be covered to build a GHG inventory for agriculture, forestry and the land use sectors.

4.3 METADATA AND EMISSIONS CATEGORIES FOR AGRICULTURE

Agriculture contains all the emissions produced in the different agricultural sub-domains. We further provide guidance on estimating emissions from energy use in agriculture, consistently with information currently provided and compiled by FAO.

GHG emissions from the AFOLU sector consist of non-CO₂ gases, namely methane (CH₄) and nitrous oxide (N₂O), generated through crop and livestock production and management activities. Emissions from energy used in agriculture consist mainly of CO₂ and, to a lesser extent, of CH₄ and N₂O. All emissions may be expressed, for convenience and easy comparison across domains, in Giga grams of gas emitted or in Giga grams of CO₂ equivalents (CO_{2eq}).

All relevant activity data from official national sources can be found in FAOSTAT, as specified in detail in Ch. 5. Following simple prescriptions, national GHG emissions can thus be computed at Tier 1 with official national data for all member countries, relative to the period 1961-present, and applying default IPCC guidelines. Exceptions are the Energy sub-domain, which given data availability through the International Energy Agency (IEA) can 'only' start from 1970, and the cultivation of organic soils and burning savanna sub-domains, which are not based on official national statistics and only cover the period 1990-present due to the nature of the underlying remote-sensing methodologies used to derive the underlying activity data. Emission projections to 2030 and 2050 can also calculated for all agriculture domains, offering an opportunity to member countries to produce reference scenarios to be used in setting future emission targets.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. They may be related to, *inter alia*, natural variability, partitioning fractions, lack of spatial or temporal coverage, or spatial aggregation.

The Table 4 below illustrates how the relevant national GHG data can be organized in a "Emissions – Agriculture" database domain. Each sub-domain identifies an emission source category. Each category includes elements, which provide the quantity of activity data (input) and the estimated emissions for each item (output). Each element is expressed with the corresponding unit of measurement.

TABLE 4
Data structure in the FAOSTAT "Emissions – Agriculture" domain

Subdomain	Item	Element	Unit
Enteric fermentation	Asses Buffalo Camels Dairy cattle Non-dairy cattle Goat Horses Llamas Mules Sheep Swine breeding Swine market Aggregated items: All GHG animals + (Total) Camels and llamas + (Total) Cattle + (Total) Mules and asses + (Total) Sheep and goats + (Total) Swine + (Total)	Stocks Implied emission factor for CH ₄ Emissions (CH ₄) Emissions (CO ₂ eq)	Head kg CH₄/head Gigagrams Gigagrams
Manure management	Asses Buffalo Camels Dairy cattle Non-dairy cattle Chickers broilers Chickens layers Ducks Goat Horses Llamas Mules Sheep Swine breeding Swine market Turkeys Aggregated items: All GHG animals + (Total) Camels and llamas + (Total) Cattle + (Total) Chickens + (Total) Mules and asses + (Total) Poultry birds + (Total) Sheep and goats + (Total) Swine + (Total)	Implied emission factor for CH ₄ Emissions (CH ₄) Emissions (CO ₂ eq) from CH ₄ Manure (N content) Implied emission factor for N ₂ O Direct emissions (N ₂ O) Indirect emissions (CO ₂ eq) Indirect emissions (CO ₂ eq) Emissions (N ₂ O) Emissions (CO ₂ eq) from N ₂ O Emissions (CO ₂ eq)	kg CH ₄ /head Gigagrams Gigagrams kg kg N ₂ O-N/kg N Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams
Rice cultivation	Rice, paddy	Area harvested Implied emission factor for CH ₄ Emissions (CH ₄) Emissions (CO ₂ eq)	ha g CH ₄ /m ² Gigagrams Gigagrams
Synthetic fertilizers GY	Nitrogen fertilizers (N total nutrients)	Consumption Implied emission factor for N_2O Direct emissions (N_2O) Direct emissions (CO_2 eq) Indirect emissions (N_2O) Indirect emissions (CO_2 eq) Emissions (N_2O) Emissions (N_2O)	Kg of nutrients kg N_2 O-N/kg N Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams

Subdomain	ltem	Element	Unit
Manure applied to soil	Asses Buffalo Camels Dairy cattle Non-dairy cattle Chickers broilers Chickens layers Ducks Goat Horses Llamas Mules Sheep Swine breeding Swine market Turkeys Aggregated items: All GHG animals + (Total) Camels and llamas + (Total) Cattle + (Total) Chickens + (Total) Mules and asses + (Total) Poultry birds + (Total) Sheep and goats + (Total) Swine + (Total)	Manure (N content) Implied emission factor for N ₂ O Direct emissions (N ₂ O) Direct emissions (CO ₂ eq) Indirect emissions (N ₂ O) Indirect emissions (CO ₂ eq) Emissions (N ₂ O) Emissions (CO ₂ eq)	Kg kg N₂O-N/kg N Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams
Manure left on pasture	Asses Buffalo Camels Dairy cattle Non-dairy cattle Chickers broilers Chickens layers Ducks Goat Horses Llamas Mules Sheep Swine breeding Swine market Turkeys Aggregated items: All GHG animals + (Total) Camels and llamas + (Total) Cattle + (Total) Chickens + (Total) Poultry birds + (Total) Sheep and goats + (Total) Swine + (Total)	Manure (N content) Implied emission factor for N ₂ O Direct emissions (N ₂ O) Direct emissions (CO ₂ eq) Indirect emissions (N ₂ O) Indirect emissions (CO ₂ eq) Emissions (N ₂ O) Emissions (CO ₂ eq)	Kg kg N₂O-N/kg N Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams
Crop residues	Barley Beans, dry Maize Millet Oats Potatoes Rice, paddy Rye Sorghum Soybeans Wheat Aggregated items: All GHG crops + (Total)	Residues (N content) Implied emission factor for N ₂ O Direct emissions (N ₂ O) Direct emissions (CO ₂ eq) Indirect emissions (CO ₂ eq) Emissions (N ₂ O) Emissions (N ₂ O) Emissions (CO ₂ eq)	Tonnes of nutrients kg N ₂ O kg ⁻¹ N in residues Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams
Cultivation of organic soils	Cropland organic soils Grassland organic soils Aggregated items: Cropland and grassland organic soils + (Total)	Area Implied emission factor for N ₂ O Emissions (N ₂ O) Emissions (CO ₂ eq)	ha kg N₂O-N/ha Gigagrams Gigagrams

Subdomain	Item	Element	Unit
Burning - Crop residues	Maize Rice, paddy Sugar cane Wheat Aggregated items: All GHG crops + (Total)	Biomass burned (dry matter) Implied emission factor for N ₂ O Implied emission factor for CH ₄ Emissions (N ₂ O) Emissions (CH ₄) Emissions (CO ₂ eq) from N ₂ O Emissions (CO ₂ eq) from CH ₄ Emissions (CO ₂ eq)	Tonnes g N ₂ O/kg dry matter g CH ₄ /kg dry matter Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams
Burning - Savanna	Savanna Woody savanna Closed Shrublands Open Shrublands Grasslands Aggregated items: Burning - All categories + (Total) Savanna and woody savanna + (Total) Closed and open Shrubland (+Total)	Burned area Biomass burned (dry matter) Implied emission factor for N ₂ O Implied emission factor for CH ₄ Emissions (N ₂ O) Emissions (CH ₄) Emissions (CO ₂ eq) from N ₂ O Emissions (CO ₂ eq) from CH ₄ Emissions (CO ₂ eq)	ha Tonnes g N ₂ O/kg dry matter g CH ₄ /kg dry matter Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams
Energy use	Gas - Diesel oils Gasoline Natural gas (including LNG) Liquefied petroleum gas (LPG) Residual fuel oil Hard coal Electricity Gas - Diesel oils in fisheries Residual fuel oil in fisheries Energy for power irrigation Aggregated items: Total energy + (Total) Transport fuel consumed in agriculture (excl. fishery) + (Total) Energy consumed in fishery + (Total)	Consumption in Agriculture Consumption in Agriculture Implied emission factor for CH ₄ Emissions (CO ₂ eq) from CH ₄ Implied emission factor for N ₂ O Emissions (N ₂ O) Emissions (CO ₂ eq) from N ₂ O Implied emission factor for CO ₂ Emissions (CO ₂) Emissions (CO ₂ eq)	Terajoule million kWh Kg/TJ Gigagrams Gigagrams Kg/TJ Gigagrams Gigagrams Gigagrams Gigagrams g/kWh Gigagrams Gigagrams

In the remainder of this document, a metadata sheet and a detailed methodology are presented for each of the subdomains above.

4.4 METADATA AND EMISSIONS CATEGORIES FOR FORESTRY AND LAND USE

Land use contains all the emissions produced in the different forestry and land use sub-domains. We further provide guidance on estimating emissions from biomass burning in all ecosystem categories other than in agriculture, based on remote-sensing information, consistently with updated FAO activity data aggregated at national level and IPCC guidelines.

GHG emissions from the Land use sector consist largely of CO_2 gas, generated mainly through cropland, grassland and forest management activities, including carbon gain and losses linked to anthopogenic land use changes. Emissions from biomass fires include in addition significant CH_4 and N_2O components. All emissions may be expressed, for convenience and easy comparison across domains, in Giga grams of gas emitted or in Giga grams of CO_2 equivalents (CO_{2eq}).

Relevant activity data from official national sources can be found in FRA, as well as in a set of ancillary geo-spatial data aggregated at national level, recently made available by FAO via FAOSTAT, as specified in detail in Ch. 5. Following simple prescriptions, national GHG emissions can thus be computed at Tier 1, land approach 1, using official national data for all member countries, relative to the period 1990-present. Exceptions are the cultivation of organic soils and burning biomass sub-domains, which are not based on official national statistics and only cover the period 1990-present due to the nature of the underlying remote-sensing methodologies and reporting characteristics of the statistics used to derive the underlying activity data.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors, carbon stocks and activity data. They may be related to, inter alia, natural variability, partitioning fractions, lack of spatial or temporal coverage, and spatial aggregation.

The Table 5 below illustrates how the relevant national GHG data can be organized in a "Emissions – Land use" database. Each sub-domain groups one or more emission source category. For each category, a number of elements provide the quantity of activity data (input) and the estimated emissions (output). Each element is expressed according to the correspondent unit measure, visible in the "Unit" column.

TABLE 5
Data structure of the "Emissions – Land Use" domain

Subdomain	Item	Element	Unit
Land use total	Forest land Cropland Grassland Burning biomass Aggregated items: Land use + (Total)	Net emissions/removals (CO_2) Net emissions/removals (CO_2 eq) Emissions (CO_2 eq) from N_2O Emissions (CO_2 eq) from CH_4	Gigagrams Gigagrams Gigagrams Gigagrams
Forest land GF	Forest land Net forest conversion Aggregated items: Forest land + (Total)	Area Implied emission factor for CO ₂ Net emissions/removals (CO ₂) Net emissions/removals (CO ₂ eq)	ha Tonnes CO₂/ha Gigagrams Gigagrams
Cropland GC	Cropland organic soils	Area Implied emission factor for C Net stock change (C) Net emissions/removals (CO ₂) Net emissions/removals (CO ₂ eq)	ha Tonnes CO ₂ /ha Gigagrams Gigagrams Gigagrams
Grassland GC	Grassland organic soils	Area Implied emission factor for C Net stock change (C) Net emissions/removals (CO ₂) Net emissions/removals (CO ₂ eq)	ha Tonnes C/ha Gigagrams Gigagrams Gigagrams
Burning - Biomass GI	Humid tropical forest Other forest Organic soils	Burned area Biomass burned (dry matter) Implied emission factor for N ₂ O Implied emission factor for CH ₄ Implied emission factor for CO ₂ Emissions (N ₂ O) Emissions (CH ₄) Emissions (CO ₂ eq) from N ₂ O Emissions (CO ₂ eq) from CH ₄ Emissions (CO ₂ eq) Emissions (CO ₂ eq)	ha Tonnes g N₂O/kg dry matter g CH₄/kg dry matter g CO₂/kg dry matter g CO₂/kg dry matter Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams Gigagrams

34

Minimum data availability for developing countries: Structure and data organization using FAOSTAT as guide

The FAOSTAT Emissions database, constructed on the basis of a robust methodology that conforms to the IPCC-2006 Guidelines, seeks to provide user-friendly support for policy decision makers and other users in Member Countries in preparing their GHG inventory of AFOLU sector. The database serves as a foundation for GHG reporting obligations, such as NCs and BURs under the UNFCCC.

In particular, the database was created to support Member Countries in their performance of the following activities:

- providing regular updates on global and regional trends in GHG emissions and removals from AFOLU;
- bridging gaps in Member Countries' capacity to assess and report GHG emissions and removals for AFOLU, in
 observance of the new requirements established by the Durban accords (BURs);
- setting up QA/QC procedures and performing data analyses, as the database provides an internationally-accepted and neutral data platform to support national reporting;
- developing indicators for further analysis, and deriving complex indexes useful for analysis and policy support.

The FAOSTAT Emissions database may be accessed from the following link http://faostat.fao.org/, as shown in Figure 13 below. The information is available in three languages: English, French and Spanish.

FIGURE 13 FAOSTAT database homepage



FAO collects, analyses, interprets, and disseminates information related to nutrition, food and agriculture¹, and furnishes such technical assistance at governments' requests. Member Countries, for their part, regularly communicate statistical and technical data, and other information published by governments (respectively, Articles I and XI of the FAO Constitution).

The Statistics Division of FAO collates and disseminates food and agricultural statistics globally². To collect and disseminate information adequately, Member Countries must cooperate in providing information, and relevant indicators on food security and agricultural development must be available to Member Countries. A calendar of disseminating activities is available every year, for each FAOSTAT domain. The global data refer to statistics on crops, livestock, irrigation, land use, fertilizers, pesticide consumption, agricultural machinery, and forestry, including statistics on imports and exports of woods and paper.

FAOSTAT is the corporate statistical database maintained by the Statistics Division, it was created pursuant to the need to provide a single point of statistical time series for data collection and entry, validation and generation, and analysis, and is fully compliant with the principles of good practice governing international statistics. Thus, FAOSTAT is the world's largest and most comprehensive statistical database on food and agriculture. The Statistics Division compiles, processes and stores statistical information by country starting from 1961. The database contains data from over 245 countries and 35 regional areas, and covers domains on agriculture (production, consumption, trade, prices and resources), nutrition, fisheries, forestry, food aid, land use, population and agro-environment.

The FAOSTAT database is based on a data collection system for agricultural and forest statistics that is implemented every year, through questionnaires delivered to Member Countries' National Statistics Offices (NSO's), and when applicable, to other relevant institutions such as Ministry of Agriculture or other specific departments. At year end, the FAO Statistics Division again sends questionnaires to Member Countries, notifying the relevant FAO country representatives and FAO Regional Offices, and collects data for the previous year. Over the subsequent year, the questionnaires are collected, analysed and interpreted.

¹ The term "agriculture" and its derivatives include fisheries, marine products, forestry and primary forestry products.

² http://www.fao.org/economic/ess/en/.


If the FAO Statistics Division is to fulfil its mandate and collect and disseminate information adequately, Member Countries must cooperate in providing information, and relevant indicators on food security and agricultural development must be available to Member Countries. A calendar of disseminating activities is available every year, for each FAOSTAT domain³. The global data refer to statistics on crops, livestock, irrigation, land use, fertilizers, pesticide consumption, agricultural machinery, and forestry, including statistics on imports and exports of woods and paper⁴.

The New FAOSTAT platform⁵ contains information organised by subject domain: Production (which includes crops, livestock), Trade, Food supply, Commodity balances, Food balance sheets, Prices, Resources, Population, Investments, Emissions-Agriculture, Emissions-Land Use (including emissions/removals from forest land), and Forestry. FAOSTAT is the main source of the time-series activity data required to compile the AFOLU GHG inventory.

FAOSTAT is also available in French and Spanish, and both platforms contain the same datasets. However, the New FAOSTAT offers easy access to data and introduces enhanced features tailored to the needs of a wider range of users. New domains in FAOSTAT, such as Emissions-Agriculture and Emissions-Land Use, address emerging data requirements related principally to global warming and environmental monitoring.

The homepage of FAOSTAT (depicted in Figure 14 below) has different options for accessing the GHG database.

FIGURE 14
FAOSTAT homepage



 $^{^3 \} http://faostat3.fao.org/home/faostatRelease Calendar.html \ (or \ Latest \ news: \ http://faostat.fao.org/site/291/default.aspx).$


⁴ http://www.fao.org/corp/statistics/en/.

⁵ http://faostat.fao.org/.

Clicking on the "Emissions – Agriculture" icon leads to the Standard download webpage (Figure 15 below), which displays the following options:


- The menu on the far left of the page lists all domains and sub-categories.
- The menu in the "Index" box links to each category's metadata.
- Go to Browse Data on the right of the page enables to visualize data.

FIGURE 15 The FAOSTAT Emissions database



The following Figure 16 shows an example of metadata for the "Enteric fermentation" category.

FIGURE 16
Metadata in the FAOSTAT Emissions database



The database has a hierarchical structure that comprises two main domains: "Emissions – Agriculture" and "Emissions – Land Use". Each domain includes a series of sub-domains, which identify emission categories. For each sub-domain, data are organized by four dimensions:

- Geographical Area: Countries, Regions, Global and Special Groups (i.e. aggregated by economic, political or other categories). The Regional and Global areas aggregate data by total and annual average;
- Elements: this section includes activity variables and estimated GHG emissions;
- Items: the set of activity data;
- Years: the time period of the data available, for instance, 1961-present.

Data can be selected in two different ways: "Browse data" and "Download data".

- a) Positioning the pointer on the "BROWSE DATA" icon at the top of the menu, the list of main categories appears. Click on the selected category to retrieve the maps and graphics required.
 - The "Browse data" option enables the user to retrieve aggregate data (annual average or sum) for immediate analysis.

The data are expressed in pie-charts, graphs and comparisons among countries and regions.

The figure 17 shows an example from the "EMISSION-AGRICULTURE" category. By selecting the item, area, year and aggregation variables on the query boxes, maps, data and graphics may be viewed.


FIGURE 17 The "Browse data" option



b) "DOWNLOAD DATA" enables actual retrieval of the data from the database.

With the "**Download data**" option, the estimated gas emissions become available for direct download. The following Figure 18 shows shows all available variables in the four sub-domains.






Example:

How can GHG emissions statistics be accessed? What is the yearly average of CO₂-equivalent emissions produced by rice cultivation in South-Eastern Asia from 2004 to 2010?


a) "Browse data" option:

FIGURE 19 Example "Browse data"

Step 1.



Step 2.




Step 3.




b) "Download data" option:

FIGURE 20 Example "Download data"


Step 1.




Step 2.



Step 3.



In the "Output Options" section, further down the same page, additional information is provided on the statistics selected.



5.1 AGRICULTURE

5.1.1 Enteric fermentation

Dataset information	
Title	Enteric fermentation
Definition	Greenhouse gas (GHG) emissions from enteric fermentation consist of methane gas produced in the digestive systems of ruminants and, to a lesser extent, of non-ruminants.
Methodology and qua	lity information
Methods and processing	GHG emissions from enteric fermentation consist of methane gas (CH4) produced in the digestive systems of ruminants and, to a lesser extent, of non-ruminants. The FAOSTAT data are computed at Tier 1, following IPCC 2006 Vol. 4, Ch. 10. The emissions are estimated at country level, using the formula:
	Emission = A * EF
	where
	• Emission = GHG emissions, in kg CH ₄ yr ⁻¹ ;
	 A = Activity data, representing the number of livestock in heads (1);
	• EF = Tier 1, default IPCC emission factors, expressed in units of kg CH ₄ head ⁻¹ yr ⁻¹ (2).
	(1) Activity data cover the following animal categories: buffaloes, sheep, goats, camels, llamas, horses, mules, asses, pigs, and dairy and non-dairy cattle*. For the period between 1961-present, the activity data are taken directly from FAOSTAT (domain: Production/Live animals). Projections of activity data for 2030 and 2050 for the categories of dairy and non-dairy cattle, buffaloes, sheep, goats, pigs and poultry, are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). Activity data for animal categories for which no FAO projection was available were set to the most recent available FAOSTAT value. The FAO projections used cover about 140 countries. Projections of activity data for countries that were not included assume the same growth rate of neighbouring countries.
	*FAOSTAT livestock data include cattle and dairy cattle. Dairy cattle data are expressed as heads of cows producing milk, and can be found under the Production/Livestock Primary domain by selecting the item "Cow milk, whole fresh" item and the "Producing animals/Slaughtered" element. Non-dairy cattle are derived from FAOSTAT categories, specifically as "cattle minus dairy cattle".
	(2) The EF values are specified by livestock category and regional grouping in the IPCC Guidelines, 2006, Vol. 4, Ch. 10, Tables 10.10 and 10.11.
	The dimensionless conversion factors used are
	 10⁻⁶, to convert the emissions from kg CH₄ to Gg CH₄; and
	• GWP-CH $_4$ = 21 (100-year time horizon global warming potential), to convert Gg CH $_4$ to Gg CO $_2$ eq (IPCC, 1996: Technical Summary, Tab 4, p. 22).
	The enteric fermentation domain contains the following data available for download: country-level GHG emissions in both Gg CH $_4$ and Gg CO $_2$ eq, by livestock species and by species aggregates, as well as their total; implied emission factors; and activity data. Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of enteric fermentation, further information is available in the IPCC Guidelines (Vol. 4, Ch. 10, Section 10.3.4).

Calculation procedure


Step 1. Activity data

In this sub-domain, activity data refers to the number of heads (Stock) of the following items: dairy and non-dairy cattle, buffaloes, sheep, goats, camels, horses, mules, asses, swine ("Pigs" in FAOSTAT nomenclature) and llamas ("Camelids, other" in FAOSTAT nomenclature).

The definitions used by FAOSTAT for the "Number of Live Animals/Stock" and for the individual items are available in the Glossary (see Annex 1), at "Definition of items-livestock" or on the FAOSTAT website, at http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E. The unit measure is expressed in terms of the number of heads.

Data (from 1961 to present) are to be found under the FAOSTAT – Production, Live Animals sub-domain:

FIGURE 21 "Number of heads of sheep" from the FAOSTAT database




For cattle and swine, in accordance with the 2006 IPCC prescriptions, the FAOSTAT Emissions provides emission data disaggregated by dairy and non-dairy cattle, breeding and market swine.

However, no corresponding stock data (heads) may be found in the FAOSTAT Production Live Animals sub-domain, because FAOSTAT does not collect this disaggregated data. However, this information can be retrieved from FAOSTAT with the following steps.

- 1. The data sets of "dairy" cattle and "non-dairy" cattle are obtained as follows:
 - the number of dairy cattle heads is in the Production Livestock Primary sub-domain, under the element "Producing Animals/Slaughtered" and the item "Milk, whole fresh":


FIGURE 22

"Number of heads of dairy cattle" from FAOSTAT database



- the number of non-dairy cattle heads may be derived by subtracting the number of heads of dairy cattle from the total cattle (Stocks), which can be found in the Production, Live Animals sub-domain, under the element "Stocks" and the item "Cattle".
- The global stock of non-dairy cattle in 2010 was 1,410,800 = 2,895,000 (total stock) -1,485,000 (dairy cattle).


FIGURE 23
"Number of heads of cattle" from the FAOSTAT database



- 2. The market swine data set may be obtained by multiplying the total heads of pigs from FAOSTAT Production Live Animals domain by 0.9, following the default 2006 IPCC Guidelines.
- 3. The breeding swine data set may be obtained by multiplying the total heads of pigs from FAOSTAT Production Live Animals domain by 0.1, following default 2006 IPCC Guidelines.

The FAOSTAT Emissions Enteric Fermentation sub-domain provides the data, disaggregated by sub-categories: dairy, non-dairy, breeding and market swine.

FIGURE 24 Livestock sub-categories of the FAOSTAT Emissions database



1.2 Projections of stocks

Projections of activity data for 2030 and 2050 for the dairy and non-dairy cattle, buffaloes, sheep, goats, pigs, categories were computed with respect to a baseline provided by FAO perspective studies⁶:

$$N_{(T)y} = \alpha_{(T)_y} \times N_{(T)_{2005-2007}}$$

- $N_{(T)y}$ = Number of heads for animal category T in the projected year y
- $\alpha_{(T)y}$ = Projection ratio for animal category T, in projected year y with respect to the baseline
- $N_{(T),2005-2007}$ = Baseline heads computed as the 2005, 2006, 2007 average heads of the animal category T
- T =Animal category
- y =Projected years 2030 and 2050.

⁶ Alexandratos, N. & Bruinsma, J. 2012. World agriculture towards 2030/2050: the 2012 revision. ESA Working Paper No. 12-03. Rome: FAO.

Step 2. Emissions (CH₄)

IPCC 2006, Equation 10.19

Emissions
$$(CH_4)_{(T)} = EF_{(T)} \times \frac{N_{(T)}}{10^6}$$

Where

- Emissions $(CH_4)_{(T)}$ = Methane emissions for animal category T, $Gg CH_4 yr^{-1}$
- $EF_{(T)}$ = emission factor for animal type T, kg CH₄ head⁻¹ (Table 1A)
- $N_{(T)}$ = number of heads for animal category T
- T = animal category

Step 3. Emissions (CO₂eq)

$$Emissions(CO_2eq)_{(T)} = Emissions(CH_4)_{(T)} \times GWP$$

Where

- Emissions $(CO_2eq)_{(T)}$ = Methane emissions in CO_2 equivalent for animal category T, $Gg CO_2eq yr^{-1}$
- Emissions $(CH_4)_{(T)}$ = Methane emissions for animal category T, Gg CH₄ yr⁻¹
- GWP = 21 (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq
- T =Animal category

Step 4. Implied emissions factor for CH₄

$$IEF_{(T)} = \frac{Emissions(CH_4)_{(T)}}{N_{(T)}} \times 10^6$$

Where

- $IEF_{(T)}$ = Implied emission factor for animal category T, kg CH₄ head⁻¹ yr⁻¹
- Emissions $(CH_4)_{(T)}$ = Methane emissions for animal category T, $Gg CH_4$ head⁻¹
- $N_{(T)}$ = Number of heads for animal category T
- T =Animal category

Example: What are the emissions from livestock enteric fermentation (cattle) in Morocco in the year 2010?

The emissions from enteric fermentation are estimated with the equation set out in Step 2 above.

We must therefore retrieve both the number of heads (the activity data) for the requested country by year, and the emission factor, as follows:

- 1. The number of heads of live animals is available in the FAOSTAT database Production domain, at http://faostat3.fao.org/faostat-gateway/go/to/download/G1/GE/E.
 - Collect the data set of the "dairy" cattle stock for Morocco.
 - Collect data of "non-dairy" cattle stock (which corresponds to "Other cattle" in IPCC-2006) for Morocco. In FAOSTAT, this is derived by subtracting, from the stock of cattle in Production/Live Animals, the number of heads for dairy cattle.

The stock of "Non-dairy" cattle in 2010 is 1,410,800 = 2,895,000 (total stock) -1,485,000 (dairy cattle). The values described in the following table result.

Livestock category	Annual Average Population (FAOSTAT) (Heads of cattle)
Dairy cows	1,485,000
Other cattle	1,410,800

2. The EF values are those specified for the livestock category and regional grouping in IPCC 2006, Vol. 4, Ch. 10, Tables 10.10 and 10.11.

TABLE 6
Tier 1 enteric fermentation emissions factors for cattle

Latin America: Commercialised dairy sector based on grazing. Separate beef cow herd grazing pastures	Dairy	72	Average milk production of 800 kg head-1 yr ⁻¹
and rangelands. Minor amount of feedlot feeding with grains. Growing non-dairy cattle comprise a large portion of the population.	Other cattle	56	Includes beef cows, bulls, and young.
Asia: Small commercialised dairy sector. Most cattle are multi-purpose, providing draft power and some milk within farming regions. Small grazing population.	Dairy	68	Average milk production of 1,650 kg head-1 yr ¹
Cattle of all types are smaller than those found in most other regions.	Other cattle	47	Includes multi-purpose cows, bulls, and young
Africa and Middle East: Commercialised dairy sector based on grazing with low production per cow. Most cattle are multi-purpose, providing draft power	Dairy	46	Average milk production of 475 kg head ⁻¹ yr ⁻¹
and some milk within farming regions. Some cattle graze over very large areas. Cattle are smaller than those found in most other regions.	Other cattle	31	Includes multi-purpose cows, bulls, and young

The solution is as follows:

TABLE 7
Results of livestock enteric fermentation (cattle) in Morocco in the year 2010

Livestock category	Annual average population (FAOSTAT) (Heads of cattle)	Emission factor for enteric fermentation (kg CH ₄ head ⁻¹ yr ⁻¹ (Table 10.11 Chapter 10, Vol. 4, 2006 IPPC Guidelines)	CH4 emission from livestock enteric fermentation (cattle) (Gg CH ₄ yr ⁻¹)
	(AD)	(EF)	(E=AD*EF)
Dairy cows	1,485,000	46	68.31
Other cattle	1,410,800	31	43.73
Total			112.04

5.1.2 Manure management

Dataset information	Dataset information		
Title	Manure management		
Definition	Greenhouse gas (GHG) emissions from manure management consist of methane and nitrous oxide gases from aerobic and anaerobic manure decomposition processes.		
Methodology and	quality information		
Methods and processing	GHG emissions from manure management consist of methane and nitrous oxide gases from aerobic and anaerobic decomposition processes. The FAOSTAT emission data are computed at Tier 1, following the IPCC 2006, Vol. 4, Ch. 10 and 11.		
	The term "manure" includes both urine and dung (i.e. both liquid and solid material) produced by livestock. More specifically, CH_4 gas is produced by the anaerobic decomposition of stored or treated manure, while N_2O is produced directly by nitrification and de-nitrification processes in the manure, and indirectly by nitrogen (N) volatilization and re-deposition processes, as well as from leaching of manure N.		
	CH ₄ emissions are estimated at the country level, using the formula:		
	Emission = A * EF		
	where		
	• Emission = GHG emissions, in kg CH ₄ yr ¹ ;		
	• A = Activity data, representing the number of livestock in heads (1);		
	 EF=Tier 1, default IPCC emission factors, expressed in units of kg CH₄ head⁻¹ yr¹ (2 (1) Activity data cover the following animal categories: buffaloes, sheep, goats, camels, 		
	llamas, horses, mules, asses, ducks, turkeys, dairy and non-dairy cattle*, chickens layers and broilers** and market and breeding swine***. For the period between 1961-present, activity data are taken directly from FAOSTAT (Production/Live animals domain). Projections of activity data for 2030 and 2050 for the categories of dairy and non-dairy cattle, buffaloes, sheep, goats, pigs and poultry, are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). Activity data for animal categories for which no FAO projections were were set to the most recent available FAOSTAT value. The FAO projections used cover about 140 countries. Projections of activity data for countries that were not included assume the same growth rate of neighbouring countries.		
	*FAOSTAT livestock data include the "cattle" and "dairy cattle" items. Dairy cattle data are expressed as heads of cows producing milk, and can be found under the Production/Livestock Primary domain by selecting the "Cow milk, whole fresh" item and the "Producing animals" element. Non-dairy cattle is derived from FAOSTAT categories, specifically as "cattle minus dairy cattle";		
	**FAOSTAT livestock data include the "chicken" and "chickens, layers" items. Chicken layers are expressed in 1000 heads of hens which have laid eggs in the reference period, and can be found under the Production/Livestock Primary domain by selecting the "Hen eggs, in shell" item and the "Producing animals" element "Chickens, broilers" is derived from FAOSTAT categories, specifically as "chickens minus chickens, layers";		
	***FAOSTAT livestock data include the "pigs" item. Market and breeding swine are calculated as 90% and 10% of the pigs item respectively (IPCC 2006, Vol. 4, Ch. 10, Table 10.19).		
	(2) EF values are assigned to each country (IPCC, 2006:Table 10.14 for cattle, buffaloes and pigs and Table 10.15 for all other animals), as a function of country-level average annual temperature (⁰ C). Average annual temperatures by country are taken from the FAO Global Agro-Ecological Zones (GAEZ) dataset (IIASA/FAO, 2012), relative to the baseline period 1961-1990.		

Methodology and quality information

Methods and processing

Direct N₂O emissions are estimated at the country level, using the formula:

Emission = A * EF

where

Emission = GHG emissions in kg N_2O-N yr⁻¹;

A = Activity data, representing the total amount of N in manure treated in Manure Management Systems (MMS) in kg N yr¹ (3);

EF=Tier 1, default IPCC emission factors, expressed in kg N₂O-N/kg N yr¹ (4).

- (3) This is the total amount of N excreted (i) for each livestock category (ii) treated in MMS (iii).
- (i) Following IPCC, 2006, Vol. 4, Ch. 10 Equation 10.30, the total amount of N excreted by each livestock category is calculated by multiplying the number of livestock heads by two coefficients: a) the Typical Animal Mass (TAM) and b) the N excretion coefficient (Nex). Both parameters vary according to the geographic region. TAM values are obtained from IPCC, 2006, Vol.4, Ch. 10, Annex 10A.2, Tables 10A-4 to 10A-9; Nex values are derived from IPCC, 2006, Vol. 4, Ch. 10, Table 10.19.
- (ii) See (1) for the livestock categories.
- (iii) Default IPCC percentages of total N treated in different MMS, by region and livestock category, are taken from IPCC, 2006, Vol. 4, Ch. 10, Annex 10A.2Tables 10A-4 to 10A-9 (for poultry: IPCC, 1997, Ch. 4Table 4.21).

(4) The EF values depend on the specific MMS, as per IPCC 2006, Vol. 4, Ch. 10, Table 10.21.

Indirect N₂O emissions are estimated at the country level, using the formula:

Emission = A * EF

where

- Emission = GHG emissions in kg N₂O-N yr⁻¹;
- A = Activity data, representing the fraction of the total amount of nitrogen (N) in manure treated in MMS that volatilises as NH₃ and NOx, and is lost through runoff and leaching, in kg N yr¹ (5);
- EF=Tier 1, default IPCC emission factors, expressed in kg N₂O-N kg -1N yr¹ (6).
- (5) The fractions for volatilisation by animals and MMS are taken from IPCC, 2006, Vol. 4, Ch. 10, Table 10.22. A mean fraction value of 10% for all countries is used for leaching (IPCC, 2006, Vol. 4, Ch. 10, note to Equation 10.28).
- (6) All countries are assigned global default EF values for volatilization and leaching (IPCC, 2006, Vol. 4, Ch. 11, Table 11.3).

The dimensionless conversion factors used are:

- 44/28, to convert the emissions from kg N₂O-N to kg N₂O gas;
- 10⁻⁶, to convert kg to Gg;
- GWP-CH₄ = 21 and GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg CH₄ or Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4 p. 22).

The manure management domain contains the following data categories, available for download: country-level GHG emissions, provided as total, direct and indirect amounts in Gg CH₄, Gg N₂O and Gg CO₂eq, by livestock species and by species aggregates, as well as their total; implied emission factors; and activity data.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of manure management, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol. 4, Ch. 10, Section 10.5.5).

Calculation procedure

Step 1. Activity data

In this domain, activity data refers to the number of heads (Stock) of the following items: dairy and non-dairy cattle, buffaloes, sheep, goats, chickens layers, chickens broilers, turkeys, ducks, market and breeding swine ("Pigs" in FAOSTAT nomenclature), camels and llamas ("Camelids, other" in FAOSTAT nomenclature). Camels and llamas data are used to calculate CH₄ emission only.


The definitions used by FAOSTAT for the "Number of Live Animals/Stock" and for the individual items are available in the Glossary (see Annex 1) at "Definition of items-livestock" or on the FAOSTAT website: http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E. The unit measure is expressed in number of heads.

Data (from 1961 to present) are to be found under the FAOSTAT – Production, Live Animals sub-domain and must be retrieved as described in the "Enteric fermentation activity data" section.

With regard to chickens layers and chickens broilers, data can be downloaded as follows:


1. the number of heads of chicken layers may be retrieved from the Production, Livestock Primary sub-domain, choosing the element "Producing Animals/Slaughtered", and the item "Eggs, hen eggs, in shell":

FIGURE 25
"Stock of laying animals" from the FAOSTAT database




- 2. the number of heads of broilers is obtained by subtracting the number of heads of layers from the stock of chickens.
- 3. The stock of chickens may be found in Production, Live Animals as follows:

FIGURE 26 "Stock of chickens" from the FAOSTAT database



The FAOSTAT Emissions manure management sub-domain provides data disaggregated by sub-categories: dairy and non-dairy cattle, breeding and market swine, chicken layers and broilers.

FIGURE 27
Manure management items from the FAOSTAT Emissions database



1.2 Projections of stocks

Projections of activity data for 2030 and 2050 for the dairy and non-dairy cattle, buffaloes, sheep, goats, pigs, poultry, etc. categories are computed with respect to a baseline provided by FAO perspective studies (Alexandratos, and Bruinsma, 2012). This is as follows:

$$N_{(T)y} = \alpha_{(T)y} \times N_{(T)_{2005-2007}}$$

Where

- $N_{(T)y}$ = Number of heads for animal category T in the projected year y
- $\alpha_{(T)y}$ = Projection ratio for animal category T, in projected year y with respect to the baseline
- $N_{(T) 2005-2007}$ = Baseline heads computed as the 2005, 2006, 2007 average heads of the animal category T
- T =Animal category
- y =Projected years 2030 and 2050.

Step 2. Emissions (CH₄)

IPCC 2006, Equation 10.22

$$Emissions(CH_4)_{(T)} = EF_{(T)} \times \frac{N_{(T)}}{10^6}$$

Where

- Emissions $(CH_4)_{(T)}$ = Methane emissions for animal category T, Gg CH₄ yr⁻¹
- $EF_{(T)}$ = Emission factor for T, CH₄ head⁻¹ yr⁻¹ (Table 2A)
- $N_{(T)}$ = Number of heads for animal category T, heads yr⁻¹
- T =Animal category

Step 3. Emissions (CO₂eq) for CH₄

$$Emissions(CO_2eq)_{(T)} = Emissions(CH_4)_{(T)} \times GWP$$

Where

- Emissions $(CO_2eq)_{(T)}$ = Methane emissions in CO_2 equivalent for animal category T, $Gg CO_2eq yr^{-1}$
- Emissions (CH₄) (T) = Methane emissions for animal category T, Gg CH₄ yr⁻¹
- GWP = 21 (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq
- T =Animal category

Step 4. Implied emissions factor for CH₄

$$IEF_{(T)} = \frac{Emissions(CH_4)_{(T)}}{N_{(T)}} \times 10^6$$

- IEF(T) = Implied emission factor for T, kg CH₄ head⁻¹
- Emissions (CH_4) $_{(T)}$ = Methane emissions for animal category T, $Gg CH_4 yr^{-1}$
- $N_{(T)}$ = Number of heads for animal category T heads yr⁻¹
- T =Animal category

Step 5. Excretion rate per animal

IPCC 2006, Equation 10.30

$$Nex_{(T)} = N_{rate(T)} \times \frac{TAM_{(T)}}{1000} \times 365$$

Where

- $Nex_{(T)} = N$ excreted in manure for animal category T, kg N animal⁻¹ yr⁻¹
- $N_{rate(T)}$ = Default N excretion rate per mass, kg N (tonnes animal mass)⁻¹ day⁻¹ (Table 3A)
- $TAM_{(T)}$ = Typical animal mass for animal category T, kg animal⁻¹ (Table 4A)
- T =Animal category

Step 6. Manure N content to manure management systems

$$NE_{MS(T)} = (N_{(T)} \times Nex_{(T)} \times MS_{(S,T)})$$

Where

- $NE_{MS(T)}$ = Total nitrogen excreted from manure management systems for animal category T, kg N yr⁻¹
- $N_{(T)}$ = Number of head of animal category T, heads yr⁻¹
- $Nex_{(T)} = Annual N excretion for animal category T, kg N animal⁻¹ yr⁻¹$
- $MS_{(S,T)}$ = Share of manure treated in each systems S for animal category T (Tables 5A to 13A)
- T =Animal category
- *S* = Manure management system (Lagoon, Slurry, Solid Storage, Drylot, Daily Spread, Digester, Pit<1Month, Pit>1Month, Other)

Step 7. Direct emissions (N2O)

IPCC 2006, Equation 10.25

$$Direct\ Emissions(N_2O)_{(T)} = \sum_{S} \left[NE_{MS} \times EF_{3(S)} \right] \times \frac{44}{28} \times 10^{-6}$$

Where

- Direct emissions(N_2O)_(T) = Direct N_2O emissions from manure management systems for animal category T, $Gg N_2O yr^{-1}$
- NE_{MS} = Total nitrogen excreted from manure management systems for animal category T, kg N yr⁻¹
- $EF_{3(S)}$ = Emission factor for direct N₂O emissions from each manure management system S, kg N₂O-N/kg N (Table 22A)
- T =Animal category
- S =Manure management system

Step 8. Direct emissions (CO₂eq)

$$Direct\ Emissions(CO_{2eq})_{(T)} =\ Direct\ Emissions(N_2O)_{(T)} \times GWP$$

- Direct emissions (CO_{2eq}) (T) = Direct N₂O emissions from manure management systems in CO₂ equivalent for animal category T, Gg CO₂eq yr⁻¹
- Direct emissions $(N_2O)_{(T)}$ = Direct N_2O emissions from manure management systems for animal category T, $Gg N_2O yr^{-1}$
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T =Animal category

Step 9. Indirect emissions (N2O)

$$Indirect \; Emissions(N_2O)_{(T)} = NE_{MS(T)} \times \left[(Frac_{GasMS(S)} \times EF_4) + (Frac_{LeachMS} \times EF_5) \right] \times \frac{44}{28} \times 10^{-6}$$

Where

- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions produced from the atmospheric deposition of N volatilised from manure management systems for animal category T, $Gg N_2O yr^{-1}$
- $NE_{MS(T)}$ = Total nitrogen excreted from manure management systems for animal category T, kg N yr⁻¹
- Frac_{GasMS(S)} = Fraction of applied organic N fertiliser materials that volatilises as NH₃ and NO_X, kg N volatilised from each system S (Tables 14A to 21A)
- EF₄ = Emission factor for indirect N₂O emissions from atmospheric deposition of N on soils and water surfaces,
 kg N-N₂O/kg NH₃-N + NO_x-N volatilised (Table 24A)
- Frac_{Leach} *= Fraction of applied organic N fertiliser materials that leaches as NH₃ and NO_X, kg N leached/kg of N applied (Table 23A)
- EF_5 = Emission factor for indirect N₂O emissions from N leaching and runoff, kg N₂O N/kg N (Table 25A)
- T = Animal category
- S = Manure management system
- * According to the note on p. 10.56 of the 2006 IPCC Guidelines: the percent of managed manure nitrogen losses for livestock category T due to runoff and leaching during solid and liquid storage of manure has a typical range of 1-20%; Tier 2 method.

Step 10. Indirect emissions (CO₂eq)

Indirect Emissions
$$(CO_{2eq})_{(T)} = Indirect Emissions (N_2O)_{(T)} \times GWP$$

Where

- Indirect emissions $(CO_2eq)_{(T)}$ = Indirect N₂O emissions from manure management systems in CO₂ equivalent, for animal category T, Gg CO₂eq yr⁻¹
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from manure management systems for animal category T, $Gg N_2O yr^{-1}$
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T =Animal category

Step 11. Emissions (N₂O)

$$Emissions(N_2O)_{(T)} = Direct \ Emissions(N_2O)_{(T)} + Indirect \ Emissions(N_2O)_{(T)}$$

- Emissions $(N_2O)_{(T)}$ = Total N₂O emissions from manure management systems for animal category T, Gg N₂O yr⁻¹
- Direct emissions(N_2O)_(T) = Direct N_2O emissions from manure management systems for animal category T, $Gg N_2O yr^{-1}$
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from manure applied on soils for animal category T, Gg N_2O yr^{-1}
- T =Animal category

Step 12. Implied emissions factor for N₂O

$$IEF_{(T)} = \frac{Emissions(N_2O)_{(T)}}{NE_{MMS(T)}} \times 10^6 \times \frac{28}{44}$$

Where

- IEF(T) = Implied emission factor for animal category T, kg $N_2O/$ kg N applied
- Emissions $(N_2O)_{(T)}$ = Total N_2O emissions from manure management systems for animal category T, Gg N_2O yr⁻¹
- $NE_{MMS(T)}$ = Total nitrogen excreted from manure management systems for animal category T, heads yr^{-1}
- T =Animal category

5.1.3 Rice cultivation

Dataset information	Dataset information		
Title	Rice cultivation		
Definition	Greenhouse gas (GHG) emissions from rice cultivation consist of methane gas from the anaerobic decomposition of organic matter in paddy fields.		
Methodology and	quality information		
Methods and processing	GHG emissions from rice cultivation consist of methane gas (CH ₄) emitted by anaerobic decomposition of organic matter in paddy fields. The FAOSTAT data are computed at Tier 1 following the criteria established in IPCC, 1997, Vol. 3, Ch. 4 and IPCC, 2000, Ch. 4.		
	The emissions are estimated at the country level, using the formula:		
	Emission = A * EF		
	where		
	• Emission = GHG emissions, in g CH ₄ m ⁻² yr ⁻¹ ;		
	• $A = Activity data$, representing the annual rice paddy harvested area in m ⁻² (1);		
	• EF = Tier 1, default IPCC emission factors, in g CH ₄ m ⁻² yr ⁻¹ (2).		
	(1) Activity data cover the following animal categories: buffaloes, sheep, goats, camels, llamas, horses, mules, asses, ducks, turkeys, dairy and non-dairy cattle*, chickens layers and broilers** and market and breeding swine***. For the period between 1961-present, activity data are taken directly from FAOSTAT (Production/Live animals domain). Projections of activity data for 2030 and 2050 for the categories of dairy and non-dairy cattle, buffaloes, sheep, goats, pigs and poultry, are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). Activity data for animal categories for which no FAO projections were were set to the most recent available FAOSTAT value. The FAO projections used cover about 140 countries. Projections of activity data for countries that were not included assume the same growth rate of neighbouring countries.		
	(2) Seasonally integrated EF values are specified for key rice producing countries in the IPCC Guidelines (IPCC, 1997, Vol. 3, Ch. 4, Table 4.13). For countries with no default EF value, the following strategy is applied: for Asia, countries are assigned an area-weighted average EF (15.7 g CH ₄ m ⁻² yr ¹); in all other regions, countries are either assigned the IPCC EF value reported for a neighbouring country, where one exists, or the IPCC global default EF value (20 g CH ₄ m ⁻² yr ¹). Also, seasonally integrated EF values are further modified by the application of a dimensionless scaling factor for water regimes (i) and a dimensionless correction factor for organic amendments (ii).		
	(i) The scaling factors for rice paddy water regimes (IPCC, 1997, Vol. 3, Ch. 4, Table 4.12) are in the range of 0-1. Specifically, for all countries, a scaling factor of 0.7 is used for rain-fed rice, and 0 for upland rice or dry conditions (IPCC, 2000, Table 3, p. 403).		

Methodology and quality information

Methods and processing

(ii) The correction factor for organic amendments is the default value of 2 for all countries, corresponding to the assumption that 40% of farmers use organic amendments (IPCC, 2000, Table 3, p. 403).

The dimensionless conversion factors used are

- 10⁻⁴, to convert the activity data from m⁻² to ha;
- 10⁻⁹, to convert the emissions from g CH₄ to Gg CH₄; and
- GWP-CH₄ = 21 (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22).

The rice cultivation sub-domain contains the following data categories available for download: country-level GHG emissions both in Gg CH_4 and Gg CO_2 eq; implied emission factors; and activity data.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of rice cultivation, more detailed information is available in the IPCC Guidelines (IPCC, 2000, Ch. 4, Section 4.9.1.6).


Calculation procedure

Step 1. Activity data

In this sub-domain, activity data refers to the paddy rice harvested area. The definition used by FAOSTAT for "Area Harvested" is available in the Glossary–Area harvested; the definition for the "Rice paddy" item may also be found at the Glossary-Rice paddy or on the FAOSTAT website, at http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E. The unit measure is expressed in hectares.

The data, from 1961-present, are disseminated in the FAOSTAT – Production, Crops sub-domain:

FIGURE 28
Paddy rice area harvested, from the FAOSTAT database



The data on the area harvested can be also downloaded from the FAOSTAT Emissions database, in particular from the Emissions – Agriculture Rice Cultivation sub-domain, at the following link http://faostat3.fao.org/faostat-gateway/go/to/download/G1/GR/E.

The area must be disaggregated into two water regimes, namely irrigated (i) and rainfed (j) regimes, using default IPCC shares (Table 29A).

Please note that the upland regime does not emit methane. Therefore so there is no need to calculate.

$$A_i = \alpha_i \times A$$

$$A_j = \alpha_j \times A$$

Where

- A_i = Area harvested of rice paddy for *i* water regimes, ha yr⁻¹
- A_i =Area harvested of rice paddy for j water regimes, ha yr⁻¹
- $A_{i,j}$ = Fraction representing the two water regimes, constant in time
- A =Rice paddy area harvested, ha yr⁻¹
- i and j = Respectively, irrigated and rainfed water regimes.

1.2 Projections of area

The projections of activity data for 2030 and 2050 are computed with respect to a baseline provided by FAO perspective studies (Alexandratos and Bruinsma, 2012), as follows:

$$A_y = \alpha_y \times A_{(2005-2007)}$$

Where:

- A_{y} = Area for rice in the projected year y
- $\alpha_{\rm v}$ = Projection ratio for crop area A, at projected year y with respect to the baseline
- $A_{2005-2007}$ = Baseline area for rice computed as the average of the 2005, 2006 and 2007 area values
- y =The projected years 2030 and 2050

Note: for the area projection too, the area should be split into different water regimes after Step 1.

Step 2. Emissions (CH₄)

The following equation was extrapolated from Table 3 of the IPCC 2002 background paper.

Emissions
$$(CH_4) = \frac{EF \times SF_o \times (A_i + [A_j \times SF_j])}{10^5}$$

- Emissions (CH_4) = Methane emissions per rice paddy, $Gg CH_4 yr^{-1}$
- EF = Seasonal methane emission factor, g m⁻² yr⁻¹ (Table 30A)
- $A_{i,i}$ = Rice paddy area harvested in the two water regimes, irrigated and rainfed, ha yr⁻¹
- $SF_o = 1.4$ correction factor for organic amendments, for all countries
- $SF_i = 0.7$ scaling factor for A_i

Step 3. Emissions (CO₂eq)

$$Emissions(CO_2eq)_{(T)} = Emissions(CH_4)_{(T)} \times GWP$$

Where

- $Emissions (CO_2eq) = Methane emissions in CO_2 equivalent per rice paddy, Gg CO_2eq yr^{-1}$
- Emissions (CH_4) = Methane emissions per rice paddy, $Gg CH_4yr^{-1}$
- GWP = 21 (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq

Step 4. Implied emissions factor for CH₄

$$EF = \frac{Emissions(CH_4)}{A} \times 10^5$$

Where

- *IEF* = Implied emission factor per rice paddy, g m⁻² yr⁻¹
- Emissions (CH_4) = Methane emissions per rice paddy, $Gg CH_4 yr^{-1}$
- A =Rice paddy harvested area, ha yr⁻¹

5.1.4 Synthetic fertilizers

Dataset information	
Title	Synthetic fertilisers
Definition	Greenhouse gas (GHG) emissions from synthetic fertilisers consist of nitrous oxide gas from synthetic nitrogen additions to managed soils.
Methodology and qua	lity information
Methods and processing	GHG emissions from synthetic fertilisers consist of direct and indirect nitrous oxide (N_2O) emissions from nitrogen (N) added to agricultural soils by farmers. Specifically, N_2O is produced by microbial processes of nitrification and de-nitrification, taking place on the addition site (direct emissions), and after volatilisation/re-deposition and leaching processes (indirect emissions). The FAOSTAT data are estimated at Tier 1, following IPCC, 2006, Vol. 4, Ch. 11.
	Direct emissions are estimated at country level, using the formula:
	Emission = A * EF
	• Emission = GHG emissions, in kg N ₂ O-N yr ⁻¹ ;
	 A = Activity data, representing the amount of annual synthetic N applications in kg N yr¹ (1);
	• EF =Tier 1, default IPCC emission factors, expressed in kg N ₂ O-N / kg N (2).
	(1) N consumption data are taken from the FAOSTAT Fertilisers Archive Dataset (1990-2001) (Inputs/ Fertilisers Archive sub domain) and the Fertilisers Dataset (2002 to present) (Inputs/Fertilizers sub domain). These are derived as an annual balance of N production and net trade. Projections of activity data for 2030 and 2050 are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). The FAO projections used cover about 140 countries. Projections of activity data for countries that are not included assume the same growth rate of neighbouring countries. The following assumption is made: N application data = FAOSTAT N consumption data.

Methodology and quality information

Methods and processing

(2) Global default EF values are taken from IPCC, 2006, Vol. 4, Ch.11, Table 11.1.

Indirect emissions are estimated at country level, using the formula:

Emission = A * EF

where

- Emission = GHG emissions, in kg N₂O-N yr⁻¹;
- A = Activity data, representing the amount of annual synthetic N applications that volatilises as NH₃ and NO_x and is lost through runoff and leaching in kg N yr¹ (3);
- EF=Tier 1, default IPCC emission factors, expressed in kg N₂O-N / kg N yr⁻¹ (4).
- (3) Obtained through the volatilisation and leaching factors in IPCC, 2006, Vol. 4, Ch. 11, Table 11.3.
- (4) Global default EF values from IPCC, 2006, Vol. 4, Ch. 11, Table 11.3.

The dimensionless conversion factors used are

- 44/28, to convert the emissions from kg N₂O-N to kg N₂O gas;
- 10⁻⁶, to convert the emissions from kg N₂O to Gg N₂O; and
- GWP-N₂O = 310 (100-year time horizon), to convert Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22).

The synthetic fertiliser sub-domain contains the following data categories available for download: country-level GHG emissions, provided as total, direct and indirect amounts in both Gg N₂O and Gg CO₂eq; implied emission factors; and activity data.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of synthetic fertilisers, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol.4, Ch. 11, Section 11.2.1.4 for direct emissions, and Section 11.2.2.4 for indirect emissions).

Calculation procedure

Step 1. Activity data

Activity data in this domain refers to the consumption of fertilisers. The definitions used by FAOSTAT of "Consumption of Fertilisers" is available in the Glossary–Consumption of Fertilizers (see Annex 1) or on the FAOSTAT website, specifically at http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E.

Data in the FAOSTAT Inputs sub domain is divided into two sub-domains that reflect the different methodologies adopted over time: the first series ("Fertilizers archive", see Figure 29 below) includes data from 1961 to 2002 and the second ("Fertilizers", Figure 30 below) lists data from 2002 to present. In the latter dataset, the element "Nitrogen Fertilizers (N total nutrients)" corresponds to the old "Nitrogenous Fertilizers"; "Consumption in nutrients (tonnes of nutrients)" corresponds to the old "Consumption".

Data collected with a revised methodology and presented in a new format. See http://faostat3.fao.org/faostat-gateway/go/to/download/E/*/E.

FIGURE 29 Fertilizers archive from the FAOSTAT database






FIGURE 30 Fertilizers dataset from the FAOSTAT database



In the FAOSTAT Emissions database, data from 1961 onwards is available in the Synthetic Fertilizers sub-domain with a unique "Nitrogen Fertilizers" item, but this query provides separate time series: the "Fertilizer Archive" data, from 1961 to 2002 (Nitrogenous fertilizers) and "Fertilizer" data from 2002 to present (Nitrogen fertilizers).

Data under "Nitrogen Fertilizers" has been converted into the nitrogen nutrients, and expressed in kg.

FIGURE 31
Synthetic fertilizers items from the FAOSTAT Emissions database



1.2 Projections of Consumption in Nutrients

The projections of activity data for 2030 and 2050 are computed with respect to a baseline provided by FAO perspective studies (Alexandratos and Bruinsma, 2012). The baseline is the following:

$$N_y = \alpha_y \times N_{2005-2007}$$

- $N_{\rm v}$ = Consumption in nutrients of nitrogen fertilizers in the projected year y
- α_v = Projection ratio for synthetic fertilizer, at projected year y with respect to the baseline
- $N_{2005-2007}$ = Baseline of the consumption in nutrients of nitrogen fertilizers computed, as the average of years 2005, 2006 and 2007, tonnes
- y =The projected years 2030 and 2050

Step 2. Direct emissions (N2O)

Direct Emissions
$$(N_2O) = N \times \frac{44}{28} \times EF_1 \times 10^{-6}$$

Where

- Direct emissions (N₂O) = Direct N₂O emissions from synthetic nitrogen additions to the managed soils, Gg N₂O yr⁻¹
- N = Consumption in nutrients of nitrogen fertilizers, kg N input yr⁻¹
- EF_I= Emission factor for N₂O emissions from N inputs, kg N₂O-N/kg N input (Table 26A)

N.B. The following assumption is made: N application data = FAOSTAT Nitrogen Consumption data.

Step 3. Direct emissions (CO₂eq)

$$Direct\ Emissions(CO_{2eq})_{(T)} = Direct\ Emissions(N_2O)_{(T)} \times GWP$$

Where

- Direct emissions (CO₂eq) = Direct N₂O emissions from synthetic nitrogen additions to the managed soils, in CO₂ equivalent, Gg CO₂eq yr⁻¹
- Direct emissions (N₂O) = Direct N₂O emissions from synthetic nitrogen additions to the managed soils, Gg N₂O yr⁻¹
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq

Step 4. Indirect emissions (N₂O)

$$Indirect\ Emissions\ (N_2O) = N \times [(Frac_{GASF} \times EF_4) + (Frac_{Leach} \times EF_5)] \times \frac{44}{28} \times 10^{-6}$$

Where

- Indirect emissions (N₂O) = Indirect N₂O emissions produced from atmospheric deposition of N, volatilised from managed soils, Gg N₂O-N yr⁻¹
- N =Consumption in nutrients of nitrogen fertilizers, kg N input
- Frac_{GASF} = Fraction of applied synthetic N fertilizer materials that volatilises as NH₃ and NO_x, kg N volatilised/ kg of N applied (Table 28A)
- EF_4 = Emission factor for N₂O emissions from atmospheric deposition of N on soils and water surfaces, kg N-N₂O/kg NH₃-N + NO_x-N volatilised (Table 24A)
- Frac_{Leach} = Fraction of applied synthetic N fertilizer materials that leaches as NH₃ and NO_X, kg N leached/kg of N additions (Table 27A)
- EF_5 = Emission factor for N₂O emissions from N leaching and runoff, kg N₂O-N/kg N (Table 25A)

Step 5. Indirect emissions (CO₂eq)

Indirect Emissions $(CO_{2eq})_{(T)} = Indirect Emissions(N_2O)_{(T)} \times GWP$

Where

- Indirect emissions (CO₂eq) = Indirect N₂O emissions from synthetic nitrogen additions to managed soils, in CO₂ equivalent, Gg CO₂eq yr⁻¹
- Indirect emissions (N₂O) = Indirect N₂O emissions from synthetic nitrogen additions to managed soils, Gg N₂O yr⁻¹
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq

Step 6. Emissions (N₂O)

Emissions
$$(N_2O)$$
 = Direct Emissions (N_2O) + Indirect Emissions (N_2O)

Where

- Emissions (N_2O) = Total N_2O emissions from synthetic nitrogen additions to managed soils, $Gg N_2O yr^{-1}$
- Direct emissions (N_2O) = Direct N_2O emissions from synthetic nitrogen additions to managed soils, $Gg N_2O yr^{-1}$
- Indirect emissions (N₂O) = Indirect N₂O emissions from synthetic nitrogen additions to managed soils, Gg N₂O yr⁻¹

Step 7. Emissions (CO₂eq)

Emissions
$$(CO_{2eq}) = Emissions(N_2O) \times GWP$$

Where

- Emissions (CO₂eq) = Total N₂O emissions from synthetic nitrogen additions to managed soils, in CO₂ equivalent, Gg CO₂eq yr⁻¹
- Emissions (N_2O) = Total N_2O emissions from synthetic nitrogen additions to managed soils, $Gg N_2O yr^{-1}$
- $GWP = 310 (1_0 0$ -year time h_orizon global warming potential), to convert Gg N₂O to Gg CO₂eq

Step 8. Implied emissions factor for N₂O

$$IEF = \frac{Emissions(N_2O)}{N} \times 10^6 \times \frac{28}{44}$$

- IEF = Implied emission factor for synthetic nitrogen additions to managed soils T, kg N₂O/kg of N applied yr⁻¹
- Emissions (N_2O) = Total N_2O emissions from synthetic nitrogen additions to managed soils, $Gg N_2O yr^{-1}$
- N =Consumption in nutrients of nitrogen fertilizers (kg)

5.1.5 Manure applied to soils

Dataset information		
Title	Manure applied to soils	
Definition	Greenhouse gas (GHG) emissions from manure applied to soils consist of nitrous oxide gas, from nitrogen additions in the treated manure that is applied to managed soils.	
Methodology and	quality information	
Methods and processing	GHG emissions from manure applied to soils consist of direct and indirect nitrous oxide (N_2O) emissions from manure nitrogen (N) added to agricultural soils by farmers. Specifically, N_2O is produced by microbial processes of nitrification and de-nitrification taking place on the application site (direct emissions), and after volatilisation/re-deposition and leaching processes (indirect emissions). The FAOSTAT data are estimated at Tier 1, following the IPCC Guidelines, 2006, Vol. 4, Ch.s 10 and 11.	
	Direct emissions are estimated at country level, using the formula:	
	Emission = A * EF	
	where	
	• Emission = GHG emissions, in kg N ₂ O-N yr ⁻¹ ;	
	 A = Activity data, representing the total amount of N in manure applied to soils in k N yr¹ (1); 	
	• EF =Tier 1, default IPCC emission factors in kg N ₂ O-N/kg N yr ⁻¹ (2).	
	(1) This is the amount of N excreted (see below, note <i>i</i>) by livestock (<i>ii</i>), treated in manure management systems (MMS) (<i>iii</i>) and net of losses through volatilisation, runoff and leaching from MMS, and other human use (<i>iv</i>), plus the N contribution from bedding materials where present (<i>v</i>).	
	(<i>i</i>) Following IPCC, 2006, Vol. 4, Ch. 10, Equation 10.30, the total amount of N excreted by each livestock category is calculated by multiplying the number of livestock heads by two coefficients: a) the Typical Animal Mass (TAM) and b) the N excretion coefficient ($N_{\rm ex}$). Both parameters vary according to geographic region. TAM values are obtained from IPCC, 2006, Vol. 4, Ch. 10, Annex 10A.2, Tables 10A-4 to 10A-9; $N_{\rm ex}$ values are derived from IPCC, 2006, Vol. 4, Ch. 10, Table 10.19.	
	(ii) Livestock data cover the following animal categories: buffaloes, sheep, goats, camels, llamas, horses, mules, asses, ducks and turkeys, dairy and non-dairy cattle*, chickens layers and broilers** and market and breeding swine***.	
	For the period 1961-present, activity data are taken directly from FAOSTAT (domain: Production/Live animals). Projections of activity data for 2030 and 2050 for the categories of dairy and non-dairy cattle, buffaloes, sheep, goats, pigs and poultry, are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). Activity data for animal categories for which no FAO projections were available were set to the most recent FAOSTAT value available. The FAO projections used cover about 140 countries. Projections of activity data for countries that were not included assume the same growth rate of neighbouring countries.	
	*FAOSTAT livestock data include the cattle and dairy cattle items. Dairy cattle data are expressed as heads of cows producing milk, and can be found under the Production/Livestock Primary domain by selecting the "Cow milk, whole fresh" item and the "Producing animals" element. Non-dairy cattle is derived from FAOSTAT categories, specifically as cattle minus dairy cattle.	
	**FAOSTAT livestock data include the items chickens and chicken layers. Chicken layers are expressed in 1000 heads of hens which have laid eggs in the reference period, and can be found under the Production/Livestock Primary domain by selecting the "Hen eggs, in shell" item. The "Producing animals. Chickens broilers" element is derived from FAOSTAT categories, specifically as chickens minus chickens layers;	
	***FAOSTAT livestock data include the "Pigs" item. Market and breeding swine are calculated, respectively, as 90% and 10% of item pigs (IPCC, 2006, Vol. 4, Ch. 10, Table 10.19).	

Methodology and quality information

Methods and processing

(*iii*) Default IPCC percentages of total N treated in different MMS, by region and livestock category, are taken from IPCC, 2006, Vol. 4, Ch. 10, Annex 10A.2, Tables 10A-4 to 10A-9 (for poultry: IPCC, 1997, Vol. 3, Ch. 4, Table 4.21).

(*iv*) Default IPCC values for total N losses from different MMS depend on the livestock category, as per IPCC, 2006, Vol. 4, Ch. 10, Tab. 23. Losses of treated manure due to use for construction, feed or fuels are set to zero, as per IPCC, 2006, Vol. 4, Ch. 11, p. 11.13. It is assumed that all treated manure, net of the above losses, is applied to soils following IPCC, 2006, Vol. 4, Ch. 11, p. 11.13.

(v) Default amounts of N used in bedding (for solid storage and deep bedding) as per IPCC, 2006, Vol. 4, Ch. 10, p. 10.66.

(2) Global IPCC default EF values are taken from IPCC, 2006: Vol. 4, Ch. 11, Table 11.1.

Indirect emissions are estimated at country level, using the formula:

Emission = A * EF

where

- Emission = GHG emissions in kg yr⁻¹;
- A = Activity data, representing the fraction of manure N applications that volatilises as NH₃ and NO_x, and is lost through runoff and leaching in kg N yr¹ (3);
- EF=Tier 1, default IPCC emission factors in kg N₂O-N/kg N yr⁻¹ (4).

(3) Obtained through the volatilisation and leaching factors in IPCC, 2006, Vol. 4, Ch. 11, Table 11.3.

(4) Global IPCC default EF values, from IPCC, 2006, Vol. 4, Ch. 11, Table 11.3.

The dimensionless conversion factors used are

- 44/28, to convert the emissions from kg N₂O-N to kg N₂O gas;
- 10⁻⁶, to convert the emissions from kg N₂O to Gg N₂O;
- GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22).

The "manure applied to soils" domain contains the following data categories available for download: country-level GHG emissions, provided as total, direct and indirect amounts in both $Gg\ N_2O$ and $Gg\ CO_2eq$; implied emission factors; and activity data.

Uncertainties in estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of manure applied to soil, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol. 4, Ch. 11, Section 11.2.1.4 for direct emissions, and Section 11.2.2.4 for indirect emissions).

Calculation procedure

Step 1. Activity data

Activity data in this subdomain refers to the number of heads (Stock) of the following items: dairy and non-dairy cattle, buffaloes, sheep, goats, chickens, turkeys, ducks, market swine and breeding swine ("Pigs" in FAOSTAT nomenclature).


The definitions used by FAOSTAT for the "Number of Live Animals/Stock" and for the individual items are available in the Glossary (see Annex 1) at "Definition of items-livestock", or on the FAOSTAT website, at http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E. The unit measure is expressed in terms of number of heads, except for poultry (chickens, ducks, and turkeys), which are expressed in terms of 1000 heads.

Data (from 1961 to present) are to be found under the FAOSTAT – Production, Live Animals sub-domain. Data of all live animals categories must be retrieved as described above in the "Enteric fermentation activity data" section. With regard to "Chickens, layers" and "Chickens, broilers", data can be retrieved as described in the "Manure management activity data" section.

The FAOSTAT Emissions database provides the data, disaggregated by the sub-categories of dairy and dairy cattle, swine, layers, and broilers.

FIGURE 32

Manure applied to soils items from the FAOSTAT Emissions database



1.2 Projections of stocks

Projections of activity data for 2030 and 2050 for the categories of dairy and non-dairy cattle, buffaloes, sheep, goats, pigs and poultry, are computed with respect to a baseline provided by FAO perspective studies (Alexandratos and Bruinsma, 2012). The baseline is as follows:

$$N_{(T)y} = \alpha_{(T)_y} \times N_{(T)_{2005-2007}}$$

Where:

- $N_{(T)}$ = Number of heads for animal category T, in the projected year y
- $\alpha_{(T)y}$ = Projection ratio for animal category T, in projected year y with respect to the baseline
- $N_{(T) 2005-2007}$ = Baseline heads computed as the average of the heads of animal category T for years 2005, 2006, 2007
- T =Animal category
- y =The projected years 2030 and 2050

Step 2. Excretion rate per animal

IPCC 2006, Equation 10.30

$$Nex_{(T)} = N_{rate(T)} \times \frac{TAM_{(T)}}{1000} \times 365$$

- $Nex_{(T)} = Annual N excretion for animal category T, kg N animal⁻¹ yr⁻¹$
- $N_{rate(T)}$ = Default N excretion rate, kg N (tonnes animal mass)⁻¹ day⁻¹ (Table 3A)
- $TAM_{(T)}$ = Typical animal mass for livestock category T, kg animal⁻¹(Table 4A)
- T =Animal category

Step 3. Nitrogen in manure available for application to managed soils and for other uses (fuel/feed/constructions)

IPCC 2006, Equation 10.30

$$N_{MS_Avb(T)} = \sum_{S} \left[N_{(T)} \times Nex_{(T)} \times MS_{(S,T)} \times \left(1 - Frac_{Loss(S,T)}\right) \right] + \left[N_{(T)} \times MS_{SolidStorage(T)} \times \frac{N_{bedding_SolidStorage(T)}}{1000} \right]$$

Where:

- $N_{MS_Avb(T)}$ = Amount of managed manure nitrogen available for application to managed soils and for other uses, for animal category T, kg N yr⁻¹
- $N_{(T)}$ = Number of heads of animal category T, heads yr⁻¹
- $Nex_{(T)} = Annual N excretion for animal category T, kg N animal⁻¹ yr⁻¹$
- $MS_{(S,T)}$ = Share of manure treated in each MS, for animal category T, (Tables 5A to 13A)
- $Frac_{Loss(S, T)}$ = Fraction of managed manure nitrogen for livestock category T that is lost in each system S (Tables 32A to 37A)
- MS_{SolidStorage(T)} = Share of manure treated in Solid Storage for animal category T (Table 7A)
- $N_{bedding_SolidStorage\ (T)}$ = Amount of nitrogen from bedding for solid storage MS for animal category T, kg N animal⁻¹ yr⁻¹ (Table 31A)
- T =Animal category
- *S* = Manure management system (Lagoon, Slurry, Solid Storage, Drylot, Daily Spread, Digester, Pit<1Month, Pit>1Month, Other)

Step 4. Nitrogen in manure applied to soils

IPCC 2006, Equation 11.4

$$F_{AM(T)} = N_{MS_Avb(T)} \times \left[1 - \left(Frac_{FEED(T)} + Frac_{FUEL(T)} + Frac_{CNST(T)}\right)\right]$$

Where

- $F_{AM(T)}$ = Amount of managed manure nitrogen applied to managed soils, for animal category T, kg N yr⁻¹
- $N_{MS_Avb(T)}$ = Amount of managed manure nitrogen available for application to managed soils or for feed fuel, or construction purposes, for animal category T, kg N yr⁻¹
- $Frac_{FFED(T)}$ = Fraction of managed manure used for feed for animal category T (Table 39A)
- $Frac_{FUEL(T)}$ = Fraction of managed manure used for fuel for animal category T (Table 40A)
- Frac_{CNST(T)} = Fraction of managed manure used for construction for animal category T (Table 41A)

Step 5. Direct emissions (N2O)

$$Direct\ Emissions(N_2O)_{(T)} = F_{AM(T)} \times EF_1 \times \frac{44}{28} \times 10^{-6}$$

- Direct emissions $(N_2O)_{(T)}$ = Direct N_2O emissions from manure applied on managed soils for animal category T, $Gg N_2O yr^{-1}$
- $F_{AM(T)}$ = Amount of animal manure N applied to managed soils, kg N yr⁻¹
- EF_I = Emission factor for N₂O emissions from N inputs, kg N₂O-N/kg N input (Table 26A)
- T =Animal category

Step 6. Direct emissions (CO2eq)

Direct Emissions(
$$CO_{2eq}$$
)_(T) = Direct Emissions(N_2O)_(T)×GWP

Where

- Direct emissions $(CO_2eq)_{(T)}$ = Direct N₂O emissions from manure applied on managed soils in CO₂ equivalent, for animal category T, Gg CO₂eq yr⁻¹
- Direct emissions $(N_2O)_{(T)}$ = Direct N_2O emissions from manure applied on managed soils for animal category T, $Gg N_2O yr^{-1}$
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T =Animal category

Step 7. Indirect emissions (N2O)

$$Indirect\ Emissions(N_2O)_{(T)} = F_{AM_{(T)}} \times [(Frac_{GASM} \times EF_4) + (Frac_{Leach} \times EF_5)] \times \frac{44}{28} \times 10^{-6}$$

Where

- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions produced from atmospheric deposition of N, volatilised from managed soils, for animal category T, $Gg N_2O yr^{-1}$
- $F_{AM(T)}$ = Amount of animal manure N applied to managed soils for animal category T, kg N yr⁻¹
- Frac_{GASM} = Fraction of applied organic N fertilizer materials that volatilises as NH₃ and NO_X, kg N volatilised/kg of N applied)(Table 38A)
- EF₄ = Emission factor for N₂O emissions from atmospheric deposition of N on soils and water surfaces, kg N₂O
 -N/kg NH₃-N + NO_x-N volatilised (Table 24A)
- Frac_{Leach} = Fraction of applied organic N fertilizer materials that leaches as NH₃ and NO_X, kg N leached/kg of N additions (Table 27A)
- EF_5 = Emission factor for N₂O emissions from N leaching and runoff, kg N₂O-N/kg N volatilized (Table 25A)
- T =Animal category

Step 8. Indirect emissions (CO₂eq)

$$Indirect\ Emissions(CO_{2eq})_{(T)} = Indirect\ Emissions(N_2O)_{(T)} \times GWP$$

- Indirect emissions (CO₂eq)_(T) = Indirect N₂O emissions from manure applied on managed soils in CO₂ equivalent for animal category T, Gg CO₂eq yr⁻¹
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from manure applied on managed soils for animal category T, Gg N_2O yr⁻¹
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T =Animal category

Step 9. Emissions (N2O)

$$Emissions(N_2O)_{(T)} = Direct \ Emissions(N_2O)_{(T)} + Indirect \ Emissions(N_2O)_{(T)}$$

Where

- Emissions $(N_2O)_{(T)}$ = Total N₂O emissions from manure applied on managed soils for animal category T, Gg N₂O yr⁻¹
- Direct emissions $(N_2O)_{(T)}$ = Direct N_2O emissions from manure applied on soils for animal category T, Gg N_2O yr^{-1}
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from manure applied on soils for animal category T, Gg N_2O yr^{-1}
- T =Animal category

Step 10. Emissions (CO₂eq)

$$Emissions(CO_{2eq})_{(T)} = Emissions(N_2O)_{(T)} \times GWP$$

Where

- Emissions $(CO_2eq)_{(T)}$ = Total N₂O emissions from manure applied on managed soils, in CO₂ equivalent, for animal category T, Gg CO₂eq yr⁻¹
- Emissions $(N_2O)_{(T)}$ = Total N₂O emissions from manure applied on managed soils for animal category T, Gg N₂O yr⁻¹
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T =Animal category

Step 11. Implied emissions factor for N₂O

$$IEF_{(T)} = \frac{Emissions(N_2O)_{(T)}}{F_{PRP(T)}} \times 10^6 \times \frac{28}{44}$$

- IEF(T) = Implied emission factor for animal category T, kg N₂O/kg of N applied
- Emissions $(N_2O)_{(T)}$ = Total N_2O emissions from manure applied on soils for animal category T, $Gg N_2O yr^{-1}$
- $F_{AM(T)}$ = Amount of managed manure nitrogen applied to managed soils for animal category T, kg N yr⁻¹
- T =Animal category

5.1.6 Manure left on pasture

Dataset information	Dataset information		
Title	Manure left on pastures		
Definition	Greenhouse Gases (GHG) emissions data from manure left on pasture consist of nitrous oxide gas from nitrogen additions, made by grazing livestock, to managed soils.		
Methodology and o	quality information		
	Direct emissions are estimated at country level, using the formula:		
	Emission = A * EF		
	where		
	• Emission = GHG emissions in kg yr ¹ ;		
	 A = Activity data, representing the total amount of manure N left on pasture in kg N yr¹ (1); 		
	• EF = Tier 1, default IPCC emission factors, expressed in kg N ₂ O-N/kg N yr ⁻¹ (2).		
	(1) Computed as per IPCC, 2006, Vol. 4, Ch. 11, Equation 11.5, as the amount of total N excreted (see below, note i) by livestock (ii) and left on pastures as urine and dung (iii).		
	(i) Following IPCC, 2006, Vol. 4, Ch. 10, Equation 10.30, the total <i>amount of nitrogen excreted in manure</i> is calculated, for each livestock category, by multiplying the number of livestock heads by two coefficients: a) the Typical Animal Mass (TAM) and b) the N excretion coefficient (N _{ex}). Both parameters vary according to geographic region. TAM values are obtained from IPCC, 2006, Vol. 4, Ch. 10, Annex 10A.2, Tables 10A-4 to 10A-9; N _{ex} values are derived from IPCC, 2006, Vol. 4, Ch. 10, Tables 10.19. (
	ii) Livestock data cover the following animal categories: buffalo, sheep, goats, camels, llamas, horses, mules, asses, ducks, turkeys, dairy and non-dairy cattle*, chickens layers and broilers** and market and breeding swine***. For the period between 1961-present, activity data are taken directly from FAOSTAT (domain: Production/Live animals). Projections of activity data for 2030 and 2050 for the categories of dairy and non-dairy cattle, buffaloes, sheep, goats, pigs and poultry, are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). Activity data for animal categories for which no FAO projections were available were set to the most recent available FAOSTAT value. The FAO projections used cover about 140 countries. Projections of activity data for countries that were not included assume the same growth rate experienced in neighbouring countries.		
	*FAOSTAT livestock data include the "Cattle" and "Dairy cattle" items. Dairy cattle data are expressed as heads of cows producing milk, and can be found under the Production/Livestock Primary domain by selecting the item "Cow milk, whole fresh" and the "Producing animals" element. Non-dairy cattle is derived from FAOSTAT categories, specifically as "Cattle" minus "Dairy cattle";		
	**FAOSTAT livestock data include the "Chickens" and "Chickens, layers" items. Chicken layers are expressed in 1000 heads of hens which have laid eggs in the reference period, and can be found under the Production/Livestock Primary domain by selecting the "Hen eggs, in shell" item and the "Producing animals" element. "Chickens, broilers" is derived from FAOSTAT categories by subtracting the values for "Chickens, layers" from chickens;		
	***FAOSTAT livestock data include the "Pigs" item. Market and breeding swine are calculated as, respectively, 90% and 10% of this item (IPCC, 2006, Vol. 4, Ch.10, Table 10.19).		

Methodology and quality information

Methods and processing

(iii) Default IPCC percentages of total excreted N in different Manure Management Systems (MMS) by region and livestock category, as per IPCC, 2006, Vol. 4, Ch. 10, Annex 10A.2, Tables 10A-4 to 10A-9 (for poultry: IPCC, 1997, Vol. 3, Ch. 4, Table 4.21). With reference to these tables specifically, the amount of manure left on pasture contributing to GHG emissions is the sum of 100% of the total excreted N left on pasture, as per IPCC percentage value "Pasture/Range/Paddock", and 50% of the total excreted N as per IPCC percentage value "Burned for Fuel" (IPCC, 2006, Vol. 4, Ch. 10, p. 10.58).

(2) Global default EF values taken from IPCC, 2006, Vol. 4, Ch. 11, Table 11.1.

Indirect emissions are estimated at country level, using the formula:

Emission = A * EF

where

- Emission = GHG emissions in kg yr⁻¹;
- A = Activity data, representing the fraction of manure N left on pastures that volatilises as NH₃ and NO_x and is lost through runoff and leaching in kg N yr¹ (3);
- EF=Tier 1, default IPCC emission factors, expressed in kg N₂O-N/kg N yr⁻¹ (4).
- (3) Obtained through the volatilisation and leaching factors in IPCC, 2006, Vol. 4, Ch. 11, Table 11.3.
- (4) Global IPCC default EF values from IPCC, 2006, Vol. 4, Ch. 11, Tab. 11.1.

The dimensionless conversion factors used are

- 44/28, to convert the emissions from kg N₂O-N to kg N₂O gas;
- 10⁻⁶, to convert the emissions from kg N₂O to Gg N₂O;
- GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22).

The "manure left on pastures" domain contains the following data categories available for download: country-level GHG emissions, provided as total, direct and indirect amounts in both Gg N₂O and Gg CO₂eq; implied emission factors; and activity data.

Uncertainties in the estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of manure left on pastures, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol. 4, Ch. 11, Section 11.2.1.4 for direct emissions, and Section 11.2.2.4 for indirect emissions).

Calculation procedure

Step 1. Activity data

In this sub-domain, activity data refers to the number of heads (Stock) of the following items: dairy and non-dairy cattle, buffaloes, sheep, goats, chickens, turkeys, ducks, market swine and breeding swine ("Pigs" in FAOSTAT nomenclature).


The definitions used by FAOSTAT for the "Number of Live Animals/Stock" and for individual items are available in the Glossary (see Annex 1) at "Definition of items-livestock" or on the FAOSTAT website, at http://faostat3.fao. org/faostat-gateway/go/to/mes/glossary/*/E. The unit measure is expressed in terms of numbers of heads, except for poultry (chickens, ducks, and turkeys), which are expressed in terms of 1000 heads.

Data (from 1961 to present) are to be found under the FAOSTAT – Production, Live Animals sub-domain. Data of all categories must be retrieved as described in the "Enteric fermentation activity data" section. With regard to the "Chickens, layers" and "Chickens, broilers", data can be retrieved as described in the "Manure management activity data" section above.

The FAOSTAT Emissions database provides the data required, disaggregated by the sub-categories of dairy and dairy cattle, swine, layers, and broilers.

FIGURE 33

Manure left on pasture items from the FAOSTAT Emissions database



1.2 Projections of stocks

Projections of activity data for 2030 and 2050 for the categories of dairy and non-dairy cattle, buffaloes, sheep, goats, pigs and poultry, are computed with respect to a baseline provided by FAO perspective studies (Alexandratos and Bruinsma, 2012). The baseline is as follows:

$$N_{(T)y} = \alpha_{(T)y} \times N_{(T)_{2005-2007}}$$

Where:

- $N_{(T)}$ = Number of heads for animal category T in the projected year y
- $\alpha_{(T)y}$ = Projection ratio for animal category T, in projected year y with respect to the baseline
- $N_{(T) 2005-2007}$ = Baseline heads computed as the average of the heads of animal category T for years 2005, 2006, 2007
- T =Animal category
- y =The projected years 2030 and 2050

Step 2. Excretion rate per animal

IPCC 2006, Equation 10.30

$$Nex_{(T)} = N_{rate(T)} \times \frac{TAM_{(T)}}{1000} \times 365$$

- $Nex_{(T)} = \text{Annual N excretion for animal category } T$, kg N animal⁻¹ yr⁻¹
- $N_{rate(T)}$ = Default N excretion rate, kg N/tonnes animal mass day⁻¹ (Table 3A)
- $TAM_{(T)}$ = Typical animal mass for livestock category T, kg animal⁻¹ (Table 4A)
- T =Animal category

Step 3. Nitrogen in urine and dung deposited by grazing animals on pastures, ranges and paddocks

$$F_{PRP(T)} = \left[\left(N_{(T)} \times Nex_{(T)} \right) \times MS_{PRP(T)} \right] + \frac{\left[N_{(T)} \times Nex_{(T)} \times MS_{BurnedForFuel(T)} \right]}{2}$$

Where:

- $F_{PRP(T)}$ = Amount of animal manure N left on pastures for animal category T, kg N yr⁻¹
- $N_{(T)}$ = Number of heads of animal category T, heads yr⁻¹
- $Nex_{(T)} = Annual N excretion for animal category T, kg N animal⁻¹ yr⁻¹$
- $MS_{PRP(T)}$ = Share of total annual N excretion for each animal category T that is deposited on pastures, ranges and paddocks (Table 42A)
- $MS_{BurnedForFuel(T)}$ = Share of total annual N excretion for each animal category T that is deposited on pastures, ranges and paddocks in the form of urine (Table 43A)
- T =Animal category

NB: the N in the urine (50%) remains in the field; the N in the dung (50%) is removed and burned (IPCC, 2006, Vol. 4 Ch. 10 p. 10.58).

Step 4. Direct emissions (N₂O)

$$Direct\ Emissions\ (N_2O)_{(T)} = F_{PRP(T)} \times EF_{3PRP(T)} \times \frac{44}{28} \times 10^{-6}$$

Where

- Direct emissions(N_2O)_(T) = Direct N_2O emissions from manure left on pastures for animal category T, $Gg\ N_2O\ yr^{-1}$
- $F_{PRP(T)}$ = Annual amount of animal manure N applied to managed soils for animal category T, kg N yr⁻¹
- $EF_{3PRP(T)}$ = Emission factor for N₂O emissions from N inputs on pastures for animal category T, kg N₂O-N/kg N input (Table 44A)
- T =Animal category

Step 5. Direct emissions (CO₂eq)

$$Direct\ Emissions(CO_{2eq})_{(T)} = Direct\ Emissions(N_2O)_{(T)} \times GWP$$

- Direct emissions (CO_2eq)_(T) = Direct N₂O emissions from manure left on pastures in CO₂ equivalent, for animal category T, Gg CO₂eq yr⁻¹
- Direct emissions, $N_2O)_{(T)}$ = Direct N_2O emissions from manure left on pastures for animal category T, $Gg\ N_2O\ yr^{-1}$
- $GWP = 310 (100 \text{-year } t_i \text{me } h_0 \text{rizon global warming potential})$, to convert Gg N₂O to Gg CO₂eq
- $T = Animal categor_v$

Step 6. Indirect emissions (N2O)

$$Indirect\ Emissions(N_2O)_{(T)} = F_{AM_{(T)}} \times [(Frac_{GASM} \times EF_4) + (Frac_{Leach} \times EF_5)] \times \frac{44}{28} \times 10^{-6}$$

Where

- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O-N emissions produced from atmospheric deposition of N volatilised from managed soils, for animal category T, $Gg N_2O-N yr^{-1}$
- $F_{PRP(T)}$ = Amount of animal manure N left on pastures for animal category, kg N yr⁻¹
- Frac_{GASM} = Fraction of applied organic N fertilizer materials that volatilises as NH₃ and NO_X, kg N volatilised/ kg of N applied (Table 38A)
- EF₄= Emission factor for N₂O emissions from atmospheric deposition of N on soils and water surfaces, kg N₂O
 N/kg NH₃-N + NO_x-N volatilised (Table 24A)
- Frac_{Leach} = Fraction of applied organic N fertilizer materials that leaches as NH₃ and NO_X,, Kg N leached/kg of N applied (Table 27A)
- EF_5 = Emission factor for N₂O emissions from N leaching and runoff, kg N₂O –N/kg N volatilised (Table 25A)
- T =Animal category

Step 7. Indirect emissions (CO₂eq)

$$Indirect\ Emissions(CO_{2eq})_{(T)} = Indirect\ Emissions(N_2O)_{(T)} \times GWP$$

Where

- Indirect emissions (CO₂eq) (T) = Indirect N₂O emissions from manure left on pastures in CO₂ equivalent, for animal category T, Gg CO₂eq yr⁻¹
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from manure left on pastures for animal category T, Gg N_2O yr^{-1}
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T =Animal category

Step 8. Emissions (N₂O)

$$Emissions(N_2O)_{(T)} = Direct\ Emissions(N_2O)_{(T)} + Indirect\ Emissions(N_2O)_{(T)}$$

- Emissions $(N_2O)_{(T)}$ = Total N₂O emissions from manure left on pastures for animal category T, Gg N₂O yr⁻¹
- Direct emissions(N_2O)_(T) = Direct N_2O emissions from manure left on pastures for animal category T, $Gg\ N_2O$ yr⁻¹
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from manure left on pastures for animal category T, Gg N_2O yr^{-1}
- T =Animal category

Step 9. Emissions (CO2eq)

$$Emissions(CO_{2eq})_{(T)} = Emissions(N_2O)_{(T)} \times GWP$$

Where

- Emissions (CO_2eq) $_{(T)}$ = Total N₂O emissions from manure left on pastures in CO₂ equivalent, for animal category T, Gg CO₂eq yr⁻¹
- Emissions $(N_2O)_{(T)}$ = Total N₂O emissions from manure left on pastures for animal category T, Gg N₂O yr⁻¹
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T = Animal category

Step 10. Implied emissions for N₂O

$$IEF_{(T)} = \frac{Emissions(N_2O)_{(T)}}{F_{PRP(T)}} \times 10^6 \times \frac{28}{44}$$

Where

- $IEF_{(T)}$ = Implied emission factor for animal category T (kg N₂O/kg of N applied)
- Emissions $(N_2O)_{(T)}$ = Total N_2O emissions from manure applied on soils for animal category T, $Gg N_2O yr^{-1}$
- $FPRP_{(T)}$ = Amount of animal manure N left on pastures for animal category T, kg N yr⁻¹
- T =Animal category

5.1.7 Crop residues

Dataset information	
Title	Crop residues
Definition	Greenhouse Gas (GHG) emissions from crop residues consist of nitrous oxide gas deriving from the decomposition of nitrogen in crop residues, left on managed soils.
Methodology and qua	lity information
Methods and processing	Greenhouse gas (GHG) emissions from crop residues consist of direct and indirect nitrous oxide (N_2O) emissions from nitrogen (N) in crop residues, and forage/pasture renewal left on agricultural fields by farmers. Specifically, N_2O is produced by microbial processes of nitrification and de-nitrification taking place on the deposition site (direct emissions), and after volatilisation/re-deposition and leaching processes (indirect emissions). The FAOSTAT data are estimated at Tier 1, following IPCC, 2006, Vol. 4, Ch.s 2 and 11.
	Direct emissions are estimated at the country level, using the formula:
	Emission = A * EF where
	 Emission = GHG emissions in units of Kg N₂O yr¹;
	• $A = Activity data$, representing the total amount of N in crop residues in Kg N yr ¹ , (1);
	• EF =Tier 1, default IPCC emission factors, expressed in Kg N ₂ O-N/Kg N yr ¹ (2).

Methodology and quality information

Methods and processing

(1) Activity data are calculated from crop yield and harvested area, and cover for the following crop categories: Barley, Beans-dry, Maize, Millet, Oats, Potatoes, Ricepaddy, Rye, Sorghum, Soybeans, and Wheat. Crop yield and harvested area data are used to estimate the amount of biomass N in above and below-ground residues, by crop and by country, using Equation 11.6 in IPCC, 2006, Vol. 4, Ch. 11, and the default crop values available in Table 11.2. In some cases where default parameters were not provided (N content of below-ground residues for Rice and Millet, and Ratio of below-ground residues to above-ground biomass for Millet, Sorghum, Rye, and Beans-dry) the corresponding default values for crops with similar biophysical characteristics were used. This biomass N amount is then reduced by the fraction of crop residue burnt on site, assumed to be 10% by area, following IPCC, 2000, Ch. 4, Section 4A.2.1.1 with specified combustion coefficients by crop, as per IPCC, 2006, Vol. 4, Ch. 2, Table 2.6. Finally, all N in crop residues, net of amount burnt, is assumed to remain on the field, as per IPCC, 2006, Vol. 4, Ch. 11, Equation 11.6.

For the period 1961-present, data relating to crop yield and harvested area are taken from FAOSTAT (domain: Production/Crops). Projections of crop yield and harvested area for 2030 and 2050 are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). The FAO projections used cover about 140 countries. Projections of activity data for countries not included assume the same growth rate experienced in neighbouring countries.

(2) Global default EF values taken from IPCC, 2006, Vol. 4, Ch. 11, Table 11.1.

Indirect emissions are estimated at country level, using the formula:

Emission = A * EF

where

- Emission = GHG emissions, in units of Gg N₂O yr⁻¹;
- A = Activity data, representing the fraction of N in crop residues forage/pasture renewal
 that is lost through runoff and leaching in kg N yr¹ (3);
- EF=Tier 1, default IPCC emission factors, expressed in kg N₂O-N / kg N yr⁻¹ (4).
- (3) Obtained through the leaching factor in IPCC, 2006, Vol. 4, Ch. 11, Table 11.3.
- (4) Global IPCC default EF values from IPCC, 2006, Vol. 4, Ch. 11, Table 11.1.

The dimensionless conversion factors used are

- 10⁻³, to convert the activity data from kg to tonnes;
- 44/28, to convert the emissions from kg N2O-N to kg N2O gas;
- 10^{-6} , to convert the emissions from kg N₂O to Gg N₂O; and
- GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22).

The Crop residues domain contains the following data categories, which are available for download: country-level GHG emissions, provided as total, direct and indirect amounts in both $Gg\ N_2O$ and $Gg\ CO_2eq$; implied emission factors; and activity data.

Uncertainties in the estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of crop residues, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol.4, Ch. 11, Section 11.2.1.4 for direct emissions, and Section 11.2.2.4 for indirect emissions).

Calculation procedure


Step 1. Activity data

In this sub-domain, activity data refers to the Area harvested and the Yield of the following crop items: Barley, Dry beans, Maize, Millet, Oats, Potatoes, Paddy rice, Rye, Sorghum, Soybeans, and Wheat.

Definitions used by FAOSTAT for Area harvested", "Yield" and for the individual items are available on Glossary, or on the FAOSTAT website at http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E. The unit measure is expressed in "ha" for the Area harvested and in "hg/ha" for the Yield.


Data from 1961 to present are disseminated from the FAOSTAT – Production, Crops sub-domain, as follows:

FIGURE 34 Area harvested and yield data



For the Crop Residues sub-domain, the FAOSTAT Emissions database only includes the selected items. The unit measure is expressed in tonnes of nutrients of nitrogen.

FIGURE 35 Crops residues items



1.2 Projections of area

The projections of activity data for 2030 and 2050 are computed with respect to a baseline provided by FAO perspective studies (Alexandratos and Bruinsma, 2012), as follows:

$$A_{(T)y} = \alpha_{(T)y} \times A_{(T)_{2005-2007}}$$

Where:

- $A_{(T)y}$ = Area for crop type T in the projected year y
- $\alpha_{(T)y}$ = Projection ratio for crop area A, in projected year y with respect to the baseline
- $A_{(T) 2005-2007}$ = Baseline area for crop type T, computed as the average of the 2005, 2006, 2007 area values
- T = Crop type
- y =The projected years, 2030 and 2050

Step 2. Harvested annual Dry Matter Yield

$$Crop_{(T)} = \frac{Crop_{fresh(T)} \times DRY_{(T)}}{10}$$

Where

- $Crop_{(T)}$ = Harvested dry matter yield for crop T, kg dry matter ha⁻¹ yr⁻¹
- $Crop_{fresh(T)}$ = Harvested fresh yield for crop T, hg ha⁻¹ yr⁻¹
- $DRY_{(T)} = Dry$ matter fraction of harvested product for crop T (Table 45A)
- T = Crop item

Step 3. Area Burned

$$AreaBurnt_{(T)} = Area_{(T)} \times Frac_{Burnt}$$

Where:

- $AreaBurnt_{(T)} = Area of crop T burnt, ha yr^{-1}$
- $Area_{(T)} = Total$ annual area harvested of crop T, ha yr⁻¹
- $Frac_{Burnt} = 0.1$ fraction of burnt area
- T = Crop items

Step 4. Above-ground residues dry matter

$$AG_{DM(T)} = \frac{Crop_{(T)}}{1000} \times slope_{(T)} + intercept_{(T)}$$

- $AG_{DM(T)}$ = Above-ground residues dry matter for crop T, kg dry matter ha⁻¹
- $Crop_{(T)}$ = Harvested dry matter yield for crop T, kg dry matter ha⁻¹
- $Slope_{(T)} = Slope$ element (Table 46A)
- $Intercept_{(T)} = Intercept element (Table 47A)$
- T = Crop items

Step 5. Ratio of above-ground residues dry matter (AGDM(T)) to harvested yield

$$R_{AG(T)} = \frac{(AG_{DM(T)} * 1000)}{Crop_{(T)}}$$

Where

- $R_{AG(T)}$ = Ratio of above-ground residues dry matter $(AG_{DM(T)})$ to harvested yield for crop T
- $AG_{DM(T)}$ = Above-ground residues dry matter for crop T, kg dry matter ha⁻¹
- $Crop_{(T)}$ = Harvested annual dry matter yield for crop T, kg dry matter ha⁻¹
- T = Crop items

Step 6. Ratio of below-ground residues dry matter to harvested yield for crop T

$$R_{BG(T)} = R_{BG-BIO(T)} \times \left[\frac{\left(\left(AG_{DM(T)} * 1000 \right) + Crop_{(T)} \right)}{Crop_{(T)}} \right]$$

Where

- $R_{BG(T)}$ = Ratio of below-ground residues to harvested yield for crop T, kg dry matter/kg dry matter
- $R_{BG-BIO(T)}$ = Ratio of below-ground residues to above-ground biomass for crop T, kg dry matter (Table 48A)
- $AG_{DM(T)}$ = Above-ground residues dry matter for crop T, kg dry matter ha⁻¹
- $Crop_{(T)}$ = Harvested annual dry matter yield for crop T, kg dry matter ha⁻¹
- T = Crop items

Step 7. Residues (N content)

IPCC 2006, Equation 11.6

$$F_{CR(T)} = Crop_{(T)} \times \left(Area_{(T)} - AreaBurnt_{(T)} \times C_f\right) \times Frac_{Renew} \times \left[R_{AG(T)} \times N_{AG(T)} \times (1 - Frac_{Remove}) + R_{BG(T)} \times N_{BG(T)}\right]$$

- $F_{CR(T)}$ = Amount of N in crop residues (above and below ground), including N-fixing crops, and from forage/pasture renewal, returned to soils annually for crop T, kg N yr⁻¹
- $Crop_{(T)}$ = Harvested annual dry matter yield for crop T, kg dry matter ha⁻¹
- $Area_{(T)}$ = Total annual area harvested of crop T, ha yr⁻¹
- Area $Burnt_{(T)}$ = Annual area of crop T that is burnt, ha yr⁻¹
- C_f = Combustion factor (dimensionless) (Table 49A)
- $Frac_{Renew(T)}$ = Fraction of total area under annual crop that is renewed annually
- $R_{AG(T)}$ = Ratio of above-ground residues dry matter $(AG_{DM(T)})$ to harvested yield, for crop $T(Crop_{(T)})$ kg dry matter/kg dry matter
- $N_{AG(T)} = N$ content of above-ground residues for crop T, kg N/kg dry matter (Table 50A)
- $Frac_{Remove(T)}$ = Fraction of above-ground residues of crop T removed annually for purposes such as feed, bedding and construction, kg N/kg dry matter
- $R_{BG(T)}$ = Ratio of below-ground residues to harvested yield for crop T, kg dry matter/kg dry matter
- $N_{BG(T)} = N$ content of below-ground residues for crop T, kg N/kg dry matter (Table 51A)
- T = Crop items

Step 8. Direct emissions (N2O)

$$Direct\ Emissions(N_2O)_{(T)} = F_{CR(T)}*EF_1 \times \frac{44}{28} \times 10^{-6}$$

Where

- Direct emissions $(N_2O)_{(T)}$ = Direct N_2O emissions from crop residues and Forage/Pasture renewal for crop T, $Gg N_2O yr^{-1}$
- $F_{CR(T)}$ = Annual amount of N in crop residues (above and below ground), including N-fixing crops, and from forage/pasture renewal, returned to soils annually for crop T, kg N yr⁻¹
- EF_I = Emission factor for N₂O emissions from N inputs, kg N₂O-N/kg N input (Table 26A)
- T = Crop item

Step 9. Direct emissions (CO₂eq)

Direct Emissions (
$$CO_2eq$$
) = Direct Emissions (N_2O)×GWP

Where

- Direct emissions $(CO_2eq)_{(T)}$ = Direct N₂O emissions from crop residues and Forage/Pasture renewal in CO₂ equivalent, for crop T, Gg CO₂eq yr⁻¹
- Direct emissions $(N_2O)_{(T)}$ = Direct N_2O emissions from crop residues and Forage/Pasture renewal for crop T, $Gg N_2O yr^{-1}$
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T = Crop items

Step 10. Indirect emissions (N2O)

$$Indirect\ Emissions(N_2O)_{(T)} = F_{CR_{(T)}} \times Frac_{Leach} \times EF_5 \times \frac{44}{28} \times 10^{-6}$$

Where

- Indirect emissions $(N_2O)_{(T)}$ = Indirect N₂O emissions produced from atmospheric deposition of N, volatilised from crop residues and Forage/Pasture renewal, for crop T, Gg N₂O yr⁻¹
- $F_{CR(T)}$ = Annual amount of N in crop residues (above and below ground), including N-fixing crops, and from forage/pasture renewal, returned to soils annually for crop T, kg N yr⁻¹
- Frac_{Leach} = Fraction of N from crop residues that leaches as NH₃ and NO_Xm, kg N leached/kg of N additions (Table 27A)
- EF_5 = Emission factor for N₂O emissions from N leaching and runoff, kg N₂O-N/kg N leached (Table 25A)
- T = Crop items

Step 11. Indirect emissions (CO₂eq)

$$Indirect\ Emissions(CO_{2eq})_{(T)}=\ Indirect\ Emissions(N_2O)_{(T)}\ x\ GWP$$

- Indirect emissions $(CO_2eq)_{(T)}$ = Indirect N_2O emissions from crop residues and Forage/Pasture renewal in CO_2 equivalent for crop T, Gg CO_2eq yr^{-1}
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from crop residues and Forage/Pasture renewal for crop T, $Gg N_2O yr^{-1}$
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T = Crop items

Step 12. Emissions (N₂O)

$$Emissions(N_2O)_{(T)} = Direct \ Emissions(N_2O)_{(T)} + Indirect \ Emissions(N_2O)_{(T)}$$

Where

- $Emissions(N_2O)_{(T)}$ = Total N₂O emissions from crop residues and Forage/Pasture renewal, for crop T, Gg N₂O yr⁻¹
- Direct emissions $(N_2O)_{(T)}$ = Direct N_2O emissions from crop residues and Forage/Pasture renewal for crop T, $Gg N_2O yr^{-1}$
- Indirect emissions $(N_2O)_{(T)}$ = Indirect N_2O emissions from crop residues and Forage/Pasture renewal for crop T, $Gg N_2O yr^{-1}$
- T = Crop items

Step 13. Emissions (CO₂eq)

$$Emissions(CO_{2eq})_{(T)} = Emissions(N_2O)_{(T)} \times GWP$$

Where

- Emissions (CO₂eq)_(T) = Total N₂O emissions from crop residues and Forage/Pasture renewal in CO₂ equivalent for crop T, Gg CO₂eq yr⁻¹
- Emissions $(N_2O)_{(T)}$ = Total N₂O emissions from crop residues and Forage/Pasture renewal for crop T, Gg N₂O yr⁻¹
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- T = Crop items

Step 14. Implied emissions factor for N₂O

$$IEF_{(T)} = \frac{Emissions(N_2O)_{(T)}}{F_{CR(T)}} \times 10^6 \times \frac{28}{44}$$

- IEF(T) = Implied emission factor for crop T, kg N₂O /kg of N applied
- Emissions $(N_2O)_{(T)}$ = Total N₂O emissions from crop residues and Forage/Pasture renewal for crop T, Gg N₂O yr⁻¹
- $F_{CR(T)}$ = Annual amount of N in crop residues (above and below ground), including N-fixing crops, and from forage/pasture renewal, returned to soils annually for crop T, kg N yr⁻¹
- T = Crop items

5.1.8 Cultivation of organic soils

Dataset information	
Title	Cultivation of organic soils
Definition	Greenhouse gas (GHG) emissions data from cultivation of organic soils are those associated with nitrous oxide gas emissions, from drained histosols under cropland and grassland.

Methodology and quality information

Methods and processing

GHG emissions data from cultivation of organic soils are those associated with nitrous oxide gas from cultivated organic soils under cropland (item: Cropland organic soils) and grassland (item: Grassland organic soils). The FAOSTAT data are computed at Tier 1, following IPCC, 2006, Vol. 4, Ch. 11.

The emissions are estimated at pixel level, using the formula:

Emission = A * EF

where

- Emission = Annual emissions, in units of kg N₂O-N yr⁻¹;
- A = Activity data, representing the annual area of cultivated organic soils, in hectares (1).
- EF=Tier 1, default IPCC emission factors, expressed in units of kg N₂O-N ha⁻¹ (2).
- (1) Data are obtained through the stratification of three different global datasets: i.The Harmonized World Soil Database (FAO *et al.*, 2012), used to estimate the area covered by Histosols classes.
- ii. The Global Land Cover dataset, GLC2000 (EU-JRC, 2003), used to estimate the amount of cropland and grassland area in each pixel.

For cropland, three "cropland" classes from GLC2000 are used, in accordance with You et al. (2008):

CLASS	NAME	CROPLAND SHARE PER PIXEL
16	Cultivated and managed areas	100%
17	Mosaic: cropland/tree cover/Other natural vegetation	50%
18	Mosaic: cropland/Shrub and/or grass cover	10%

For grassland, two "herbaceous" and two mosaics GLC2000 classes are used, in line with the FAO Land Cover Classification Scheme (LCCS):

CLASS	NAME	GRASSLAND SHARE PER PIXEL
13	Herbaceous Cover, closed-open	100%
14	Sparse herbaceous or sparse shrub cover	50%
17	Mosaic: Cropland/Tree Cover/Other natural vegetation	25%
18	Mosaic: Cropland/Shrub and/or grass cover	45%

iii. The Gridded Livestock of the World for cattle and sheep (Wint and Robinson, 2007), used as an additional mask over grassland as a proxy to estimate drained area. With this mask, only those pixels with non-zero drained grassland histosols area and a livestock density > 1 head/ha are included.

For the period between 1990-present, the activity data reported in this sub-domain are a constant value, representing the year 2000, i.e. the reference year of the GLC2000 database.

(2) The EF values are those specified in IPCC, 2006, Vol. 4, Ch. 11, Table 11.1. The EF were assigned at the pixel level, to distinguish the relevant climate zone, as defined in IPCC, 2006, Vol. 4, Ch. 3, Annex 3A.5, using the climatic zones map from the Joint Research Centre of the European Commission (EC-JRC, 2010).

The analysis was carried out in GIS, combining the above datasets. The GHG estimates made at the pixel level were subsequently aggregated at country level, using the FAO Global Administrative Unit Layers (GAUL) data set.

Methodology and quality information

Methods and processing

The dimensionless conversion factors used are

- 44/28, to convert the emissions from kg N₂O-N to kg N₂O gas;
- 10^{-6} , to convert the emissions from kg N₂O to Gg N₂O; and
- GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22).

The "Cultivation of organic soils" sub-domain contains the following data categories, available for download: country-level GHG emissions in both Gg N_2O and Gg CO_2eq ; implied emission factors; and activity data.

Uncertainties in the estimates of GHG emissions are due to uncertainties in the emission factors and activity data. In the case of cultivation of organic soils, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol. 4, Ch. 11, Section 11.2.1.4).

Calculation procedure

Step 1. Activity data


Activity data are represented by the annual area of organic soils covered by cropland or grassland organic soils, in hectares. The $Area_{(T)}$ is calculated in a GIS for T = cropland and T = grassland.

The cropland area has been identified from the global land cover dataset, GLC2000 (EU-JRC, 2003) available at http://bioval.jrc.ec.europa.eu/products/glc2000/glc2000.php, and which uses the three "cropland" classes. The mosaic classes have been reduced, proportionally to the general share of the cropland per pixel for each class, according to the values reported in the metadata above.

The grassland area has been identified from the global land cover dataset, GLC2000 (EU-JRC, 2003), using four classes. The mosaic classes have been reduced, proportionally to the general share of the grassland per pixel for each class, according to the values reported in the metadata above.

The area of organic soils has been identified from the Harmonized World Soil Database, available at http://www.fao. org/nr/lman/abst/lman_080701_en.htm. All the pixels in which histosols are present are selected, with the relative pixel share indicated as a percentage.

The $Area_{(T)}$ was been calculated by overlaying the organic soils with the land cover layers. Only for T = Grassland, the Gridded Livestock of the World was used as a proxy to identify the areas of managed grasslands. The values are fixed, calculated for each country and accessible in the FAOSTAT Emissions database from the Emissions-Land Use/Grassland domain. The data concerns the years between 1990 and 2050 and may be retrieved as follows:



1.2 Projections of area

The area values for 2030 and 2050 are produced simply by replicating the country values for 2011.

Step 2. Emissions (N₂O-N)

The emissions are calculated in a GIS at the pixel level, using the formula:

Emissions
$$(N_2O - N)_{(T)} = Area_{(T)} \times EF_{(T)}$$

Where

- Emission $(N_2O-N)_{(T)}$ = Annual emissions of N₂O-N for item T, kg N₂O-N yr⁻¹
- $Area_{(T)}$ = Activity data, representing the area of organic soils under item T, ha
- $EF_{(T)}$ = Tier 1, default IPCC emission factors for item T, kg N₂O-N ha⁻¹ yr⁻¹ (Table 57A)

The EF values are specified in IPCC, 2006, Vol. 4, Ch. 11, Table 11.1. The EF were assigned at pixel level to the relevant climate zone, as defined in IPCC, 2006, Vol. 4, Ch. 3, Annex 3A.5. The climatic zones map used was developed by the Joint Research Centre of the European Commission (EC-JRC, 2010), following the IPCC's prescriptions.

The emissions are relate to the year 2000, which is the reference year of the land cover map used in the estimations. These values are also used for subsequent years, to cover the period between 1990 and 2011. Additional country-level disaggregations that have occurred over the period 1990-2011 are also provided.

As long as a country's boundary does not change, the area values for years following 2011 can be produced by simply replicating the country's 2011 values.

Step 3. Emissions (N₂O)

Emissions
$$(N_2O)_{(T)} = Emissions (N_2O - N)_{(T)} \times \frac{44}{28} \times 10^{-6}$$

Where:

- Emission $(N_2O)_{(T)}$ = Annual emissions of N_2O for item T, $Gg N_2O yr^{-1}$
- Emission $(N_2O-N)_{(T)}$ = Annual emissions of N_2O-N for item T, kg N_2O-N yr⁻¹

Step 4. Emissions (CO₂eq)

$$Emissions(CO_{2eq})_{(T)} = Emissions(N_2O)_{(T)} \times GWP$$

Where

- Emissions $(CO_2eq)_{(T)} = N_2O$ emissions from item T, in CO_2 equivalent, $Gg CO_2eq yr^{-1}$
- Emissions $(N_2O)_{(T)} = N_2O$ emissions from item T, Gg N₂O yr⁻¹
- GWP = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq

Step 5. Implied emissions factor for N₂O-N

$$IEF_{(T)} = \frac{Emissions(N_2O - N)_{(T)}}{Area_{(T)}}$$

Where

- $IEF_{(T)}$ = Implied emission factor for item T, kg N₂O-N ha⁻¹
- Emissions $(N_2O-N)_{(T)}$ = Total N_2O-N emissions from item T, kg N_2O-N yr⁻¹
- $Area_{(T)}$ = Area of Cultivated Organic Soils under item T

5.1.9 Burning crop residues

Dataset information		
Title	Burning crop residues	
Definition	Greenhouse Gas (GHG) emissions from burning crop residues consist of methane and nitrous oxide gases, produced by the combustion of a percentage of the crop residues burnt on-site.	
Methodology and qua	Methodology and quality information	
Methods and processing	Greenhouse Gas (GHG) emissions from burning crop residues consist of methane (CH ₄) and nitrous oxide (N ₂ O) gases produced by the combustion of a percentage of crop residues burnt on-site. The mass of fuel available for burning should be estimated, taking into account the fractions removed before burning due to animal consumption, decay in the field, and use in other sectors (e.g. biofuel, domestic livestock feed, building materials, <i>etc.</i>). The FAOSTAT data are estimated at Tier 1 in accordance with IPCC, 2006, Vol. 4, Chapters 2 and 5.	
	The CH_4 and N_2O emissions are estimated at country level, using the formula:	
	Emission = A * EF where	
	 Emission = GHG emissions in units of g CH₄ and g N₂O; 	
	• A = Activity data, representing the total amount of biomass burned, kg of dry matter (1);	
	 EF = Tier 1, default IPCC emission factors, expressed in g CH₄/kg of dry matter and g N₂O/kg of dry matter (2). 	

Methodology and quality information

Methods and processing

(1) Activity data are calculated from harvested area statistics and cover the following crops: wheat, maize, rice, and sugarcane. For the period between 1961-present, the harvested area data is taken from FAOSTAT (domain: Production/crops). Projections of harvested area for 2030 and 2050 are computed with respect to a baseline, defined as the 2005-2007 average of the corresponding FAOSTAT activity data, and by applying percentage growth rates from FAO perspective studies (Alexandratos and Bruinsma, 2012). The FAO projections used cover about 140 countries. Projections of activity data for countries that were not included assume the same growth rate experienced in neighbouring countries. The harvested area is used to estimate the amount of biomass burned, using mean default crop values of the mass of fuel available for combustion (MB) and the combustion factor (Cf) available in IPCC, 2006, Vol. 4, Ch. 2, Table 2.4. The mass is then reduced by the fraction of crop residue burnt on-site, assumed to be 10%, following IPCC, 2000, Ch.4, Section 4A.2.1.1.

(2) Global default EF values taken from IPCC, 2006, Vol. 4, Ch. 2, Table 2.5.

The dimensionless conversion factors used are

- 10⁻⁹, to convert the emissions from g CH₄ to Gg CH₄ and g N₂O to Gg N₂O;
- GWP- $CH_4 = 21$ (100-year time horizon global warming potential), to convert $Gg \ CH_4$ to $Gg \ CO_2eq$ and
- GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4 p. 22).

The burning crop residues domain contains the following data categories, available for download: country-level GHG emissions, provided as total, in Gg CH₄, Gg N₂O and Gg CO₂eg; implied emission factors; and activity data.

Uncertainties in the estimates of GHG emissions are due to uncertainties in the emission factors and activity data. In the case of burning crop residues, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol. 4, Ch. 5, Section 5.2.4.4).

Calculation procedure

Step 1. Activity data


In this sub-domain, activity data refers to the Area harvested of the following crop items: Maize, Paddy Rice, Sugar Cane and Wheat.

The definitions used by FAOSTAT for the term "Area harvested", and for individual items, are available at the Glossary, or on the FAOSTAT website at http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E. The unit of measurement is expressed in hectares ("Ha").

The data from 1961-present are disseminated from the FAOSTAT – Production, Crops sub-domain, as follows:

FIGURE 37

Area harvested data from the FAOSTAT database



1.2 Projections of area

The projections of activity data for 2030 and 2050 are computed with respect to a baseline provided by FAO perspective studies (Alexandratos and Bruinsma, 2012). This baseline may be computed as follows:

$$A_{(T)y} = \alpha_{(T)_y} \times A_{(T)_{2005-2007}}$$

Where

- $A_{(T)y}$ = Area for crop type T in the projected year y
- $\alpha_{(T)y}$ = Projection ratio for crop area A, in projected year y with respect to the baseline
- $A_{(T) 2005-2007}$ = Baseline area for crop type T computed as the average of the 2005, 2006 and 2007 area values
- T = Crop type
- y =The projected years, 2030 and 2050

Step 2. Area burned

$$AreaBurnt_{(T)} = Area_{(T)} \times Frac_{Burnt}$$

- AreaBurned_(T) = Annual area of crop T burned, ha yr⁻¹
- $Area_{(T)}$ = Total annual area harvested of crop T, ha yr⁻¹
- Frac_{Burned} = Fraction of burned area
- T = Crop category

Step 3. Biomass burned (dry matter)


$$BB_{(T)} = AreaBurned_{(T)} \times Fuel_{(T)}$$

Where

- $BB_{(T)}$ = Biomass burned for crop T, tonnes dry matter yr⁻¹
- AreaBurned_(T) = Annual area of crop T burned, ha yr⁻¹
- $Fuel_{(T)} = Mb \times Cf$, mass of fuel available x combustion factor (tonnes dry matter ha⁻¹) (Table 52A)
- T = Crop category.

The FAOSTAT Emissions database provides data relating to the biomass burned for the selected items:

FIGURE 38 Burning crops items



Step 4. Emissions (CH₄) and (N₂O)

IPCC 2006, Equation 2.27

$$Emissions(gas)_{(T)} = BB_{(T)} \times G_{ef(gas)} \times 10^{-6}$$

- $Emissions(gas)_{(T)}$ = Amount of emissions of Gas from crop residues burning for crop T, $Gg CO_2eq yr^{-1}$
- $BB_{(T)}$ = Biomass burned for crop T, tonnes yr⁻¹
- $G_{ef(gas)}$ = Emission factor g/kg dry matter burnt (Table 53A)
- $Gas = (CH_4, N_2O)$
- T = Crop category

Step 5. Emissions (CO₂eq) from CH₄ and N₂O

$$Emissions(CO_2eq)_{(T)} from (gas) = Emissions(gas)_{(T)} \times GWP_{(gas)}$$

Where

- $Emissions(CO_2eq)_{(T)}$ from (gas) = Amount of Gas emissions in CO_2eq from crop residues burning for crop T, $Gg CO_2eq yr^{-1}$
- Emissions $(gas)_{(T)}$ = Amount of Gas emissions from crop residues burning for crop T, Gg yr⁻¹
- $GWP_{(N2O)} = 310$ (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- $GWP_{(CH4)} = 21$ (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq
- T = Crop category
- $Gas = Gas (CH_4, N_2O)$

Step 6. Emissions (CO₂eq)

$$Emissions(CO_2eq)_{(T)} = Emissions(CO_2eq)_{(T)} from N_2O + Emissions(CO_2eq)_{(T)} from CH_4$$

Where

- $Emissions(CO_2eq)_{(T)}$ = Amount of emissions in CO₂eq from crop residues burning for crop T, Gg CO₂eq yr⁻¹
- $Emissions(CO_2eq)_{(T)}$ from N_2O = Amount of N_2O emissions in CO_2eq from crop residues burning for crop T, $Gg CO_2eq yr^{-1}$
- $Emissions(CO_2eq)_{(T)}$ from CH_4 = Amount of CH_4 emissions in CO_2eq from crop residues burning for crop T, $Gg CO_2eq yr^{-1}$
- T = Crop category.

Step 7. Implied emissions factors for N2O and CH4

$$IEF_{(T,gas)} = \frac{Emissions(gas)_{(T)}}{BB_{(T)}} \times 10^6$$

- $IEF_{(T, gas)}$ = Implied emission factor for Gas for crop T, g/kg applied
- $Emissions(gas)_{(T)} = Amount of Gas emissions from crop residues burning for crop T, Gg yr⁻¹$
- $BB_{(T)}$ = Biomass burned for crop T, tonnes yr⁻¹
- T = Crop category
- $Gas = Gas (CH_4, N_2O)$

5.1.10 Burning-Savanna

Dataset information		
Γitle	Burning-Savanna	
Definition	Greenhouse Gas (GHG) emissions from burning of savanna consist of methane and nitrous oxide gases from biomass combustion.	
Methodology and o	quality information	
Methods and processing	Greenhouse Gas (GHG) emissions from burning of savanna consist of methane (CH ₄) and nitrous oxide (N ₂ O) gases produced from the burning of vegetation biomass in the following five land cover types: Savanna, Woody Savanna, Open Shrublands, Closed Shrublands, and Grasslands. The FAOSTAT data are estimated at Tier 1, in accordance with IPCC, 2006, Vol. 4, Ch. 2, Equation 2.27.	
	CH_4 and N_2O emissions are estimated on a spatial grid, at a 0.25 degree resolution (approximately 25 km at the Equator), using the formula:	
	Emission = A * EF where	
	 Emission = GHG emissions in g CH₄ and g N₂O; 	
	 A = Activity data, representing the total mass of fuel burned in each pixel, kg of dimatter (1); 	
	 EF = Tier 1 IPCC emission factor, expressed in g CH₄ or g N₂O per kg of burned dimatter (2). 	
	(1) Total mass of fuel burned is computed by multiplying the burned area (i) by the fue biomass consumption values (ii).	
	(i) Yearly composite burned area values are produced from monthly statistics of the Global Fire Emission Database v.4, based on MODIS remote-sensing data (see GFED4, Giglio <i>et al.</i> , 2013). The data set provides data on burned area by land cover classes, as identified by the MODIS Land Cover product (MCD12Ω1, Hansen et al., 2000).	
	(ii) Fuel biomass consumption values were assigned at pixel level to the relevant climate zone, as defined in IPCC, 2006, Vol. 4, Ch. 2, Table 2.4. The climatic zones map used was developed by the Joint Research Centre of the European Commission (EC-JRC, 2010), and follow IPCC prescriptions.	
	The data period available on FAOSTAT is 1990-present, with annual updates, and projections to 2030 and 2050. Since GFED4 data are not available before 1996 and for future periods, yearly values for the period between 1990 and 1995, as well as for 2030 and 2050, are set as a constant, and estimated as the average of the period between 1996 and 2012.	
	(2) Default EF values are taken from IPCC, 2006, Vol. 4, Ch. 2, Table 2.5.	
	GHG estimates made at pixel level were subsequently aggregated at the country level using the FAO Global Administrative Unit Layers (GAUL) data set.	
	The dimensionless conversion factors used are	
	 10⁻⁹, to convert the emissions from g CH₄ to Gg CH₄ and g N₂O to Gg N₂O; GWP- CH₄ = 21 (100-year time horizon global warming potential), to convert Gg Ch to Gg CO₂eq GWP-N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂ to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4 p. 22). 	
	The Burning-Savanna sub-domain contains the following data categories available for download, for each land cover class: country-level GHG emissions, provided as totals in Gg CH ₄ , Gg N ₂ O and Gg CO ₂ eq; implied emission factor; burned area; and tota mass of fuel available. Data is available for all countries and territories with an area large enough to be monitored at the resolution of GFED4 data, as well as for standard FAOSTAT regional aggregations, plus Annex I and non-Annex I groups.	
	Uncertainties in the estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of biomass burning, more detailed information is available in the IPCC guidelines (IPCC, 2006, Vol. 4, Ch. 5, Section 5.2.4.4).	

Calculation procedure

Step 1. Activity data

The Burning-Savanna domain considers emissions from fires in the following land cover classes: "Savanna", "Woody Savanna", "Closed Shrublands", "Open Shrublands" and "Grasslands". The GHG considered in this domain are CH_4 and N_2O .

The annual burned area for each land cover is derived from the Global Fire Emission Database burned area data set, version 4 (GFED4). This is available at http://www.globalfiredata.org/


GFED4 provides global monthly burned area maps at 0.25° spatial resolution, from mid-1995 to present (the most recent year currently available is 2012; see Giglio *et al.* 2013). The GFED4 dataset combines a 500m Moderate Resolution Imaging Spectro-radiometer (MODIS) burned area data set with active fire data sets from the Tropical Rainfall Measuring Mission (TRMM) Visible and Infrared Scanner (VIRS), and the Along-Track Scanning Radiometer (ATSR) family of sensors. The annual burned area for each land cover class is calculated by adding up the GFED4 monthly data by country.

The Biomass Burned (BB) activity data for each land cover class, as well as the resulting emissions of N_2O and CH_4 , are estimated using the IPCC 2006 default methodology. This is based on Equation 2.27, with default parameters for biomass consumption per hectare in different climates and vegetation types provided by Table 2.4, and emission factors for each vegetation type taken from Table 2.5. The biomass burned and emission computations are performed at pixel level in a GIS environment, using additional data sets to associate the correct IPCC parameters to the pixel's specific climate and vegetation characterization.

The methodology set out below enables the calculation of annual GHG emissions for the period between 1990-1995, based on averages for the period between 1996-present; the conversion of non-CO₂ emissions into CO₂ equivalents; and the calculation of implied emission factors for N₂O and CH₄.

The calculated data are available in the FAOSTAT Emissions database, under the Emission-Agriculture/Burning Savanna domain, as shown below:





1.2 Projections of area

Since GFED4 data are not available for years before 1996 and, obviously, for future periods, yearly values for the period 1990-1995, as well as for 2030 and 2050, are set as a constant, and estimated as the average of the period between 1996 and 2012.

Step 2. Production of 1990-1995 data

For each country, yearly values for the period 1990-1995, X_i , are to be estimated, as the average value over the period between 1996-present:

$$X_i = \overline{X_{1996-present}}$$

Where

- X = Variable (Burned Area, Biomass Burned, Emissions)
- $X_{(1996-present)}$ = Average value for the years 1996-present
- i = Year (period between 1990-1995)

Step 3. Emissions (CO₂eq) from CH₄ and N₂O

$$Emissions(CO_2eq)_{(T)} from (gas) = Emissions(gas)_{(T)} \times GWP_{(gas)}$$

Where

- Emissions $(CO_2eq)_{(T)}$ from (gas) = Amount of GHG emissions in CO_2eq from biomass burning for land cover T, Gg CO_2eq vr^{-1}
- $Emissions(gas)_{(T)} = Amount of Gas emissions from biomass burning for crop T, Gg yr⁻¹$
- $GWP_{(N2O)} = 310$ (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- $GWP_{(CH4)} = 21$ (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq
- T = Land cover class
- *Gas*= CH₄, N₂O

Step 4. Emissions (CO₂eq)

$$Emissions(CO_2eq)_{(T)} = Emissions(CO_2eq)_{(T)} from N_2O + Emissions(CO_2eq)_{(T)} from CH_4$$

Where

- Emissions(CO_2eq)_(T) = Amount of emissions in CO_2eq from biomass burning, for land cover T, Gg CO_2eq yr⁻¹
- $Emissions(CO_2eq)_{(T)}$ from N_2O = Amount of N_2O emissions in CO_2eq from biomass burning for crop T, Gg CO_2eq yr⁻¹
- Emissions(CO₂eq)_(T) from CH₄ = Amount of CH₄ emissions in CO₂eq from biomass burning for crop T, Gg CO₂eq yr⁻¹
- T = land cover class

Step 5. Implied emission factor for N2O and CH4

$$IEF_{(T,gas)} = \frac{Emissions(gas)_{(T)}}{BB_{(T)}} \times 10^6$$

- $IEF_{(T, gas)}$ = Implied emission factor for Gas for land cover T, $g/kg yr^{-1}burnt$
- $Emissions(gas)_{(T)} = Amount of Gas emissions from biomass burning for land cover T, Gg yr⁻¹$
- $BB_{(T)}$ = Biomass burned for land cover T, tonnes yr⁻¹
- T = Land cover class
- $Gas = Gas (CH_4, N_2O)$

5.1.11 Energy use

Dataset information	Dataset information	
Title	Energy use	
Definition	Greenhouse gas (GHG) emissions from direct energy use consist of carbon dioxide, methane and nitrous oxide gases associated with fuel burning and the generation of electricity used in agriculture (including fisheries).	
Methodology and	quality information	
Methods and processing	GHG emissions from direct energy use consist of carbon dioxide (CO_2), methane (CH_4) and nitrous oxide (N_2O) gases associated with fuel burning and generation of electricity used in agriculture (including fisheries). The FAOSTAT data are computed at Tier 1, following the criteria established in IPCC 2006, Vol. 2, Ch. 2 and 3, with the exception of the country electricity-only emission factors (see below).	
	Data are estimated for each main energy carrier (fuels and electricity) used in agriculture, for the following carriers: Motor gasoline, Gas-Diesel oil, Natural gas, Residual fuel oil, Liquefied petroleum gas, Coal and Electricity, and for selected agricultural sub-domains (i.e. irrigation and fisheries). Renewable energy for electricity indirectly contributes to the GHG emission calculation, via country emission factors, which depend on a country's chosen energy mix for electricity generation.	
	The emissions are estimated at country level, using the formula:	
	Emission = A * EF where	
	 A = Activity data, representing the amount of fuel consumed in Kt (1000 metric tons) yr¹, TJ (terajoules) yr¹ or ktoe (k tonnes of oil equivalent) yr¹ for fuels, and GWh yr¹ for electricity (1); 	
	 EF=Tier 1, default IPCC emission factors, expressed as kg of gas emitted per terajoule (for fuels) or gram per kilowatt hour (for electricity) of energy used per year (2). 	
	(1) The activity data relating to fuels are taken from the United Nations Statistics Division (UNSD) Energy Statistics database; the data relating to fuels used in fisheries, specifically "Gas-Diesel oil used in fisheries" and "Residual fuel oil used in fisheries" are taken from the International Energy Agency (IEA) database of "World energy balances"; for energy used for power irrigation, the irrigated area is taken from FAO AQUASTAT (*). The activity data (**) cover the following:	
	 a. Gas-Diesel oil (including fisheries); b. Motor gasoline; c. Natural gas (including Liquefied Natural Gas); d. Liquefied petroleum gas; e. Residual fuel oil (including fisheries); f. Coal (other bituminous coal or, if not available, Hard coal); g. Electricity; h. Gas-diesel oil used in fisheries; i. Residual fuel oil used in fisheries; 	
	i. Residual fuel oil used in fisheries; j. Energy for power irrigation.	
	(*)The irrigated area considered is the sum of the "Total area equipped for full control irrigation: sprinkler" and the "Total area equipped for full control irrigation: localized", values taken from the FAO AQUASTAT database. Annual values in FAOSTAT are estimated by repeating the last-available value reported in AQUASTAT within the relevant time interval. Regional factors indicating the energy required to irrigate one hectare of land (Stout, 1990) have been applied to estimate the energy consumed for irrigation. Specifically, for Europe, Northern America and Oceania, this is 1,929 kWh/ha; for Africa and Asia, 2,411 kWh/ha; for Central America, South America and the Caribbean, 2,170 kWh/ha.	
	(**) For items a, b, d, e and f, the amount of energy from the mass of fuels was derived using the Net Calorific Values reported in IPCC 2006, Vol. 2, Ch. 1, Table 1.2;	
	(2) The EF values are those reported by energy carrier in IPCC 2006, Vol. 2, Ch. 2, Table 2.5, and Ch. 3, Table 3.31. A summary table is shown below:	

Methodology and quality information

Methods and processing

Energy carrier	Use	CO ₂	CH₄ kg/TJ	N ₂ O
Gas/Diesel oil	Mobile	74,100	4.15	28.6
Motor gasoline	Mobile	69,300	80	2
Natural gas liquids	Stationary	64,200	10	0.6
Residual fuel oil	Stationary	77,400	10	0.6
Liquefied petroleum gases	Stationary	63,100	5	0.1
Coal (Other Bituminous Coal)	Stationary	94,600	300	1.5

For electricity, country emission factors for CO2, namely electricity-only factors, are those reported by the IEA (OECD/IEA, 2012). These are based on 1996 IPCC Guidelines for National Greenhouse Gas Inventories; the 1990 country emission factors are applied to previous years. Non-CO2 emission factors for electricity used for all countries are estimated on the basis of the IEA global electricity emission factor for CO2, considering these emissions as a fraction of the CO2 emitted (CH4: 300/98,300 for coal, 5/56,100 for natural gas, 10/73,300 for crude oil; N2O: 1.5/98,300 for coal, 0.1/56,100 for natural gas, and 0.6/73,300 for crude oil). The relevant energy mix for electricity is: 48% coal, 22% natural gas, and 6% crude oil.

The dimensionless conversion factors used are

- 10⁻⁶, to convert kg to Gg; and
- GWP-CH₄= 21; GWP-N₂O= 310 (100-year time horizon global warming potential), to convert Gg CH₄ or Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22);
- 1 GWh = 3.6TJ

The energy use domain contains the following data categories, available for download: country-level GHG emissions, provided as a total, in Gg CH_4 , Gg N_2O and Gg CO_2eq ; implied emission factors; and activity data.

Calculation procedure

Step 1. Activity data

The following items are included as activity data: Gas-diesel oils, motor gasoline, natural gas (including LNG), liquefied petroleum gas (LPG), residual fuel oil, coal, electricity, gas - diesel oils in fisheries, residual fuel oil in fisheries, and energy for power irrigation. Their definitions are available in the Glossary.

The activity data for gas-diesel oils, motor gasoline, natural gas (including LNG), liquefied petroleum gas (LPG), residual fuel oil, coal (hard coal or other bituminous coal), and electricity, are taken from the UNSD energy database, which is publicly available (for data after 1990) at the UN DATA portal (http://data.un.org/Explorer.aspx?d=EDATA).


Data relating to *energy consumed in fisheries*, specifically "Gas-diesel oils used in fisheries" and "Residual fuel oil used in fisheries", are taken from the IEA Statistics 2013 (http://www.iea.org/statistics/).

For *Energy for power irrigation*, the source of the activity data is the sum of the values for the "Area equipped for full control irrigation: sprinkler" and for the "Area equipped for full control irrigation: localized" that can be found in the Glossary and on the FAO AQUASTAT database website, at http://www.fao.org/nr/water/aquastat/data/query/index.html.

Figure 40 below shows how to access to the data of the "Area equipped for full control irrigation", both sprinkler and localized.

FIGURE 40

Data on the area equipped for power irrigation from the AQUASTAT database



Step 2. Calculate consumption in agriculture

a) For the Gas-Diesel oil, Gasoline, Liquefied petroleum gas (LPG), Residual fuel oil, and Coal (Hard coal or Other bituminous coal) items, consumptions are converted to TJ by applying the parameters reported in (Table 54A) in Annex 3, and the formula:

$$CA_{(F)}(Tj) = MF(Kt)x NCV_{(F)}$$

Where

- $CA_{(F)}$ = Consumption of fuel F in agriculture (TJ)
- MF = Mass of fuel consumed (1000 metric tons)
- $NCV_{(F)}$ = Default net calorific value of fuel F (Table 54A) and consistent with IPCC 2006 Guidelines)
- F = Fuel

Natural gas (including LNG) is provided in TJ from the original source. Residual fuel oil and Gas-Diesel oil used in fisheries are provided in Ktoe from the original source and converted to TJ (1 Ktoe = 41.868 TJ). Electricity consumption is provided in GWh from the original source and converted to TJ (1 GWh = 3.6 TJ).

2.2 For the "Energy for power irrigation" item, the following formula was applied:

$$CA_{(el)}(KWh) = Api(ha)xTC$$

Where

- $CA_{(el)}$ = Consumption of power irrigation in agriculture (KWh)
- Api = Total area equipped for full control irrigation: sprinkler and localized (ha)
- TC = Technical coefficient indicating the energy needed to irrigate one hectare of land (adapted from Stout, 1990) as follows: Europe, Northern America, Oceania 1,929 kWh/ha; Africa, Asia 2,411 kWh/ha; Central America, South America, Caribbean 2,170 kWh/ha

FAO's regional groups were applied.

Step 3. Calculating Emissions of CH₄, N₂O and CO₂

a) For the Gas-Diesel oils, Gasoline, Natural Gas (including LNG), Liquefied petroleum gas (LPG), Coal, Residual fuel oil, Gas-Diesel oils in fisheries, and Residual fuel oil in fisheries items, the formula is:

$$Emissions(gas)_{(F)} = CA_{(F)} \times G_{ef(F,gas)} \times 10^{-6}$$

Where

- $Emissions(gas)_{(F)}$ = Amount of emissions of Gas from energy use in agriculture of energy carrier F (Gg yr⁻¹)
- $CA_{(F)}$ = Energy consumption in agriculture (TJ yr⁻¹)
- $G_{ef(gas)}$ = Implied emission factor (kg TJ⁻¹)
- $Gas = (CO_2, CH_4, N_2O)$
- F =Energy carrier

The emission factors used are specified by energy carrier, in IPCC 2006, Vol. 2, Ch. 2, Table 2.5, and Ch. 3, Table 3.31; they are reported in Table 54A in Annex 3 to this publication. For Coal, the default NCV and emission factor of "Other bituminous coal" is applied.

b) For the Electricity and Energy for power irrigation items, the formula used was:

$$Emissions(gas)_{(el)} = CA_{(el)} \times G_{ef(el)} \times 10^{-3}$$

Where

- $Emissions(gas)_{(el)}$ = Amount of emissions of Gas from electricity use in agriculture, Gg yr⁻¹)
- CA_(el) = Electricity consumption in agriculture, GWh yr⁻¹
- $G_{ef(el)}$ = Country emission factor (electricity-only) of country C, g KWh⁻¹
- *Gas*= (CO₂, CH₄, N₂O)

For electricity, country emission factors for CO_2 – namely, electricity-only factors – are those reported by the IEA (OECD/IEA, 2012). These are based on the 1996 IPCC Guidelines for National Greenhouse Gas Inventories; the 1990 country emission factors are applied to previous years. Non- CO_2 emission factors for electricity used for all countries are estimated on the basis of the IEA global electricity emission factor for CO_2 , considering these emissions as a fraction of the CO_2 emitted⁸, and considering the following energy mix for electricity: 48% coal, 22% natural gas, and 6% crude oil.

Step 4. Calculate Emissions (CO₂eq) from CH₄ and N₂O

The relevant formula was:

$$Emissions(CO_2eq)_{(T)} from (gas) = Emissions(gas)_{(T)} \times GWP_{(gas)}$$

Where

- Emissions(CO₂eq) from (gas) = Amount of GHG emissions, in CO₂eq, from energy use in agriculture, Gg CO₂eq yr⁻¹
- $Emissions(gas)_{(F)}$ = Amount of Gas emissions from energy use in agriculture for energy carrier F, Gg yr⁻¹
- $GWP_{(CO2)} = 1$
- $GWP_{(CH4)} = 21$ (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq

⁸ CH₄: 300/98,300 for coal, 5/56,100 for natural gas, 10/73,300 for crude oil; N₂O: 1.5/98,300 for coal, 0.1/56,100 for natural gas, 0.6/73,300 for crude oil.

- $GWP_{(N2O)} = 310$ (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- F =Energy carrier
- $Gas = CH_4, N_2O$

Step 5. Calculate Overall CO2eq Emissions

The relevant formula is:

 $Emissions(CO_2eq)_{(F)} = Emissions\left(CO_2(F) + Emissions(CO_2eq)_{(F)} from \ CH_4 + \ Emissions(CO_2eq)_{(F)} from \ N_2O_2(F) + CO_2(F) from \ N_2O_2(F)$

Where

- Emissions $(CO_2eq)_{(F)}$ = Amount of emissions in CO_2eq from energy use for energy carrier F, Gg CO_2eq yr⁻¹
- Emissions $(CO_2)_{(F)}$ = Amount of CO_2 emissions from energy use for energy carrier F, $Gg CO_2eq yr^{-1}$
- Emissions $(CO_2eq)_{(F)}$ from N_2O = Amount of N_2O emissions in CO_2eq from energy use for energy carrier F, $Gg CO_2eq yr^{-1}$
- Emissions $(CO_2eq)_{(F)}$ from CH_4 = Amount of CH_4 emissions in CO_2eq from energy use for energy carrier F, $Gg CO_2eq yr^{-1}$
- F =Energy carrier

The items are also provided as the following groups:

Total energy = Gas-diesel oils + Motor gasoline + Natural gas (including LNG) + Liquefied petroleum gas (LPG) + Residual fuel oil + Coal + Electricity.

Transport fuel consumed in agriculture (excluding fisheries) = Gas-Diesel oil + Motor gasoline - Gas-Diesel oil in fisheries.

Energy consumed in fisheries = Gas-Diesel oil in fisheries + Residual fuel oil in fisheries.

5.2 FORESTRY AND LAND USE

5.2.1 Forest land

Dataset information	
Title	Forest land
Definition	The annual net CO ₂ emissions/removals from Forest land consist of the net carbon stock gain/loss in the living biomass pool (aboveground and belowground biomass) associated with <i>Forest</i> and <i>Net Forest Conversion</i> .
Methodology and qua	lity information
Methods and processing	The net CO ₂ emissions/removals from Forest land consist of the net carbon stock change in the living biomass pool (aboveground and belowground) associated with: i) Forest, referring to C stock changes occurring on Forest land in the reported year; and ii) Net Forest conversion from Forest land to Other land uses. The FAOSTAT data are computed at Tier 1, with the stock difference method, following the criteria established in IPCC 2006, Vol. 4, Ch.s 2 and 4.
	The net CO_2 emissions/removals (E/R) are estimated at country level, using the formula:
	E/R = A * CSCF * -44/12 / 1,000
	 where E/R = Net CO₂ emission/removal, in Gg CO₂ yr¹ (3);
	 A = Activity data, representing the forest area under forest management or the forest
	area net change, in ha (1);
	 CSCF = per-hectare carbon stock change in the living biomass pool (aboveground + belowground) of Forest land, expressed in units of t C/ha (2);
	(1) The area of forest, A, is calculated from the annual areas of Forest land taken directly from the Global Forest Resource Assessment (FRA) conducted by FAO (available at http://www.fao.org/forestry/fra/fra2010/en/). Data for the years 1990, 2000, 2005 and 2010, as provided by FRA, for the categories of <i>Primary forest, Other naturally regenerated forest</i> and <i>Planted forest</i> were linearly interpolated to compile, for each country, complete time series of areas for each category, for the period between 1990-2010. The FRA categories of <i>Primary forest</i> and <i>Other naturally regenerated forest</i> were aggregated, while <i>Planted forest</i> was considered separately, to compute the following forest area components at year t:
	 a. Area of forest that was still forest in the previous year (area type SFA), computed as Min [A(t), A(t-1)]; b. For Forest land (Forest Management): the new net area converted to forest in the same year (area type NAD), computed as: Max[A(t)-A(t-1),0], thus including only positive net forest area change; or c. For Net Forest Conversion: the net area loss converted from forest to Other land uses (area type NAD), computed as: Min[A(t)-A(t-1),0]], thus including only positive net forest area change.
	(2) CSCF is computed from the per-hectare carbon stock in the living biomass (aboveground + belowground) pool in the country in year t , $b(t)$. The latter is obtained from data on per-hectare carbon stocks taken directly from the FRA database for the years 1990, 2000, 2005 and 2010. These were linearly interpolated to compile, for each country, a complete time series of per-hectare average carbon stock in the living biomass pool, $b(t)$, for the period between 1990-2010. For countries for which FRA carbon stock data were not available, the relevant FRA regional carbon stock (FRA 2010, Table T2.21) was applied.
	For each year t, and each forest area type above, the CSCF is calculated as follows:
	For Forest:
	i. $CSCF(t, SFA) = \Delta b(t) = b(t) - b(t-1)$, for SFA-type forest areas; ii. $CSCF*(t, NAD) = b(t)$, for NAD-type forest areas.
	The overall net carbon stock change factor at year t , $CSCF(t)$, is computed as: $CSCF(t) = [CSCF(t,SFA)*SFA + CSCF(t,NAD)*NAD]/A$
	For Net Forest Conversion:
	i. $CSCF*(t, NAD) = b \ (t-1)$, for NAD-type forest areas

Methodology and quality information

Methods and processing

The overall net carbon stock change factor at year t, CSCF(t), is computed as: CSCF(t) = CSCF(t, NAD)*NAD/A

The dimensionless conversion factors used were

- -44/12, to convert from carbon mass to CO₂ emissions;
- 10⁻³, to convert tonnes into Gg.

The Forest land sub-domain contains the following data, available for download: country-level Net CO₂ emissions/removals in Gg CO₂, and Carbon stock change in Gg C; implied emission factors (i.e. the CSCFs); and underlying activity data. Data are available for all individual countries and territories, as well as for standard FAOSTAT regional aggregations, and Annex I and non-Annex I groups. The data period covers 1990-present, with periodic updates in line with FRA releases.

For estimating uncertainty, it should be noted that information on growing stock and basic wood densities is typically used to compute living biomass carbon stocks. FRA 2005 estimates uncertainties in growing stock at $\pm 8\%$ for industrialized countries and $\pm 30\%$ for non-industrialized countries, and uncertainties for basic wood density around 10 to 40%. FAOSTAT estimates area uncertainties at $\pm 10\%$.

Calculation procedure


Step 1. Activity data

In this sub-domain, the activity data are taken from the "Global Forest Resources Assessments" website, available at http://www.fao.org/forestry/fra/en/. The website includes datasets for 1990, 2000, 2005 and 2010 (FRA 2010), by item and country.

The FRA is the most comprehensive assessment of forests and forestry to date, in terms of the number of countries and individuals involved. It examines the current status and recent trends for about 90 variables, covering the extent, condition, uses and values of forests and other wooded land, with the aim of assessing all benefits that may be derived from forest resources. All methodological documentation referenced in this section is available at the Global Forest Resources Assessments website (http://www.fao.org/forestry/fra/67094/en/).

The FRA 2010 database may be accessed at http://www.fao.org/forestry/fra/fra2010/en/.

FIGURE 41
The "Global Forest Resources Assessment 2010" homepage




FRA data are distributed according to the following categories:

- Primary forest (ha) (PA)
- Secondary forest (ha) (SA)
- Planted forest (ha) (PLA)
- Carbon density of living biomass (tC/ha) (CD)


1.1 Data for the **Primary forest**, secondary forest (other naturally regenerated forest), and planted forest categories are accessible as follows:






However, the FAOSTAT Emissions database uses the complete time series (1990-2011) of total forest area which is available in FAOSTAT database (Inputs Land sub domain http://faostat3.fao.org/faostat-gateway/go/to/download/R/RL/E) and derived from FRA data.

FIGURE 43 Total forest area in FAOSTAT



1.2 The data relating to the **Carbon density of living biomass** are accessible as follows:

FIGURE 44 Carbon stock dataset from the FRA database 1) Click on "Volume, biomass and carbon" ABOUT KEY INDICATORS METADATA NEWS Volume, biomass and carbon Extent and characteristics Ownership and management ,O csv Forest establishment Volume, biomass and carbon Growing stock (Million m3 over bark) Forest health T06F0000 by Forest/Other wooded land, FRA category **Employment and human resources** Biomass stock (Million metric tonnes) by Forest/Other wooded land, FRA categories, Year, Country **Education and research** T07F0000 Revenues and expenditures Carbon stock (Million metric tonnes) All T08F0000 2) Click on "Carbon stock"



FRA data are based on the country reports compiled by officially appointed country correspondents, in collaboration with FAO staff. Prior to finalization, these reports are subject to validation by the forestry authorities of the relevant countries.

The official national reports are available at the following link http://www.fao.org/forestry/fra/67090/en/.

The FRA data were used to compute the following items and components:

Items	Net Forest Conversion	
Components	Net Area Difference (ha)	NAD
	• Carbon Stock Change (Gg C)	CSC
	• CO ₂ Emissions (Gg CO ₂)	CEM
Items	• Forest	
Components	• Steady Forest Area (ha)	SFA
	Net Area Difference (ha)	NAD
	•Total Forest Area (ha)	TFA
	Carbon Stock Change (Gg C)	CSC
	• Net CO ₂ Emissions/Removals (Gg CO ₂)	NER

The methodology is illustrated below.

Step 2. Net Forest Conversion

2.1 Calculation of Net Area Difference (NAD)

• An area difference based on primary and secondary forest areas must be computed:

$$\begin{split} NAD^{PS}_{\ (i,j)} &= Min \ [\ PA_{(i,j)} + SA_{(i,j)} - PA_{(i-1,j)} - SA_{(i-1,j)}, 0] \\ for \ i &= 1991, \dots, 2010; \ j = country \ j \\ with \ NAD^{PS}_{\ } (1990,j) &= NAD^{PS}_{\ } (1991,j) \end{split}$$

An area difference based on planted areas only must then be computed:

NADPLA
$$_{(i,j)}$$
 = Min [PLA $_{(i,j)}$ - PLA $_{(i-1,j)}$, 0]
for i =1991, ..., 2010; j = country j
with NADPLA (1990, j) = NADPLA (1991, j)

• Finally, compute the sum of these two values:

$$NAD_{(i,j)} = NADPS_{(i,j)} + NADPLA_{(i,j)}$$
 for $i = 1990,...,2010$; $j = country j$

Note: NAD, NAD^{PS} and NAD^{PLA} are always <= 0

2.2 Calculation of Carbon Stock Change (CSC)

• This is simply computed as

$$CSC_{(i,j)} = NAD_{(i,j)} \times CD_{(i-1,j)}$$

for i =1991, ..., 2010; j = country j

Where

• $CD_{(i-1,j)} = Carbon Density in Living Biomass in the previous year, from the parameter file.$

Note: in this case, CSC is always negative or zero.

2.3 Calculation of CO₂ Emissions, CEM [CO₂, xx] [CO₂eq, xx]

• This is computed simply, as:

```
CEM_{(i,j)} = -CSC_{(i,j)} \times 44/12
for i =1990, ..., 2010; j = country j
```

Note: The CEM amount in CO₂eq is equal to the NER in CO₂.

Step 3. Forest land

3.1 Calculation of Net Area Difference (NAD)

• An area difference based on primary and secondary forest areas must be computed:

$$\begin{split} &NAD^{PS}_{\ (i,j)} = Max \ [\ PA_{(i,j)} + SA_{(i,j)} - PA_{(i-1,j)} - SA_{(i-1,j)}, 0] \\ &for \ i = 1991, \dots, 2010; \ j = country \ j \\ &with \ NAD^{PS} \ (1990,j) = NAD^{PS} \ (1991,j) \end{split}$$

• An area difference based on planted areas only must be computed:

$$NAD^{PLA}_{(i,j)} = Max [PLA_{(i,j)} - PLA_{(i-1,j)}, 0]$$

for i =1991, ..., 2010; j = country j
with NAD^{PLA} (1990, j) = NAD^{PLA} (1991, j)

• The total must then be calculated:

$$NAD_{(i,j)} = NAD^{PS}_{(i,j)} + NAD^{PLA}_{(i,j)}$$

for i =1990, ..., 2010; j = country j

Note: NAD, NAD^{PS} and NAD^{PLA} are always ≥ 0

Step 4. Calculation of Steady Forest Area (SFA)

• First, compute the steady forest area of primary and secondary forests:

```
\begin{split} SFA^{PS}_{\ (i,j)} &= Min \ [ \ PA_{(i,j)} + SA_{(i,j)}, PA_{(i-1,j)} + SA_{(i-1,j)} \ ] \\ for \ i &= 1991, \dots, 2010; \ j = country \ j \\ With \ SFA^{PS}_{\ } (1990,j) &= SFA^{PS}_{\ } (1991,j) \end{split}
```

Then, compute an area difference based on planted areas only:

$$\begin{split} & SFA^{PLA}{}_{(i,j)} = Max \; [PLA_{(i,j)}, PLA_{(i-1,j)}] \\ & for \; i = 1991, \dots, 2010; \; j = country \; j \\ & With \; SFA^{PLA}(1990,j) = SFA^{PLA}(1991,j) \end{split}$$

• Finally, compute the total:

$$SFA_{(i,j)} = SFA^{PS}_{(i,j)} + SFA^{PLA}_{(i,j)}$$

for i =1990, ..., 2010; j = country j

Step 5. Calculation of Total Forest Area (TFA)

• This may be simply computed, as:

$$TFA_{(i,j)} = NAD_{(i,j)} + SFA_{(i,j)}$$
 for $i = 1990, ..., 2010$; $j = country j$

Step 6. Calculation of Carbon Stock Change (NCSC)

• The following formula must be applied for $NAD_{(i,j)}$:

$$CSC^{NAD}_{(i,j)} = NAD_{(i,j)} \times CD_{(i,j)}$$

for i =1990, ..., 2010; j = country j

Where

 $CD_{(i,j)}$ = Carbon Density in Living Biomass, from the parameter files of the Average Carbon Stock in Forest Living Biomass (t ha⁻¹), available in the FRA 2010, by geographical group area and year (Table 55A), and of the Total Carbon Stock in Forest Living Biomass (M t), available in the FRA 2010 by Country (Table 56A).

Note: CSC^{NAD} is always positive or zero in this case.

• The following must be computed for SFA_(i,j):

$$CSC^{SFA}_{(i,j)} = [CD_{(i,j)} - CD_{(i-1,j)}] * SFA_{(i,j)}$$

Where

$$CSC(1990,j) = CSC(1991,j);$$

 $CD_{(i,j)}$ = Carbon Density in Living Biomass, from the parameter of the Average Carbon Stock in Forest Living Biomass (t ha-1), available in the FRA 2010 by geographical group area and year (Table 55A), and the Total Carbon Stock in Forest Living Biomass (M t), available in the FRA 2010 by Country (Table 56A).

• Finally, compute the total:

$$\begin{aligned} &CSC_{(i,j)} = CSC^{NAD}_{\ (i,j)} + CSC^{SFA}_{\ (i,j)} \\ &for\ i = 1990, \dots, 2010; \ j = country\ j \end{aligned}$$

Step 7. Net CO₂ Emissions/Removals, NER [CO₂, xx] [CO₂eq, xx])

• This may be simply computed as:

Note: The NER amount in CO_2 eq is equal to the NER in CO_2 .

5.2.2 Cropland

Dataset informatio	11			
Title	Cropland			
Definition	emission	use gas (GHG) emissions data from cro s from cropland organic soils. They are ass nistosols under cropland.		
Methodology and	quality informa	tion		
Methods and processing	organic s	ssions data from cropland are currently limicals. They are associated with carbon lossed data are computed at Tier 1, following the case.	s from drained organic soils. The	
	The emis	sions are estimated at the pixel level, using	the formula:	
		Emission = A * EF		
	where			
	• Emissi	on = Annual emissions, in units of tonnes C	yr ⁻¹ ;	
	 A = Act 	ivity data, representing the annual area of cul	tivated organic soils, in hectares (´	
	• <i>EF</i> =Tie	er 1, default IPCC emission factors, express	ed in units of tonnes C ha ⁻¹ (2).	
	(1)The da	ata are obtained through the stratification o	f two different global datasets:	
		rmonized World Soil Database (FAO <i>et al.</i> , by Histosols classes.	2012), used to estimate the area	
		obal Land Cover dataset, GLC2000 (EU-J of cropland and grassland area in each pixe		
	For cropl (2008):	For cropland, three "cropland" classes from GLC2000 are used, following Yo (2008):		
	CLASS	NAME	CROPLAND SHARE PER PIXEL	
	16	Cultivated and managed areas	100%	
	17	Mosaic: cropland/tree cover/Other natural vegetation	50%	
	18	Mosaic: cropland/Shrub and/or grass cover	10%	
	sub-dom of the GL (2) The E assigned Vol. 4, Cl Research	eriod 1990-present and for 2030 and 2050, ain are a constant value, representing the y C2000 database. F values are specified in IPCC, 2006, Vol. at the pixel level to the relevant climate in 3, Annex 3A.5. The climatic zones map us Centre of the European Commission (EC scriptions.	year 2000, i.e. the reference year 4, Ch. 5, Table 5.6. The EF were zone, as defined in IPCC, 2006 sed was developed by the Joint	
	The glob	ysis was carried out in a GIS environment, or al emissions dataset was summarized by co ensionless conversion factors used were		
	 44/12, 1 10⁻³, to 	to convert the emissions from tonnes C to to convert the emissions from tonnes C to Gg	onnes CO ₂ gas; _I C	
	downloa	pland sub-domain contains the following d: country-level GHG emissions in both Gg (factors; and activity data.		
	factors a	nties in the estimates of GHG emissions are and activity data. In the case of cultivate ion is available in the IPCC Guidelines (IPCC	d organic soils, more detailed	

5.2.3.5).

Calculation procedure

Step 1. Activity data


Activity data are the annual area of organic soils covered by cropland, expressed in hectares. The $Area_{(T)}$ is calculated in a GIS environment.

The cropland area was identified from the Global Land Cover data set GLC2000 (EU-JRC, 2003), available at http://bioval.jrc.ec.europa.eu/products/glc2000/glc2000.php. The three "cropland" classes were used. The mosaic classes were reduced proportionally to the general share of the cropland per pixel for each class, according to the values reported in the metadata above.

The area of organic soils was identified from the Harmonized World Soil Database, available at http://www.fao. org/nr/lman/abst/lman_080701_en.htm, selecting all the pixels where histosols are present, with the relative pixel share expressed as percentages.

The $Area_{(T)}$ has been calculated by overlaying the soil and the land cover layers. The values are fixed, calculated per each country and accessible in FAOSTAT Emissions database on Emissions-Land Use/Cropland domain from 1990 to 122012 as follows:

FIGURE 45 Cropland items from the FAOSTAT Emissions database



Step 2. Emissions (C)

The emissions are calculated in a GIS environment at the pixel level, using the formula:

Emissions
$$(C)_{(T)} = Area_{(T)} \times EF_{(T)}$$

Where

- Emission $(C)_{(T)}$ = Annual emissions of C for item T, tonnes C yr⁻¹
- $Area_{(T)}$ = Activity data, representing the area of cropland organic soils under item T, ha
- $EF_{(T)} = \text{Tier 1}$, default IPCC emission factors for item T, tonnes C ha⁻¹ yr⁻¹ (Table 58A)
- T = The item of this category source is Cropland organic soils.

The EF values are specified in IPCC, 2006, Vol. 4, Ch. 5, Table 5.6. The EF were assigned at the pixel level to the relevant climate zone, as defined in IPCC, 2006, Vol. 4, Ch. 3, Annex 3A.5. The climatic zones map used was developed by the Joint Research Centre of the European Commission (EC-JRC, 2010), in accordance with IPCC prescriptions.

The emissions are related to the year 2000, which is the reference year of the land cover map used in the estimations. These values are also used for subsequent years, to cover the period 1990-2011. Additional country-level disaggregations that have occurred over the period 1990-2011 are also provided.

As long as a country's boundary does not change, the area values for years following 2011 can be produced by simply replicating the country's 2011 values.

Step 3. Emissions (CO₂)

Emissions
$$(CO_2)_{(T)} = Emissions (C)_{(T)} \times \frac{44}{12} \times 10^{-3}$$

Where

- Emission $(CO_2)_{(T)}$ = Annual emissions of CO_2 for item T, $Gg CO_2yr^{-1}$
- Emission $(C)_{(T)}$ = Annual emissions of C for item T, tonnes C yr⁻¹

Step 4. Emissions (CO₂eq)

Emissions
$$(CO_2eq)_{(T)} = Emissions (CO_2)_{(T)}$$

Where

- Emissions $(CO_2eq)_{(T)} = CO_2$ emissions from item T in CO_2 equivalent, $Gg CO_2eq yr^{-1}$
- Emissions $(CO_2)_{(T)} = CO_2$ emissions from item T, $Gg CO_2 yr^{-1}$

Step 5. Implied emissions factor for C

$$IEF_{(T)} = \frac{Emissions(C)_{(T)}}{Area_{(T)}}$$

Where

- $IEF_{(T)}$ = Implied emission factor for item T, tonnes C ha⁻¹
- Emissions $(C)_{(T)} = C$ emissions from item T, Gg C yr⁻¹
- A =Area of Cropland Organic Soils under item T ha

Step 6. Replication

Emissions are assumed to be constant in any given year, unless changes in the boundaries of the countries occur. In that case, a new GIS analysis would be required.

5.2.3 Grassland

Title Grassland Definition Greenhouse gas (GHG) emissions data from grassland are currently limited to emissions from grassland organic soils. They are associated with carbon losses from drained histosols under grassland.

Methodology and quality information

Methods and processing

GHG emissions data from grassland are currently limited to emissions from grassland organic soils. They are associated with carbon losses from drained organic soils. The FAOSTAT data are computed at Tier 1, following the criteria established in IPCC, 2006, Vol. 4, Ch. 6.

The emissions are estimated at pixel level, using the formula:

Emission = A * EF

where

- Emission = Annual emissions, in units of tonnes C yr¹;
- A = Activity data, representing the annual area of grassland organic soils, in hectares (1).
- EF=Tier 1, default IPCC emission factors, expressed in units of tonnes C ha⁻¹ (2).
- (1) The data are obtained through the stratification of two different global data sets:
- i. the Harmonized World Soil Database (FAO et al., 2012), used to estimate the area covered by histosols classes, and
- ii. the Global Land Cover dataset, GLC2000 (EU-JRC, 2003), used to estimate the amount of cropland and grassland area in each pixel.
- For grassland, two "herbaceous" and two mosaics GLC2000 classes are used, in line with the FAO Land Cover Classification Scheme (LCCS):

CLASS	NAME	GRASSLAND SHARE PER PIXEL
13	Herbaceous Cover, closed-open	100%
14	Sparse herbaceous or sparse shrub cover	50%
17	Mosaic: Cropland/Tree Cover/Other natural vegetation	25%
18	Mosaic: Cropland/Shrub and/or grass cover	45%

iii. the Gridded Livestock of the World for cattle and sheep (Wint and Robinson, 2007), used as an additional mask over grassland histosols as a proxy to estimate drained area. With this mask, only pixels with non-zero drained grassland histosols area and livestock density >1 head/ha are included.

For the period 1990-present as well for 2030 and 2050, the activity data reported in this sub-domain are a constant value, representing the year 2000, i.e., the reference year of the GLC2000 database.

(2) The EF values are those specified in IPCC, 2006, Vol. 4, Ch. 6, Table 6.3. The EF were assigned at pixel level to the relevant climate zone, as defined in IPCC, 2006, Vol. 4, Ch. 3, Annex 3A.5. The climatic zones map used was developed by the Joint Research Centre of the European Commission (EC-JRC, 2010), following IPCC prescriptions.

The analysis was carried out in a GIS environment, combining the above datasets; the global emissions dataset was summarized by country.

The dimensionless conversion factors used were:

- 44/12, to convert the emissions from tonnes C to tonnes CO₂ gas;
- 10⁻³, to convert the emissions from tonnes C to Gg C

The cultivated organic soils domain contains the following data categories, available for download: country-level GHG emissions in both Gg C, Gg CO₂ and Gg CO₂eq; implied emission factors; and activity data.

Uncertainties in the estimates of GHG emissions are due to uncertainties in emission factors and activity data. In the case of cultivated organic soils, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol. 4, Ch. 6, Section 6.2.3.5).

Calculation procedure

Step 1. Activity data


The activity data are the annual area of organic soils covered by grassland, in hectares. The $Area_{(T)}$ is calculated in a GIS environment.

The grassland area was identified from the Global Land Cover dataset, GLC2000 (EU-JRC, 2003), available at http://bioval.jrc.ec.europa.eu/products/glc2000/glc2000.php, using four classes. The mosaics classes were reduced proportionally to the general share of the grassland per pixel for each class, according to the values reported in the metadata above.

The area of organic soils was identified from the Harmonized World Soil Database, available at http://www.fao. org/nr/lman/abst/lman_080701_en.htm, selecting all the pixels where histosols were present, with the related pixel share expressed as a percentage.

The Area_(T) was calculated by overlaying the soil and the land cover layers, and using the Gridded Livestock of the World as a proxy to identify the areas of managed grasslands. The values are fixed and calculated for each country. The results are accessible in the FAOSTAT Emissions database, in the Emissions-Land Use/Grassland domain from 1990 to 2012:

FIGURE 46
Grassland items from the FAOSTAT Emissions database



Step 2. Emissions (C)

The emissions are calculated in a GIS environment at the pixel level, using the formula:

Emissions
$$(C)_{(T)} = Area_{(T)} \times EF_{(T)}$$

Where

- $Emission(C)_{(T)} = Annual emissions of C for item T, tonnes C yr⁻¹$
- $Area_{(T)}$ = Activity data, representing the area of grassland organic soils under item T, ha
- $EF_{(T)} = \text{Tier 1}$, default IPCC emission factors for item T, tonnes C ha⁻¹ yr⁻¹ (Table 59A)
- T =The item of this category source is Grassland organic soils.

The EF values are those specified in IPCC, 2006, Vol. 4, Ch. 5, Table 5.6. The EF values were assigned at the pixel level to the relevant climate zone, as defined in IPCC, 2006, Vol. 4, Ch. 3, Annex 3A.5. The climatic zones map used was developed by the Joint Research Centre of the European Commission (EC-JRC, 2010), following IPCC prescriptions.

The emissions concern the year 2000, which is the reference year of the land cover map used in the estimations. These values are also used for subsequent years, to cover the period 1990-2011. Additional country-level disaggregation occurring over the period 1990-2011 are also provided.

As long as a country's boundary does not change, the area values for years following 2011 can be produced by simply replicating the country's 2011 values.

Step 3. Emissions (CO₂)

Emissions
$$(CO_2)_{(T)} = Emissions (C)_{(T)} \times \frac{44}{12} \times 10^{-3}$$

Where

- Emission $(CO_2)_{(T)}$ = Annual emissions of CO_2 for item T, $Gg CO_2yr^{-1}$
- Emission $(C)_{(T)}$ = Annual emissions of C for item T, tonnes C yr⁻¹

Step 4. Emissions (CO₂eq)

Emissions
$$(CO_2eq)_{(T)} = Emissions (CO_2)_{(T)}$$

Where

- Emissions $(CO_2eq)_{(T)} = CO_2$ emissions from item T in CO_2 equivalent, $Gg CO_2eq yr^{-1}$
- Emissions $(CO_2)_{(T)} = CO_2$ emissions from item T, Gg CO_2 yr⁻¹

Step 5. Implied emissions factor for C

$$IEF_{(T)} = \frac{Emissions(C)_{(T)}}{Area_{(T)}}$$

Where

- $IEF_{(T)}$ = Implied emission factor for item T, tonnes $C ha^{-1}$
- Emissions $(C)_{(T)} = C$ emissions from item T, tonnes C yr⁻¹
- A =Area of Grassland organic soils under item T

Step 6. Replication

Emissions are assumed to be constant in any given year, unless changes in the boundaries of the countries occur. In that case, a new GIS analysis would be required.

5.2.4 Burning-Biomass

Dataset information			
Title	Burning-Biomass		
Definition	Greenhouse Gas (GHG) emissions from burning of biomass consist of methane and nitrous oxide gases, from biomass combustion of forest land, covering the classes "Humid and Tropical Forest" and "Other Forests". The gases consist of methane, nitrous oxide, and carbon dioxide gases, from combustion of organic soils.		
Methodology and qua	lity information		
Methods and processing	Greenhouse Gas (GHG) emissions from burning of biomass consist of gases produced from the burning of biomass for the following items: "Humid Tropical Forest," "Other Forests" and "Organic Soils."They consist of methane (CH ₄), nitrous oxide (N ₂ O), and, only in the case of Organic Soils, also carbon dioxide (CO ₂).		
	The "Humid Tropical Forest" item is defined by aggregating the following FAO-FRA Global Ecological Zones (FAO 2012): "Tropical Rainforest" and "Tropical moist deciduous forest".		
	The "Other Forests" area contains the rest of the FAO-FRA Global Ecological Zones for Forests.		
	The "Organic Soils" item is defined as the histosols class in the Harmonized World Soil Database (FAO et al., 2012), consistently with the FAOSTAT Emissions database "Cultivation Organic Soils" definition.		
	The emissions are estimated following the IPCC, 2006, Vol. 4, Ch. 2, Equation 2.27, by aggregation at the national level of spatially distributed information, produced on a spatial grid at 0.25 degree resolution (approximately 25 km at the Equator).		
	For each item, and for each pixel, the following formula is used:		
	Emission = A * EF		
	where		
	 Emission = GHG emissions in g CH₄, g N₂O and g CO₂. 		
	 A = Activity data, representing the total mass of fuel burned, in kg of dry matter (1); 		
	 EF=Tier 1 IPCC emission factor, expressed in g CH₄, g N₂O or g CO₂, per kg of burned dry matter (2). 		
	(1) The total mass of fuel burned is computed by multiplying the burned area (i) by the fuel biomass consumption values (ii).		
	(i)Yearly composite values of burned area are obtained as the sum of the monthly burned area data, from the Global Fire Emission Database v.4 (GFED4; Giglio <i>et al.</i> 2013). For forests, the GFED4 burned forest area is an aggregate of the burned areas in the following MODIS land cover classes (MCD12Q1, Hansen <i>et al.</i> , 2000): Evergreen Needle-leaf, Evergreen Broadleaf, Deciduous Needle-leaf, Deciduous Broadleaf, and Mixed Forest.		
	For "HumidTropical Forest", the burned area is obtained by overlapping the GFED4 Burned Forest area data with the relevant FAO-FRA Global Ecological Zones.		
	For "Other Forest", the burned area is obtained as the GFED4 burned Forest area minus the Humid Tropical Forest burned area.		
	For "Organic Soils", the burned area is obtained by overlapping the GFED4 Burned area data with the Histosols class information of the Harmonized World Soil Database (FAO <i>et al.</i> , 2012), assuming the even distribution of organic soils within the grid cell.		
	The projections of activity data for 2030 and 2050 are the average of the entire duration of the burned area satellite data time series (1996-present).		
	(ii) Fuel biomass consumption values are taken from IPCC, 2006, Vol. 4, Ch. 2, Table 2.4. The different values are climate-dependent, and were geographically allocated using the JRC Climate Zones map (EC-JRC, 2010).		
	(2) EF values are taken from IPCC, 2006, Vol. 4, Ch. 2, Table 2.5. The different values were assigned at the pixel level to the relevant climate zone. The climatic zones map used was developed by the Joint Research Centre of the European Commission (EC-JRC, 2010), following IPCC prescriptions.		

Methodology and quality information

Methods and processing

For each item, the emissions at the pixel level were subsequently aggregated at the country level, using the FAO Global Administrative Unit Layers (GAUL) data set.

The dimensionless conversion factors used were

- 10⁻⁹, to convert the emissions from g CH₄ to Gg CH₄ and g N₂O to Gg N₂O;
- GWP- CH₄ = 21 (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq;
- GWP.N₂O = 310 (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq (IPCC, 1996, Technical Summary, Table 4, p. 22).

The Burning-Biomass sub-domain contains the following data categories, available for download for each land cover class: country-level GHG emissions, provided as totals, in Gg CH₄, Gg N₂O and Gg CO₂eq; the implied emission factor; the burned area; and total mass of fuel available.

Since GFED4 data are not available before 1996 and for future periods, a single yearly value for the period 1990-1995, as well as for projections for 2030 and 2050, were estimated for each country, as the average of the period 1996-2012.

Uncertainties in the estimates of GHG emissions are due to uncertainties in the emission factors and activity data. These may be related to, *inter alia*, natural variability, partitioning fractions, lack of spatial or temporal coverage, spatial aggregation, or errors in satellite data. In the case of biomass burning, more detailed information is available in the IPCC Guidelines (IPCC, 2006, Vol. 4, Ch. 5, Section 5.2.4.4).

Calculation procedure

Step 1. Activity data

The items under the Burning-Biomass domain include the following land cover classes: *Humid tropical forest*, *Other forest*, and *Organic soils*.

Greenhouse gases considered in this domain are CH₄, N₂O, and (only for organic soils) CO₂.

The annual burned area for each land cover is derived from the Global Fire Emission Database burned area data set, Version 4 (GFED4).


GFED4 provides global monthly burned area maps at 0.25° spatial resolution, from mid-1995 to present (currently 2012, see Giglio *et al.* 2013). The GFED4 dataset combines a 500m Moderate Resolution Imaging Spectroradiometer (MODIS) burned area dataset with active fire datasets from the Tropical Rainfall Measuring Mission (TRMM) Visible and Infrared Scanner (VIRS) and the Along-Track Scanning Radiometer (ATSR) family of sensors. The annual burned area for each land cover class is calculated by adding up the GFED4 monthly data, by country.

The biomass burned (BB) activity of the Biomass burned (dry matter) for each land cover class, as well as the resulting emissions of N_2O , CH_4 , and (only for organic soils) CO_2 , are estimated using IPCC (2006) default methodology, based on Equation 2.27 therein; reference is made to default parameters for biomass consumption per hectare in different climates and vegetation types provided by Table 2.4, and to emission factors for each vegetation type taken from Table 2.5. The Biomass burned and emission computations are performed at the pixel level in a GIS environment, using additional data sets to associate the correct parameters to the specific climate and vegetation characterization of the pixel.

The methodology set out below enables the calculation of annual GHG emissions for the period 1990-1995, based on averages of the period 1996-present; of the conversion of non-CO₂ emissions into CO₂ equivalents; and of implied emission factors for N_2O , CH_4 , and (only for organic soils) CO_2 .

The data is available in the FAOSTAT Emissions database, in the Emissions-Land Use/Burning-Biomass domain from 1990 to 2050:

FIGURE 47



Step 2. Production of 1990-1995 data

For each country, yearly values for the period 1990-1995, X_i , may be estimated simply as the average value over the period 1996-present:

$$X_i = \overline{X_{1996-present}}$$

Where

- X = Variable (Burned area, Biomass burned, Emissions)
- $X_{(1996-present)}$ = Average value for the years 1999-present
- i = Year (period 1990-1995)

Step 3. Emissions (CO₂eq) from CH₄ and N₂O

$$Emissions(CO_2eq)_{(T)} from (gas) = Emissions(gas)_{(T)} \times GWP_{(gas)}$$

Where

- *Emissions* $(CO_2eq)_{(T)}$ from (gas) = Amount of GHG emissions in CO_2eq from biomass burning for land cover T, Gg CO_2eq yr⁻¹
- Emissions $(gas)_{(T)}$ = Amount of Gas emissions from biomass burning for crop T, Gg yr⁻¹
- $GWP_{(N2O)} = 310$ (100-year time horizon global warming potential), to convert Gg N₂O to Gg CO₂eq
- $GWP_{(CH4)} = 21$ (100-year time horizon global warming potential), to convert Gg CH₄ to Gg CO₂eq
- T = Land cover class
- *Gas*= CH₄, N₂O

Step 4. Emissions (CO₂eq)

$$Emissions(CO_2eq)_{(T)} = \\ Emissions(CO_2eq)_{(T)}from \ N_2O + Emissions(CO_2eq)_{(T)}from \ CH_4 + \\ Emissions(CO_2)_{(only \ for \ T=Organic \ Soils)}$$

Where

- Emissions $(CO_2eq)_{(T)}$ = Amount of emissions in CO_2eq from biomass burning for land cover T, $Gg CO_2eq yr^{-1}$
- Emissions $(CO_2eq)_{(T)}$ from N_2O = Amount of N_2O emissions in CO_2eq from biomass burning for land cover T, $Gg CO_2eq yr^{-1}$
- Emissions (CO₂eq)_(T) from CH_4 = Amount of CH₄ emissions in CO₂eq from biomass burning for land cover T, Gg CO₂eq yr⁻¹
- Emissions (CO₂)_(T=Organic Soils) = Amount of CO₂ emissions from biomass burning for organic soils, Gg CO₂eq yr⁻¹
- T = Land cover class

Step 5. Implied emission factor for N₂O, CH₄ and CO₂

$$IEF_{(T,gas)} = \frac{Emissions(gas)_{(T)}}{BB_{(T)}} \times 10^6$$

Where

- $IEF_{(T, gas)}$ = Implied emission factor for Gas for land cover T, g/kg burned
- Emissions $(gas)_{(T)}$ = Amount of Gas emissions from biomass burning for land cover T, Gg yr⁻¹
- $BB_{(T)}$ = Biomass burned for land cover T, tonnes yr⁻¹
- T = Land cover class
- $Gas = Gas (CH_4, N_2O, CO_2)$

Annex 1

GLOSSARY: CONCEPTS, DEFINITIONS AND SYMBOLS

Title	Definition	Source
Production	The domain of production includes the figures relating to total domestic production, within or outside the agricultural sector, i.e. including non-commercial production and production from kitchen gardens. Unless otherwise indicated, production is reported at the farm level for crop and livestock products (i.e. in the case of crops, excluding harvesting losses) and in terms of live weight for fish items (i.e. the actual ex-water weight, at the time of the catch). With regard to livestock, all data relates to total meat production, from both commercial and farm slaughter. Data are expressed in terms of dressed carcass weight, excluding offal and slaughter fats. Production of beef and buffalo meat includes veal; mutton and goat meat include meat from lambs and kids; and pig meat includes bacon and ham in fresh equivalent. Poultry meat includes meat from all domestic birds and refers, wherever possible, to ready-to-cook weight.	FAOSTAT Glossary Concepts common to FAOSTAT & CountrySTAT
Production Quantity of Primary Crops	Data refer to the actual harvested production from fields, orchards or gardens, excluding harvesting and threshing losses and those parts of crops that were not harvested for any reason. Production therefore includes the quantities of the commodity sold in markets (marketed production) and the quantities consumed or used by producers (auto-consumption). When the production data available refers to a production period falling in two successive calendar years, and it is not possible to allocate the amounts produced to the respective years, the production data is usually referred to the year in which the bulk of production took place. Crop production data are recorded in tonnes (t). In many countries, crop production data are obtained as a function of the estimated yield and the total area. If countries enforce such a compilation method of production statistics, they must ensure that the total area does not refer to sown or planted area, which would thus give the "biological production," but to the area that was actually harvested during the year. It is recommended that primarily production in terms of harvested production be reported; when this is not possible, the concept adopted in reporting production (and yield) figures should be clearly indicated.	
Area Harvested	The data refer to the area from which a crop is gathered. Therefore, the area harvested excludes the area from which, although sown or planted, there was no harvest due to damage, failure, etc. It is usually net for temporary crops and sometimes gross for permanent crops. The net area differs from the gross area insofar as the latter includes uncultivated patches, footpaths, ditches, headlands, shoulders, shelterbelts, etc. If the crop under consideration is harvested more than once during the year, as a consequence of successive cropping (i.e. the same crop is sown or planted more than once in the same field during the year), the area is counted as many times as it is harvested. On the contrary, the area harvested will be recorded only once if the crop is successively gathered during the year from the same standing crops. With regard to mixed and associated crops, the area sown relating to each crop should be reported separately. When the mixture refers to particular crops, generally grains, it is recommended to treat the mixture as if it were a single crop; therefore, the area sown would be recorded only for the crop reported. The area harvested is reported in hectares (ha).	
Area Sown	This refers to the area on which sowing or planting has been carried out, for the crop under consideration, on the soil prepared for that purpose. The area is usually reported net of uncultivated patches, footpaths, ditches, headlands, shoulders, shelterbelts, etc. For tree crops, the gross concept may be applied. With regard to mixed and associated crops, countries are requested to report the area sown for each crop separately. When the mixture refers to particular crops, generally grains, it is recommended to treat the mixture as if it were a single crop. Data are recorded in hectares (ha). The information on area sown allows for a particular application of the SUA system, in which the quantity allotted for the next year's sowing, that enters the accounts for the current year, is calculated as a seeding rate times the area sown of the next year.	

Title	Definition	Source
Crop yield	The harvested production per unit of harvested area, for crop products. In most cases, yield data are not recorded, but are obtained by dividing the production data by the data on area harvested. The data on the yields of permanent crops are not as reliable as those for temporary crops, either because most of the area information may correspond to planted area, as for grapes, or because the area figures reported by the countries are scarce or unreliable, as for example for cocoa and coffee. Data are expressed as Production (Hg)/Area harvested (ha).	
Number of Live Animals (Stocks)	This variable indicates the number of animals of the species present in the country at the time of enumeration. It includes animals raised either for draft purposes, for meat, eggs and dairy production, or kept for breeding. Live animals in captivity for fur or skin such as foxes, minks etc. are not included in the system, although fur skin trade is reported. The enumeration to be chosen when more than one survey is taken, is that closest to the beginning of the calendar year. Livestock data are reported in terms of the number of heads (units), except for poultry, rabbits and other rodents, which are reported in thousand units. For FAO, figures for the year N relate to animals enumerated by the country any day between October of year N-1 and September of year N. The statistics are related to the total stock and the number of female animals.	
Milk Animals	Milk animals are those which, in the course of the reference period, have been milked. This concept is related to that applied for the production of milk, excluding the milk sucked by young animals. If, for example, the whole milk of a cow is sucked by the calf, the cow is not considered to be a "milk animal". Data are reported in terms of the number of heads (units).	
Laying Animals	Covers the number (in thousands) of all domestic birds which have contributed to egg production during the year, wherever they lay, and the corresponding total production, including eggs intended to be used for hatching. However, the figure excludes waste on farms. Some countries have no statistics on egg production, and estimates must be derived from related data such as chicken or total poultry numbers, and reported or assumed rates of egg laying.	
Primary Crops	Primary crops are those which come directly from the land, and which have not undergone any real processing apart from cleaning. These crops maintain all the biological qualities possessed when they were still on the plants. Certain primary crops can be aggregated, with their actual weight, into totals offering meaningful figures on area, yield, production and utilization; for example, cereals, roots and tubers, nuts, vegetables and fruits. Other primary crops can be aggregated only in terms of one or the other component common to all of them. For example, primary crops of the oil-bearing group can be aggregated in terms of oil or oil cake equivalent. Primary crops are divided into temporary and permanent crops. Temporary crops are those which are both sown and harvested during the same agricultural year, sometimes more than once; permanent crops are sown or planted once and not are replanted after each annual harvest.	FAOSTAT commodity list
Cereals:	These are generally of the gramineous family. For FAO purposes, the term refers to crops harvested for dry grain only.	
Wheat	"Triticum spp.": common (T. aestivum), durum (T. durum) spelt (T. spelta)". Common and durum wheat are the main types. The main varieties of common wheat are spring and winter, hard and soft, and red and white. At the national level, different varieties should be reported separately, reflecting their different uses. Used mainly for human food.	
Rice Paddy	"Oryza spp." mainly oryza sativa. Rice grain after threshing and winnowing. Also known as rice in the husk and rough rice. Used mainly for human food.	
Barley	"Hordeum spp.: two-row barley (H. disticum), six-row barley (H. hexasticum), four-row barley (H. vulgare)." Tolerates poorer soils and lower temperatures better than does wheat. Varieties include with husk and without (naked). Used as a livestock feed, for malt and for preparing foods. The roasted grains are a coffee substitute.	
Maize	"Zea mays Corn, Indian corn, mealies". A grain with high germ content. At the national level, hybrid and ordinary maize should be reported separately, owing to their widely different yields and uses. Used largely for animal feed and commercial starch production.	
Rye	"Secale cereal". A grain that is tolerant of poor soils, high latitudes and altitudes. Mainly used in making bread, whisky and beer. When fed to livestock, it is generally mixed with other grains.	

Title	Definition	Source
Oats	"Avena spp., mainly Avena sativa". A plant with open, spreading panicle-bearing large spikelets. Used primarily in breakfast foods. Makes excellent fodder for horses.	FAOSTAT commodity list
Millets	These include, inter alia: "barnyard or Japanese millet (Echinocloa frumentacea); ragi, finger or African millet (Eleusine coracana); teff (Eragrostis abyssinica); common, golden or proso millet (Panicum miliaceum); koda or ditch millet (Paspalum scrobiculatum); pearl or cattail millet (Pennisetum glaucum); foxtail millet (Setaria italic)". Small-grained cereals that include a large number of different botanical species. Originated by the domestication of wild African grasses in the Nile valley and the Sahel zone, millets were subsequently taken to China and India. These cereals tolerate arid conditions and possess a small, highly nutritious grain that stores well. Used locally, both as a food and as a livestock feed. In all areas where they are cultivated, millets are used in traditional beer brewing. Also used as a feed for birds.	
Sorghum	"Sorghum spp.": guinea corn (S. guineense); common, milo, feterita, kaffir corn (S. vulgare); durra, jowar, kaoliang (S. dura)." A cereal that has both food and feed uses. Sorghum is a major food grain in most of Africa, where it is also used in traditional beer brewing. It is desirable hybrid and other varieties to be reported separately.	
Roots and Tubers	Plants yielding starchy roots, tubers, rhizomes, corms and stems. Used mainly for human food (as such or in processed form), for animal feed and for manufacturing starch, alcohol and fermented beverages including beer. The denomination "roots and tubers" excludes crops cultivated mainly for feed (mangolds, swedes) or for processing into sugar (sugar beets), and those classified as "roots, bulb and tuberous vegetables" (onions, garlic and beets). The term does include starch and the starchy pith and flour obtained from the trunk of the sago palm and the stem of the Abyssinian banana (Musa ensete). Certain root crops, notably bitter cassava, contain toxic substances, particularly in the skins. As a result, certain processes must be undertaken to make the product safe for human consumption. Apart from their high water content (70-80 percent), these crops contain mainly carbohydrates (largely starches that account for 16-24 percent of their total weight), with very little protein and fat (0-2 percent each). Methods of propagating root crops vary. A live potato tuber or seed must be planted, but only part of the live yam tuber and a piece of the stalk (not the root) in the case of cassava. Production data of root crops should be reported in terms of clean weight, i.e. free of earth and mud. FAO distinguishes between seven primary root and tuber crops.	
Potatoes	"Solanum tuberosum, Irish potato". A seasonal crop grown in temperate zones all over the world, but primarily in the northern hemisphere.	
Oil Crops:	Include both annual (usually called oilseeds) and perennial plants whose seeds, fruits or mesocarp and nuts are valued, mainly for the edible or industrial oils that are extracted from them. Dessert and table nuts, although rich in oil, are listed under Nuts. Annual oilseed plants that are either harvested green or are used for grazing and for green manure are included with Fodder Crops. Some of the crops included in this chapter are also fiber crops, in that both the seeds and the fibers are harvested from the same plant. Such crops include: coconuts, yielding coir from the mesocarp; kapok fruit; seed cotton; linseed; and hempseed.	
Soybeans	"Glycine soja". The most important oil crop. Also widely consumed as a bean and in the form of various derived products because of its high protein content, e.g. soya milk, meat substitutes, etc.	
Livestock – Live Animals	The term "livestock" is used in a broad sense to cover all grown animals, regardless of age, location or purpose of breeding. Non-domesticated animals are excluded under this definition unless they are kept or raised in captivity. Domestic animals included are large and small quadrupeds, poultry, insects (bees) and larvae of insects (silkworms). Figures on livestock numbers should refer to live animals enumerated on a given day or on several consecutive days. The FAO practice is that figures for an indicated year relate to animals reported by countries for any day between October of the previous year and September of the year indicated. Statistics on live animals by age, sex and utilization are generally not included in the resulting list, even though such breakdowns are extremely desirable in terms of national statistics. For each animal species, FAO proposes that information be maintained on changes in national herds during the year, according to the following equation: initial herd + animals born + imports of live animals - exports of live animals - natural losses - slaughter = closing herd.	

Title	Definition	Source
Cattle	"Cattle Common ox (<i>Bos taurus</i>)"; zebu, humped ox (<i>Bos indicus</i>); Asiatic ox (subgenus Bibos); Tibetan yak (<i>Poephagus grunniens</i>)." Animals of the genus listed, regardless of age, sex, or purpose raised. Data are expressed in number of heads.	FAOSTAT commodity list
Buffaloes	"Buffalo Indian, Asiatic, pygmy, water buffalo (<i>Bubalus bubalus</i> ; <i>B. arnee</i> ; <i>B. depressicornis</i>); African buffalo (<i>genus Syncerus</i>); American bison (<i>Bison bison</i>); European bison (<i>Bison bonasus</i>); beeffalo (a cross between a bison and a domesticated beef animal)". Animals of the genus listed, regardless of age, sex, or purpose raised. Wild bisons and buffaloes are excluded. Data are expressed in terms of the number of heads.	
Sheep	"Ovis spp.:"Animals of the genus listed, regardless of age, sex, or purpose raised. It includes Uriel, Argali, Bighorn, Karakul and Astrakhan. Data are expressed in terms of number of heads.	
Goats	"Goat, Capra spp". Animals of the genus listed, regardless of age, sex, or purpose. Hircus, Ibex, Nubiana, Pyrenaica, Tibetana, Kashmir and Angora are included. Data are expressed in terms of number of heads.	
Pigs	"Pigs: domestic pig (<i>Sus domestica</i>); wild boar (<i>Sus scrofa</i>)". Animals of the genus listed, regardless of age, sex, or purpose raised. Non-domesticated wild boars are excluded. Data are expressed in terms of number of heads.	
Chickens	"Chicken fowl (<i>Gallus domesticus</i>); Guinea fowl (<i>Numida meleagris</i> " Data are expressed in terms of 1000 heads.	
Ducks	"Ducks, Anas spp." Data are expressed in terms of 1000 heads.	
Turkeys	"Geese, Anser spp." Data are expressed in terms of 1000 heads.	
Horses	"Horses, <i>Equus caballus spp</i> ." Animals of the genus listed, regardless of age, sex, or purpose raised. Data are expressed in terms of number of heads.	
Asses	"Asses, Equus asinus spp." Animals of the genus listed, regardless of age, sex, or purpose raised. Data are expressed in terms of number of heads.	
Mules	Includes hinnies. Mules are the offspring of a male ass and a female horse (mare); a hinny is the offspring of a female ass and a male horse (stallion). Both are sterile. Animals of the genus listed, regardless of age, sex, or purpose raised. Data are expressed in terms of number of heads.	
Other Camelids	Various species of Lama: e.g. <i>glama pacos</i> (alpaca); <i>peruana</i> (llama); <i>huanacos</i> (guanaco); <i>vicugna</i> (vicuna)". Animals of the genus listed, regardless of age, sex, or purpose raised. Data are expressed in terms of number of heads.	
Camels	"Bactrian camel (<i>Camelus bactrianus</i>); Arabian camel (<i>C. dromedarius</i>)". Animals of the genus listed, regardless of age, sex, or purpose raised. Data are expressed in terms of number of heads.	
Dairy cows and Buffalo	Dairy cows are defined in this method as mature cows that are producing milk in commercial quantities for human consumption. This definition corresponds to the dairy cow population reported in the FAO ProductionYearbook. In some countries, the dairy cow population consists of two well-defined segments: (i) high-producing (also called improved) breeds in commercial operations; and (ii) low-producing cows, managed with traditional methods. These two segments can be combined, or can be evaluated separately by defining two dairy cow categories. However, the dairy cow category does not include cows kept principally to produce calves for meat or to provide draft power. Low productivity multi-purpose cows should be considered as "Other cattle."	IPCC 2006- Vol.4- CH10 page 10.10
Broiler chickens	Typically grown approximately 60 days before slaughter. Data are obtained by multiplying data from FAOSTAT (available in the Production – Live Animals domain of the FAOSTAT database) by 1000.	IPCC 2006- Vol.4- Ch. 10 page 10.8
Market swine	Considered as a "dynamic" animal, and may vary from nursery pigs weighing less than 30 kilograms to finished pigs weighing over 90 kilograms. Data are obtained multiplying the total heads of pigs (available in the Production – Live Animals domain of the FAOSTAT database) by 0.9.	IPCC 2006- Vol.4- Ch. 10, Table 10.19
Breeding swine	Considered as a "static" animal. Data are obtained by multiplying the total heads of pigs (available in the Production – Live Animals domain of the FAOSTAT database) by 0.1.	IPCC 2006- Vol.4- Ch. 10, Table 10.19

Title	Definition	Source
Annual average population	Annual average populations are estimated in various ways, depending on the data available and on the nature of the animal population. In the case of static animal populations (e.g. dairy cows, breeding swine, layers), estimating the annual average population may be as simple as obtaining data related to one-time animal inventory data. However, estimating annual average populations for a growing population (e.g. meat animals, such as broilers, turkeys, beef cattle, and market swine) requires further evaluation. Most animals in these growing populations are alive for only part of a complete year. Animals should be included in the populations regardless of whether they were slaughtered for human consumption or they die of natural causes.	IPCC 2006- Vol.4- Ch. 10 page 10.8
Fertilizers	The domain includes data on Fertilizers. They are compiled in terms of fertilizer products and then converted into nutrients and classified according to the HS coding systems. Finally, total N (Nitrogen), P_2O_5 (Phosphate) and K_2O (Potash) are calculated for: Production, Imports, Exports, Non-fertilizer use and Consumption. Fertilizers may be organic, inorganic or mineral. Organic fertilizers play an important role in crop production and are derived from animals, plants and compost. Mineral fertilizers are available to farmers in solid or liquid form, and are delivered to the farm either in bulk, in bags or in pressurized containers. All fertilizers contain at least one of the major plant nutrients: nitrogen (N), phosphorus (P) or potassium (K).	FAOSTAT Glossary Concepts common to FAOSTAT & CountrySTAT
Consumption of Fertilizers	Refers the total quantity of fertilizers applied to the soil to increase crop yields (for crop production use). This refers to the "Actual Consumption, defined as actual deliveries to agriculture by the manufacturers or actual quantity consumed by a country for agriculture production".	
Synthetic Fertilizers	Inorganic compounds usually derived from by-products of the petroleum industry. Examples are ammonium nitrate, ammonium phosphate, superphosphate, and potassium sulphate.	
Ammonia, anhydrous (NH ₃)	A material mostly produced by a synthetic process; at standard temperature and pressure, it is a gas. Fertilizer-grade anhydrous ammonia contains about 82% of nitrogen (HS code 2814).	
Ammonium nitrate (NH ₄ NO ₃)	Produced by neutralizing nitric acid (HNO ₃) with ammonia (NH ₃). Ammonium nitrate may be in white or off-white granular or prilled form, and coated with a suitable material to prevent absorption of moisture and caking during storage. Pure ammonium nitrate may have a total nitrogen content of about 35%, of which one-half is present as ammoniac nitrogen and the other half as nitrate nitrogen (HS code 310230).	
Ammonium sulphate (NH ₄) ₂ SO ₄	Produced by reacting ammonia with sulphuric acid (H_2SO_4). It is produced in the form of fine white granules or crystals, and contains no less than 20.6% nitrogen in ammoniac form (HS code 310221).	
Calcium ammonium nitrate (NH ₄ NO ₃ +CaCO ₃)	Produced from ammonium nitrate and finely pulverized calcium carbonate (CACO ₃). It contains no less than 20.5% and up to 28% of nitrogen, half of which is in the form of ammoniac nitrogen and the other half in the form of nitrate nitrogen. It is produced in the form of white, off-white or grey granules or prills (HS code 310240).	
Urea – CO(NH ₂) ₂	Produced from synthetic ammonia and carbon dioxide (CO_2) and contains 46% nitrogen in ammoniac form. Urea may be in granular, prilled or crystalline form (HS code 310210).	
Urea and ammonium nitrate solutions	Produced from concentrated solutions of urea and ammonium nitrate, by chemical or blending processes (HS code 310280).	
Diammonium phosphate (DAP) – (NH ₄) ₂ HPO ₄	Produced by evaporating a solution of phosphoric acid with an excess of ammonia (HS code 310530).	
Monoammonium phosphate (MAP) – NH ₄ H ₂ PO ₄	Formed when a solution of phosphoric acid is added to ammonia until the solution is distinctly acid (HS code 310540).	
Other NP compounds	Can be produced as the result of a chemical reaction of nitric acid on phosphate rock, with or without added ammonia and/or phosphoric and/or sulphuric acid; a reaction between sulphuric acid and ammonia; or by simple mechanical mixing or blending. Other NP compounds may also include some AN grades with small amounts of phosphates (HS code 310500).	

Title	Definition	Source
Other nitrogen & phosphates compounds	Can be produced as the result of a chemical reaction of nitric acid on phosphate rock, with or without added ammonia and/or phosphoric and/or sulphuric acid; a reaction between sulphuric acid and ammonia; or by simple mechanical mixing or blending. Other NP compounds may also include some AN grades with small amounts of phosphates (HS code 310551).	FAOSTAT Glossary Concepts common to FAOSTAT &
Other nitrogen & phosphorus compounds	Can be produced as the result of a chemical reaction of nitric acid on phosphate rock, with or without added ammonia and/or phosphoric and/or sulphuric acid; a reaction between sulphuric acid and ammonia; or by simple mechanical mixing or blending. Other NP compounds may also include some AN grades with small amounts of phosphates (HS code 310559).	CountrySTAT
NPK complex	Can be produced as the result of a chemical reaction of nitric acid on phosphate rock – the nitro phosphate route, with or without added ammonia and/or phosphoric and/or sulphuric acid – or between sulphuric acid and ammonia, i.e. the ammoniation route (HS code 310610).	
NPK complex <=10kg	Can be produced as the result of a chemical reaction of nitric acid on phosphate rock – the nitro phosphate route, with or without added ammonia and/or phosphoric and/or sulphuric acid – or between sulphuric acid and ammonia, i.e. the ammoniation route (HS code 310510).	
NPK complex >10kg	Can be produced as the result of a chemical reaction of nitric acid on phosphate rock – the nitro phosphate route, with or without added ammonia and/or phosphoric and/or sulphuric acid – or between sulphuric acid and ammonia, i.e. the ammoniation route (HS code 310520).	
NPK blends	Consist of two or more intermediate granular fertilizer materials of matching physical characteristics, blended together (HS code 310620).	
Potassium nitrate (KNONH ₃ H ₂ PO ₄)	Can be produced from naturally occurring sodium nitrate and potassium chloride, and typically contains 13%N and 45% $\rm K_2O$ (HS code 283421).	
Organic Fertilizers	Animal or vegetable fertilizers, whether or not mixed together or chemically treated; or fertilizers produced by the mixing or chemical treatment of animal or vegetable product. These play an important role in crop production and are derived from animals, plants and compost. Mineral fertilizers are available to farmers in solid or liquid form, and are delivered to farms either in bulk, in bags or in pressurized containers. They include Cattle manure, Horse manure, Pig manure, Sheep manure, Poultry manure, Guano, Composts and Sewage waste.	
Forest	Land spanning more than 0.5 hectares with trees higher than 5 metres and a canopy cover of more than 10 percent, or trees able to reach these thresholds in situ. It does not include land that is predominantly under agricultural or urban land use.	FRA 2010 Terms and Definitions
Primary Forest	Naturally regenerated forest of native species, where there are no clearly visible indications of human activities and the ecological processes are not significantly disturbed.	
Other Naturally Regenerated Forest	Naturally regenerated forest with clearly visible indications of human activities and where the trees are predominantly of introduced species.	
Planted Forest	Forest predominantly composed of trees established through planting and/or deliberate seeding, and where the planted/seeded trees are predominantly of introduced species (Rubber plantations, mangroves and bamboo).	
Carbon density of living biomass	Carbon in all living biomass above the soil, including stem, stump, branches, bark, seeds, and foliage.	
Forest land	Includes all land with woody vegetation consistent with the thresholds used to define forest land in the national GHG inventory, sub-divided at the national level into managed and unmanaged land and also by ecosystem type, as specified in the IPCC Guidelines. This category also includes systems with vegetation that currently falls below, but is expected to exceed, the threshold of the forest land category.	IPCC 2006- Glossary
Net Forest Conversion	Change from forest land use to another use. For example, deforestation, reforestation and afforestation activities.	
Dry Matter	Dry matter refers to biomass that has been dried to an oven-dry state, often at $70^{\circ}\text{C}.$	

Title	Definition	Source
Area equipped for power irrigation	Area equipped for irrigation where pumps are used for water supply, from the source to the scheme. The term also includes areas where water is drained out with human- or animal-driven water lifting devices. The term does NOT refer to pumping required for the technology used within the field (such as sprinkler irrigation or localized irrigation, which require pressure and thus pumping).	AQUASTAT database of FAO
Area equipped for full control irrigation: sprinkler irrigation	A sprinkler irrigation system consists of a pipe network through which water moves under pressure, before being delivered to the crop via sprinkler nozzles. The system basically simulates rainfall in that water is applied through overhead spraying. These systems are also known as overhead irrigation systems. Unit measure: 1000 hectares.	
Area equipped for full control irrigation: localized irrigation	Localized irrigation is a system where the water is distributed under low pressure through a piped network, in a pre-determined pattern, and applies water as a light discharge to each plant or adjacent to it. There are three main categories of localized irrigation: drip irrigation (where drip emitters are used to apply water slowly to the soil surface), spray or micro-sprinkler irrigation (where water is sprayed to the soil near individual plants or trees) and bubbler irrigation (where a light stream is applied to flood small basins or the soil adjacent to individual trees). The following terms are also sometimes used to refer to localized irrigation: micro-irrigation, trickle irrigation, daily flow irrigation, drop-irrigation, sip irrigation, and diurnal irrigation. Unit measure: 1000 hectares.	
Motor gasoline	Gasoline (or motor gasoline) is a light hydrocarbon oil for use in internal combustion engines such as motor vehicles, excluding aircraft. It distills between 35°C and 200°C, and is treated to reach a sufficiently high octane number, generally between 80 and 100 RON. The substance may be treated by reforming, blending with an aromatic fraction, or by adding to it benzole or other additives (such as tetraethyl lead).	UNSD Glossary (Energy balances and electricity profiles)
Residual fuel oil	A heavy oil that makes up the distillation residue. It comprises all fuels (including those obtained by blending) with a kinematic viscosity above 27.5 cSt at 38oC. Its flash point is always above 50oC and its specific gravity is higher than 0.90. It is commonly used by ships and industrial large-scale heating installations as a fuel in furnaces or boilers.	
Coal	Coal corresponds to 'Other bituminous coal' or, when not available for the specific country, to 'Hard coal' (which includes the latter). Hard coal is a coal that has a high degree of coalification, with a gross calorific value above 23,865 KJ/kg (5,700 kcal/kg) on an ash-free but moist basis, and a mean random reflectance of vitrinite of at least 0.6. Slurries, middling and other low-grade coal products, which cannot be classified according to the type of coal from which they are obtained, are included under hard coal. There are two sub-categories of hard coal: (i) coking coal and (ii) other bituminous coal and anthracite (also known as steam coal). Coking coal is a hard coal with a quality that allows the production of coke suitable to support a blast furnace charge. Steam coal is coal used for steam raising and space heating purposes and includes all anthracite coals and bituminous coals not classified as coking coal. 'Other bituminous coal' has a higher volatile matter and a lower carbon content than anthracite.	
Anthracite	Coal that has a high degree of coalification with a gross calorific value above 23,865 KJ/kg (5,700 kcal/kg) on an ash-free but moist basis, and a mean random reflectance of vitrinite of at least 0.6. Slurries, middling and other low-grade coal products, which cannot be classified according to the type of coal from which they are obtained, are included under hard coal. There are two sub-categories of hard coal: (i) coking coal and (ii) other bituminous coal and anthracite (also known as steam coal). Coking coal is a hard coal with a quality that allows the production of coke suitable to support a blast furnace charge. Steam coal is coal used for steam raising and space heating purposes, and includes all anthracite coals and bituminous coals not classified as coking coal.	
Electricity	Final electricity consumption by the agriculture sector.	

Title	Definition	Source
Gas-diesel oils used in fisheries	Gas-diesel oils include heavy gas oils used in fishery. Gas oils are obtained from the lowest fraction from atmospheric distillation of crude oil, while heavy gas oils are obtained by vacuum re-distillation of the residue from the atmospheric distillation. Gas-diesel oil distils between 180°C and 380°C. Several grades are available depending on the uses: diesel oil for diesel compression ignition (cars, trucks, marine, etc.), light heating oil for industrial and commercial uses, and other gas oil including heavy gas oils which distil between 380°C and 540°C and which are used as petrochemical feed stocks.	Adapted from IEA Oil products definitions
Residual fuel oil used in fisheries	Oils that make up the distillation residue. This comprises all residual fuel oils, including those obtained by blending, used in fisheries. Its kinematic viscosity is above 10 cSt at 80°C. The flash point is always above 50°C and the density is always higher than 0.90 kg/l.	
Gas-diesel oils	Heavy oils distilling between 200oC and 380oC, but distilling less than 65% in volume at 250oC, including losses, and 85% or more at 350oC. Its flash point is always above 50oC and its specific gravity is higher than 0.82. Heavy oils obtained by blending are grouped together with gas oils on the condition that their kinematic viscosity does not exceed 27.5 cSt at 38oC. Also included are middle distillates intended for the petrochemical industry. Gas-diesel oils are used as a fuel for internal combustion in diesel engines, as a burner fuel in heating installations, such as furnaces, and for enriching water gas to increase its luminosity. Other names for this product are diesel fuel, diesel oil, gas oil and solar oil.	UNSD Glossary (Energy balances and electricity profiles)
Natural gas (including LNG)	Gases consisting mainly of methane occurring naturally in underground deposits. It includes both non-associated gas (originating from fields producing only hydrocarbons in gaseous form) and associated gas (originating from fields producing both liquid and gaseous hydrocarbons), as well as methane recovered from coal mines and sewage gas. Production of natural gas refers to dry marketable production, measured after purification and extraction of natural gas liquids and sulphur. Extraction losses and the amounts that have been re-injected, flared, and vented are excluded from the data on production.	Natural gas (including LNG)
Liquefied petroleum gas (LPG)	Hydrocarbons which are gaseous under conditions of normal temperature and pressure but are liquefied by compression or cooling to facilitate storage, handling and transportation. They are (i) extracted by stripping of natural gas at crude petroleum and natural gas sources; (ii) extracted by stripping of imported natural gas in installations of the importing country; and (iii) produced both in and outside of refineries, in the course of processing crude petroleum or its derivatives. It comprises propane (C_3H_8) , butane (C_4H_{10}) , or a combination of the two. Also included is ethane (C_2H_6) from petroleum refineries or natural gas producers' separation and stabilization plants.	UNSD Glossary (Energy balances and electricity profiles)
Country emission factor (electricity-only)	The CO ₂ emissions in the numerator of this indicator include emissions from fossil fuels, industrial waste and non-renewable municipal waste that are consumed for electricity generation; electricity output in the denominator includes electricity generated from fossil fuels, nuclear, hydropower (excluding pumped storage), geothermal, solar, biofuels, etc. As a result, the emissions per kWh may vary from year to year, depending on the generation mix. A fixed-heat-efficiency approach is used, which fixes the efficiency of the heat part of the generation, and calculates the electricity part of the input accordingly. More information is available in Chapter 4 of the IEA's CO ₂ Emissions from Fuel Combustion Highlights, available at https://www.iea.org/co2highlights/co2highlights.pdf.	IEA, 2012, CO ₂ Emissions from Fuel Combustion – Highlights

Symbols

С	Carbon
C	
CO ₂	Carbon dioxide
N	Nitrogen
N ₂ O	Nitrous oxide
CH ₄	Methane
CaCO ₃	Calcic limestone
CaMg(CO ₃) ₂	Dolomite
CO(NH ₂) ₂	Urea
NH ₄ +	Ammonium
OH-	Hydroxyl ion
HCO ₃ -	Bicarbonate
CO ₂ eq	Carbon dioxide equivalent
NOX	Nitrogen oxides
N ₂ O-N	Units of measurement expressed in molecular nitrogen (N_2). - Molar mass $N_2O = 14+14+16 = 44$ g/mol - Molar mass $N_2 = 14*2 = 28$ g/mol 1 kg of N_2O -N corresponds to 1 kg * 44/28 = 1.57 kg N_2O
Gg	A unit of mass equal to 1,000,000,000 grams
Tonnes	A unit of mass equal to 1,000 kilograms
PJ	A petajoule is equal to 1 joule x 10 ¹⁵
TJ	A terajoule is equal to 1 joule x10 ¹²
KWh	Kilowatt hour
GWP (N ₂ O)	Global Warming Potential for N ₂ O, kg-CO ₂ -e (kg-N ₂ O)-1 (IPCC default = 310)

Annex 2

DATA ACTIVITIES: EXAMPLE OF NATIONAL QUESTIONNAIRE ON DATA COLLECTION

Countries are requested to provide agricultural data coming from primary sources, i.e. agricultural censuses, surveys and administrative records, including related metadata (the methodology, the contact name responsible of the office, *etc.*).

The data are organized in order to respond to a standard methodology of concepts, definitions and classification of commodities adopted by FAO (FAOSTAT) in line with the international requirements of data quality, so that the data is comparable and reliable. Figure 47 below shows a sample Excel file of the production domain, which includes statistics on the area harvested by crop and on livestock production.

FIGURE 48 Example of a FAOSTAT Questionnaire

	FOOD AND AGRICULTURE ORG. STATISTICS DIVISION	ANIZATION OF THE UNITED NATI	ONS
	Questionnaire on Cro	pp and Livestock Production and Excel Version	Utilization
			XX - Country Reference Years: 2010-2012
-	National	Reporting Office and Contact name	
Reporter	name.	CHORESON MELO AN JOWNSON CONTINUE CHORESON - JOH DC.	
Title:			
STEEL STREET STREET	ration and Office:		
Address			
Web site			
Signature	K:		
Tel:	Fax	E-mail:	Date:
	- Description of commodities as well as instr	ock Production) lities overage. i included in this questionnaire. Please revise outlions are given in supporting sheets. op and livestock production we would highly a	55 550 85 85 U.S.
	or via the FAO Representative Office in your country of	erably by e-mail, to the following address: Proc	fuction-Data@fao.org; rme di Caracalla, 00163 Rome, Italy.

			Se	ection 1: PRII	MARY CROP P	RODUCTION			
extends ectually of the wh elate to commod	into the si harvested role nut in fermented tries for m	annual production of cor ubsequent year, products and the corresponding to the shell. Coconuts data I and dried beans. For Bu- sere details. Production of esponds to several FAO	on should be allocated to stal harvested production a refer to the weight of the anana's and Plantains, the lata for groups of items:	to the calender year in Cereals and Pulse no whole nut, exclud- he weight of the cent	in which the bulk of the is are reported in dry gr ing the fibrous outer hur ral stalk of the buncher	harvest takes place rains. Roots and Tub sk. Seed Cotton pro- s should not be inclu-	Area and production ers are reported in fres suction should include ded in the production of	of crops should inc h weight. Nuts data both fibre and seed lata. Please refer to	lude all areas relate to the weigh Cocoa Beans data the description of
	Cross		2010	0	2011	1	201	2	Notes:
Code	ref to CPC Ver.2	COMMODITY	Area Horrested (Ha)	Production (MT)	Area Harvested (Ha)	Production (MT)	Area Harvested (Ha)	Production (MT)	If a different unit is enclusively used to that specified, pleas indicate the unit use- as well as any other selevant notes
		Coreals		3.270.364		3.636.000			
15	0111	Wheat (including meslin)	12.000	20,000	13,000	23,000	14,000	20,000	
27	0113	Rice, paddy	87,000	218,111	90,000	233,000	92,000	212,000	
56	0112	Maize	1,032,000	2,373,501	1,063,000	2,551,000	1,094,000	2,734,000	
79	0118	Millet	167,000	267,973	172,000	292,000	175,000	244,000	
83	0114	Sorghum Other cereals, please specify	355,000	390,779	364,000	437,000	373,000	336,000	
		Roots and Tubers							
125	01592	Cassava	794,000	3,017,118	822,000	2,712,000	851,000	2,807,000	
116	01510	Potatoes	36,000	167,153	37,000	180,000	39,000	185,000	
122	01591	Sweet potatoes Other mots and tubers, please specify	442,000	1,987,203	450,000	1,798,000	452,000	1,852,000	

The FAOSTAT questionnaires are also sent to compile Fertilizers, Pesticides, Land Irrigation, Producer Prices and other agricultural domains; for trade, countries provide data coming from the national customs.

This information is provided by National Statistical Offices or Ministries. Detailed information on questionnaires is available in the "Questionnaires" section of the FAO Statistics Division's website (http://www.fao.org/economic/ess/ess-home/questionnaires/en/).

The FAOSTAT commodity list of Production is based on the Standard International Trade Classification of the UN (SITC), to provide a framework for the harmonization of production and trade data and the compilation of SUA/FBS, and to calculate various indexes (Agricultural Production Index Number – food and non-food). Details on the relevant concepts and definitions are available in the Glossary section.

Although FAOSTAT, as an international source, has adopted a process of data harmonization and validation required by international standards, during the data collection process (i.e. questionnaire compilation), experience with countries has shown that national statistical structures have had to face the following issues, at different levels:

- Data scattering coming from multiple structures responsible for producing statistics;
- Production of the same kind of statistics by different structures;
- Incompleteness of statistics;
- Absence or incompleteness of national classifications;
- Discrepancies between national and international classifications of products;
- Lack of correspondence between national and international nomenclatures;
- Absence of an organized national level for the validation and harmonization of data;
- · Weakness of data organization;
- Lack of management tools and digital archiving of statistical data;
- Weakness of the technical documentation that must accompany the production data (metadata);

With the objective to produce good quality food and agriculture statistics, the FAO Statistics Division implements projects to promote data quality, which in turn enables effective decision and policy making. In this context, national technical working groups have been organized, composed of the main national institutions. In collaboration with the Statistics Division, it is envisaged that these groups will address and eventually solve the following issues encountered in practice:

- 1. Incoherent data among national sources;
- 2. Missing data;
- 3. Anomalies in historical trends;
- 4. Incoherence between related indicators;
- 5. Incoherence between CORE and Sub-national data;
- 6. Inconsistencies between local and international concepts and definitions;
- 7. Lack of correspondence between national and international classifications.

In collaboration with Member Countries, the FAO Statistics Division has provided suggestions to address each of the abovementioned issues. These may be viewed as part of the E-Learning course, available at http://www.fao.org/economic/ess/ess-capacity/countrystathome/foundations/en/#.Uuul0T1dWHc

Annex 3

PARAMETERS FOR THE DIFFERENT EMISSION CATEGORIES

TABLE 1A Emission factor for methane (kg CH₄ head⁻¹), by animal category and IPCC area

IPCC Area	Dairy	Non-dairy	Buffalo	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	58.00	27.00	55.00	18.00	10.00	10.00	46.00	46.00
Eastern Europe	99.00	58.00	55.00	18.00	10.00	10.00	46.00	46.00
Africa	46.00	31.00	55.00	18.00	10.00	10.00	46.00	46.00
Oceania	90.00	60.00	55.00	18.00	10.00	10.00	46.00	46.00
Western Europe	117.00	57.00	55.00	18.00	10.00	10.00	46.00	46.00
Latin America	72.00	56.00	55.00	18.00	10.00	10.00	46.00	46.00
Middle east	46.00	31.00	55.00	18.00	10.00	10.00	46.00	46.00
Northern America	128.00	53.00	55.00	18.00	10.00	10.00	46.00	46.00
Asia	68.00	47.00	55.00	18.00	10.00	10.00	46.00	46.00

Developed/developing	Pigs	Sheep	Goats
Developed	1.50	8.00	8.00
Developing	1.00	5.00	5.00

TABLE 2A Emission factor for methane for each animal category, kg $\mathrm{CH_4}$ head $^{-1}$, by country

FAO Country Name	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Horses ,	Asses N	Mules C	Camels L	Llamas	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys
Afghanistan	2.00	2.00	4.00	3.00	3.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Albania	13.00	7.00	2.00	3.00	2.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Algeria	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
American Samoa	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Andorra	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Angola	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Anguilla	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Antigua and Barbuda	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Argentina	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Armenia	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Aruba	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Australia	29.00	2.00	2.00	13.00	24.00	0.28	0.20	2.34	1.10	1.10	2.37	2.37	0.02	0.03	0.03	0.02
Austria	21.00	0.00	4.00	00.9	9.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Azerbaijan	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Bahamas	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Bahrain	3.00	1.00	5.00	5.00	5.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Bangladesh	2.00	2.00	2.00	00.9	00'9	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Barbados	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Belarus	11.00	0.00	2.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Belgium	21.00	0.00	4.00	00.9	9.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Belgium-Luxembourg	21.00	0.00	4.00	00.9	9.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Belize	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Benin	1.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Bermuda	93.00	2.00		19.00	37.00	0.28	0.20	2.34	1.10	1.10	2.37	2.37	0.02	0.03	0.03	0.02
Bhutan	2.00	2.00	4.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Bolivia (Plurinational State of)	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Bosnia and Herzegovina	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Botswana	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Brazil	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
British Virgin Islands	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Brunei Darussalam	31.00	1.00	2.00	7.00	7.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Bulgaria	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Burkina Faso	1.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Burundi	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Cambodia	31.00	1.00	2.00	7.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02

open 100 <th>FAO Country Name</th> <th>Cattle, dairy</th> <th>Cattle, non-dairy</th> <th>Buffalo</th> <th>Swine, market</th> <th>Swine, breeding</th> <th>Sheep</th> <th>Goats</th> <th>Horses</th> <th>Asses</th> <th>Mules (</th> <th>Camels 1</th> <th>Llamas</th> <th>Chickens, Broilers</th> <th>Chickens, Layers</th> <th>Ducks</th> <th>Turkeys</th>	FAO Country Name	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Horses	Asses	Mules (Camels 1	Llamas	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys
tep 48.00 1.00 1.00 18.00 0.18 0.18 1.56 stream 1.00 <th< th=""><th>noc</th><th>1.00</th><th>1.00</th><th>5.00</th><th>1.00</th><th>1.00</th><th>0.15</th><th>0.17</th><th>1.64</th><th>06.0</th><th>06.0</th><th>1.92</th><th>1.92</th><th>0.02</th><th>0.02</th><th>0.02</th><th>0.02</th></th<>	noc	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
syle 100 0.10 100 0.10		48.00	1.00		10.00	19.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
African Republic 1.00 1.00 1.00 1.00 1.00 1.01 1.04 1.00 1.00 1.01 1.00 1.01 1.00 1.00 0.15 0.17 1.04 African Republic 1.00 1.00 1.00 1.00 0.00 0.13 0.13 1.04 Al Slands 2.100 1.00 1.00 1.00 1.00 0.00 0.13 0.13 1.16 Hong Korng SAR 3.00 1.00 1.00 1.00 0.10 0.10 0.11 1.00 Hong Korng SAR 3.00 1.00 1.00 1.00 2.00 0.10 0.11 0.10 0.10 0.11 0.10 0.11 0.10 0.11 0.10 0.10 0.11 0.10 0.10 0.11 0.11 0.10 0.10 0.11 0.11 0.10 0.10 0.11 0.11 0.10 0.10 0.11 0.11 0.11 0.11 0.11 0.10 0.11 0.11 0.11<	erde	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
African Republic 1.00 0.10 0.10 0.11 1.09 1.00 1.00 0.10 0.10 0.10 0.10 0.10 0.10 0.11 1.00 1.00 0.10 0.10 0.11 1.00 1.00 0.10 0.10 0.11 1.00 1.00 0.10 0.11 1.00 0.10 0.11 1.00 1.00 0.10 0.11 1.00 1.00 1.00 0.11 1.00 1.00 0.11 1.00 1.00 0.11 1.00 1.00 0.10 0.11 1.00 1.00 0.11 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00	n Islands	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
1.00 1.00 1.00 1.00 1.00 1.00 0.20 0.22 2.19	African Republic	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
1,00 1,00		1.00	1.00	5.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Hong Kong SAR 1,00	el Islands	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	97.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Hong Kong SAR 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 Maeao SAR 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 mainland 9.00 1.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 pia 1.00 1.00 1.00 1.00 1.00 1.00 0.10 0.11 1.09 pia 1.00 1.00 1.00 1.00 1.00 0.10 0.11 1.09 pia 1.00 1.00 1.00 1.00 0.15 0.17 1.64 tea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 horie 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 kea 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 kea 1.00 1.00		1.00	1.00	1.00	1.00	1.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Macao SAR 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 ainial and province of signal and plants 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 sia 1.00 1.00 1.00 1.00 1.00 0.10 0.11 1.09 se 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 se 1.00 1.00 1.00 1.00 0.15 0.17 1.64 lica 1.00 1.00 1.00 1.00 0.15 0.17 1.64 voire 1.00 1.00 1.00 1.00 0.15 0.17 1.64 voire 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 voire 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 voire 1.00 1.00 1.00 1.00 <th>Hong Kong SAR</th> <th>9.00</th> <th>1.00</th> <th>1.00</th> <th>2.00</th> <th>2.00</th> <th>0.10</th> <th>0.11</th> <th>1.09</th> <th>09.0</th> <th>09.0</th> <th>1.28</th> <th>1.28</th> <th>0.01</th> <th>0.01</th> <th>0.01</th> <th>0.01</th>	Hong Kong SAR	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
mainland 9.00 1.00 1.00 2.00 2.00 0.11 1.09 sia 9.00 1.00 1.00 1.00 2.00 0.10 0.11 1.09 ss 1.00 1.00 1.00 1.00 1.00 0.10 0.11 1.09 ss 1.00 1.00 1.00 1.00 1.00 0.10 0.11 1.09 ss 1.00 1.00 1.00 1.00 0.10 0.11 1.04 tea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 Noire 1.00 1.00 1.00 1.00 0.15 0.17 1.64 Hoolie 1.00 1.00 1.00 1.00 0.10 0.11 1.04 Noire 1.00 1.00 1.00 1.00 0.10 0.11 1.04 Isoa 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 <t< th=""><th>Macao SAR</th><th>9.00</th><th>1.00</th><th>1.00</th><th>2.00</th><th>2.00</th><th>0.10</th><th>0.11</th><th>1.09</th><th>09.0</th><th>09.0</th><th>1.28</th><th>1.28</th><th>0.01</th><th>0.01</th><th>0.01</th><th>0.01</th></t<>	Macao SAR	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
laivan Province of side 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 side 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 se 1.00 1.00 1.00 1.00 0.10 0.15 0.17 1.64 lands 1.00 1.00 1.00 1.00 0.10 0.15 0.17 1.64 kica 1.00 1.00 1.00 1.00 0.10 0.17 1.64 kica 1.00 1.00 1.00 1.00 0.10 0.17 1.64 kica 1.00 1.00 1.00 1.00 0.10 0.17 1.64 storatic People's Republic 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 storatic People's Republic of the 1.00 1.00 1.00 1.00 1.00 0.10 0.11 0.11 1.64 storatic People's Republic of the	mainland	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
pia 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 lands 1.00 1.00 1.00 1.00 1.00 0.16 0.17 1.64 kica 1.00 1.00 1.00 1.00 0.16 0.17 1.64 koire 1.00 1.00 1.00 1.00 0.16 0.17 1.64 koire 1.00 1.00 1.00 1.00 0.16 0.17 1.64 koire 1.00 1.00 1.00 1.00 0.19 0.17 1.64 static People's Republic 1.00 1.00 1.00 1.00 0.19 0.19 0.13 1.56 story 1.00 1.00 1.00 1.00 0.19 0.19 0.19 1.54 story 1.00 1.00 1.00 0.19 0.19 <t< th=""><th>Taiwan Province of</th><th>9.00</th><th>1.00</th><th>1.00</th><th>2.00</th><th>2.00</th><th>0.10</th><th>0.11</th><th>1.09</th><th>09.0</th><th>09.0</th><th>1.28</th><th>1.28</th><th>0.01</th><th>0.01</th><th>0.01</th><th>0.01</th></t<>	Taiwan Province of	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64	bia	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
lands 1.00 <t< th=""><th>80</th><th>1.00</th><th>1.00</th><th>5.00</th><th>1.00</th><th>1.00</th><th>0.15</th><th>0.17</th><th>1.64</th><th>06.0</th><th>06.0</th><th>1.92</th><th>1.92</th><th>0.02</th><th>0.02</th><th>0.02</th><th>0.02</th></t<>	80	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
lands 30.00 2.00 13.00 24.00 0.15 0.15 0.17 1.64 Noire 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.05 0.01 0.15 0.17 1.64 Republic 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 slovakia 1.100 6.00 5.00 3.00 4.00 0.15 0.17 1.64 static People's Republic 1.100 6.00 5.00 3.00 4.00 0.15 0.17 1.64 static People's Republic 1.100 6.00 5.00 3.00 4.00 0.15 0.17 1.64 static People's Republic of the 1.00 1.00 1.00 2.00 2.00 0.10 0.11 1.64 rk 1.00 1.00<		1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
tica 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.15 0.17 1.64 Noire 1.00 1.00 1.00 1.00 1.00 1.00 0.02 0.22 2.19 Noire 1.00 1.00 1.00 1.00 1.00 0.01 0.01 0.10 0.15 0.17 1.64 Hepublic 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 stowakia 11.00 6.00 5.00 3.00 4.00 0.15 0.17 1.64 ratic People's Republic of the 1.00 6.00 5.00 3.00 4.00 0.15 0.17 1.64 ratic Republic of the 1.00 1.00 1.00 1.00 1.00 0.10 0.11 0.11 1.04 ratic Republic of the 1.00 1.00 1.00 1.00 0.10 0.10 0.11 1.04 1.04 ratic Republic of the 1	slands	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Noire 1.00 1.00 5.00 1.00 1.00 0.02 2.19 Holist 12.00 6.00 5.00 3.00 5.00 0.19 0.13 1.56 Holist 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 Holist 1.00 6.00 5.00 3.00 0.15 0.17 1.64 Indicates 1.00 6.00 5.00 3.00 0.19 0.13 1.56 Indicates 1.00 6.00 5.00 3.00 0.19 0.13 1.56 Indicates 1.00 1.00 1.00 1.00 0.10 0.11 1.00 Indicates 1.00	Sica	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
12.00 6.00 5.00 5.00 0.19 0.13 1.56 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 2.00 3.00 3.00 0.15 0.17 1.64 1.00 1.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 1.00 1.00 1.00 2.00 3.00 4.00 0.19 0.13 1.56 1.00 1.00 1.00 2.00 2.00 0.10 0.13 1.56 1.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 0.10 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1.00 1.	lvoire	1.00	1.00	5.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
tepublic 1.00		12.00	00.9	2.00	3.00	2.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
tepublic 16.00 1.00 2.00 3.00 3.00 0.15 0.17 1.64 slovakia 11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 artic People's Republic 9.00 1.00 5.00 3.00 4.00 0.19 0.13 1.56 ratic People's Republic 9.00 1.00 1.00 2.00 2.00 2.00 0.19 0.11 1.09 ratic People's Republic of the 1.00 1.00 1.00 2.00 2.00 0.10 0.11 1.00 ratic Republic of the 1.00		1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
sepublic 11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 slovakia 11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 ratic People's Republic 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 ratic Republic of the 1.00 1.00 1.00 2.00 0.10 0.11 1.09 rk 21.00 6.00 4.00 6.00 9.00 0.15 0.17 1.64 rk 1.00 1.00 1.00 1.00 0.15 0.17 1.64 can Republic 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rad Guinea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rad Guinea 1.00 1.00 1.00 0.15 0.17 1.64 rad Can 1.00 1.00 1.00 1.00 0.15 0.17		16.00	1.00	2.00	3.00	3.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
slovakia 11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 ratic People's Republic 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 ratic Republic of the 1.00 1.00 5.00 1.00 1.00 0.15 0.17 1.64 rk 21.00 6.00 4.00 6.00 9.00 0.15 0.17 1.64 rk 1.00 1.00 1.00 2.00 2.00 0.19 0.13 1.56 can Republic 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rad Guinea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rad Guinea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 ca 1.00 1.00 1.00 1.00 0.15 0.17 1.64 can Republic 1.00 1.00 1.00 1.00 </th <th>Republic</th> <th>11.00</th> <th>00.9</th> <th>2.00</th> <th>3.00</th> <th>4.00</th> <th>0.19</th> <th>0.13</th> <th>1.56</th> <th>0.76</th> <th>92.0</th> <th>1.58</th> <th>1.58</th> <th>0.02</th> <th>0.03</th> <th>0.02</th> <th>0.02</th>	Republic	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
ratic People's Republic 9.00 1.00 1.00 2.00 2.00 0.10 0.11 1.09 ratic Republic of the ratic Republic of 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rk 21.00 6.00 4.00 6.00 9.00 0.19 0.13 1.64 can Republic 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rador 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rial Guinea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 can Republic 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rad Guinea 1.00 1.00 1.00 0.15 0.17 1.64 ca 1.00 1.00 1.00 0.15 0.17 1.64 can I 1.00	slovakia	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
ratic Republic of the 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rk 21.00 6.00 4.00 6.00 9.00 0.19 0.13 1.56 ca 1.00 1.00 1.00 2.00 2.00 0.20 0.22 2.19 can Republic 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 r 1.00 1.00 1.00 1.00 0.15 0.17 1.64 ador 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rial Guinea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 ca 1.00 1.00 1.00 1.00 0.15 0.17 1.64 ca 1.00 1.00 1.00 1.00 0.15 0.17 1.64 ca 1.00 1.00 1.00 1.00 0.15 0.17 1.64	ratic People's Republic	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
rk 21.00 6.00 4.00 6.00 9.00 0.19 0.13 1.56 ca 1.00 1.00 1.00 2.00 2.00 0.20 0.22 2.19 can Republic 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 r 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 ador 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 rial Guinea 1.00 1.00 1.00 1.00 0.15 0.17 1.64 ca 1.00 1.00 1.00 0.15 0.17 1.6	ratic Republic of the	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
1.00	rk	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	97.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
lic 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.01 1.04 1.04 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.01 1.04 1.04 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.04	i	1.00	1.00	5.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
lic 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.01 1.04 1.04 1.00 1.00 1.01 1.04 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.01 1.04 1.04 1.00 1.01 1.04 1.00 1.01 1.04 1.04 1.00 1.01 1.04	ica	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
1.00 1.00 <th< th=""><th>ican Republic</th><th>1.00</th><th>1.00</th><th>1.00</th><th>1.00</th><th>1.00</th><th>0.15</th><th>0.17</th><th>1.64</th><th>06.0</th><th>06.0</th><th>1.92</th><th>1.92</th><th>0.02</th><th>0.02</th><th>0.02</th><th>0.02</th></th<>	ican Republic	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
1.00	ır	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.01 1.04 1.04 1.04 1.04 1.04 1.00 <th< th=""><th></th><th>1.00</th><th>1.00</th><th>5.00</th><th>1.00</th><th>1.00</th><th>0.15</th><th>0.17</th><th>1.64</th><th>06.0</th><th>06.0</th><th>1.92</th><th>1.92</th><th>0.02</th><th>0.02</th><th>0.02</th><th>0.02</th></th<>		1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
1.00 1.00 5.00 1.00 1.00 0.15 0.17 1.64 1.00 1.00 1.00 1.00 1.00 0.20 0.22 2.19 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 1ic 1.00 1.00 1.00 1.00 0.15 0.17 1.64	ador	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
a 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.20 0.22 2.19 an Republic 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64	rial Guinea	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
a 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 an Republic 1.00 1.00 1.00 1.00 0.15 0.17 1.64		1.00	1.00	2.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
an Republic 1.00 1.00 1.00 1.00 0.15 0.17 1.64	ica	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
	ican Republic	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
1.00 1.00 1.00 1.00 0.15 0.17 1.64	ır	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02

FAO Country Name	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Horses	Asses I	Mules (Camels L	Llamas	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys
Egypt	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
El Salvador	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Equatorial Guinea	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Eritrea	1.00	1.00	2.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Estonia	11.00	00.9	5.00	3.00	4.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Ethiopia	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Ethiopia PDR	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Falkland Islands (Malvinas)	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Faroe Islands	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Fiji	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Finland	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
France	23.00	7.00	4.00	00.9	10.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
French Guiana	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
French Polynesia	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Gabon	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Gambia	1.00	1.00	5.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Georgia	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Germany	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Ghana	1.00	1.00	5.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Gibraltar	27.00	8.00	5.00	7.00	11.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Greece	29.00	8.00	5.00	8.00	12.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Greenland	48.00	1.00		10.00	19.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Grenada	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Guadeloupe	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Guam	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Guatemala	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Guinea	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Guinea-Bissau	1.00	1.00	5.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Guyana	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Haiti	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Holy See	34.00	10.00	00.9	9.00	13.00	0.28	0.20	2.34	1.10	1.10	2.37	2.37	0.02	0.03	0.03	0.02
Honduras	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Hungary	11.00	00.9	5.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Iceland	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
India	2.00	2.00	5.00	5.00	2.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Indonesia	28.00	1.00	2.00	6.00	00.9	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Iran (Islamic Republic of)	15.00	1.00	2.00	3.00	3.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02

FAO Country Name	Cattle,	Cattle,	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Horses	Asses	Mules (Camels 1	Llamas	Chickens, Broilers	Chickens, Lavers	Ducks	Turkeys
Iraq	2.00	1.00	2.00	00	4.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Ireland	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	97.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Isle of Man	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Israel	2.00	1.00	2.00	3.00	3.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Italy	25.00	7.00	5.00	7.00	10.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Jamaica	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Japan	10.00	1.00	1.00	2.00	2.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Jordan	2.00	1.00	5.00	3.00	3.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Kazakhstan	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Kenya	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Kiribati	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Kuwait	2.00	1.00	2.00	2.00	2.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Kyrgyzstan	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Lao People's Democratic Republic	24.00	1.00	2.00	2.00	5.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Latvia	11.00	00.9	5.00	3.00	4.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Lebanon	2.00	1.00	4.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Lesotho	1.00	0.00	4.00	0.00	0.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Liberia	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Libya	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Liechtenstein	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Lithuania	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Luxembourg	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Madagascar	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Malawi	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Malaysia	28.00	1.00	2.00	00.9	00.9	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Maldives	17.00	1.00	2.00	4.00	4.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Mali	1.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Malta	43.00	13.00	8.00	11.00	16.00	0.28	0.20	2.34	1.10	1.10	2.37	2.37	0.02	0.03	0.03	0.02
Marshall Islands	28.00	2.00	2.00	13.00	23.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Martinique	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Mauritania	1.00	1.00	5.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Mauritius	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Mayotte	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Mexico	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Micronesia (Federated States of)	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Monaco	27.00	8.00	2.00	7.00	11.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
Mongolia	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01

FAO Country Name	Cattle, dairv	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Horses	Asses I	Mules (Camels 1	Llamas	Chickens, Broilers	Chickens, Lavers	Ducks	Turkeys
Montenegro	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02		0.02	0.02
Montserrat	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Morocco	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Mozambique	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Myanmar	23.00	1.00	2.00	2.00	2.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Namibia	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Nauru	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Nepal	5.00	2.00	4.00	3.00	3.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Netherlands	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Netherlands Antilles	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
New Caledonia	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
New Zealand	23.00	1.00	1.00	11.00	20.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Nicaragua	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Niger	1.00	1.00	5.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Nigeria	1.00	1.00	5.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Niue	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Norfolk Island	31.00	2.00	2.00	13.00	24.00	0.37	0.26	3.13	1.52	1.52	3.17	3.17	0.02	0.03	0.03	0.02
Northern Mariana Islands	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Norway	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Occupied Palestinian Territory	2.00	1.00	2.00	3.00	3.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Oman	3.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Pacific Islands Trust Territory	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Pakistan	2.00	2.00	2.00	4.00	4.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Palau	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Panama	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Papua New Guinea	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Paraguay	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Peru	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Philippines	28.00	1.00	2.00	00.9	0.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Pitcairn Islands	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Poland	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Portugal	34.00	10.00	00.9	9.00	13.00	0.28	0.20	2.34	1.10	1.10	2.37	2.37	0.02	0.03	0.03	0.02
Puerto Rico	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Oatar	3.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Republic of Korea	10.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Republic of Moldova	11.00	00.9	5.00	3.00	4.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Réunion	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Romania	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02

11.00 6.00 5.00 3 1.00 1.00 5.00 1 1.00 1.00 5.00 1 2.00 1.00 5.00 1 2.00 1.00 2.00 2 31.00 2.00 2.00 1.00 1.00 1.00 5.00 1 2.00 1.00 5.00 1 2.00 1.00 5.00 1 1.00 1.00 5.00 3 1.00 1.00 5.00 3 1.00 1.00 5.00 3 1.00 1.00 5.00 3 1.00 1.00 5.00 1 1.00 1.00 5.00 1 1.00 1.00 5.00 1 1.00 1.00 5.00 1 1.00 1.00 5.00 1 2.00 1.00 5.00 1 2.00 1.00 5.00 1 2.00 1.00 5.00 6.0	FAO Country Name	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Horses	Asses Mules		Camels L	Llamas	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys
act 1.00 1.00 5.00 1.00	Russian Federation	11.00	00:9	5.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
	Rwanda	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
tires and Nevise 200 1000 2.00 2.00 2.00 0.20 2.01 2.9 120 120 2.66 2.66 0.02 0.00 0.00 0.00 0.00 0.	Saint Helena	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
ucia 1 2 1 2 2 1 2	Saint Kitts and Nevis	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
	Saint Lucia	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
	Saint Pierre and Miquelon	48.00	1.00		10.00	19.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
with color 200 200 2400 0.20 0.22 219 120 120 0.00 <t< th=""><th>Saint Vincent and the Grenadines</th><th>1.00</th><th>1.00</th><th>1.00</th><th>1.00</th><th>1.00</th><th>0.15</th><th>0.17</th><th>1.64</th><th>0.90</th><th>0.90</th><th>1.92</th><th>1.92</th><th>0.02</th><th>0.02</th><th>0.02</th><th>0.02</th></t<>	Saint Vincent and the Grenadines	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
virthology 2700 8100 5.00 1100 0.19 0.13 1.56 0.76 1.56 1.56 1.56 1.56 1.56 1.56 1.56 1.56 1.56 1.50 1.00	Samoa	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
trabilation 100 100 5.00 100 100 100 100 100 100 100 100 100 100 100 100 100 101 100 <t< th=""><th>San Marino</th><th>27.00</th><th>8.00</th><th>5.00</th><th>7.00</th><th>11.00</th><th>0.19</th><th>0.13</th><th>1.56</th><th>92.0</th><th>0.76</th><th>1.58</th><th>1.58</th><th>0.02</th><th>0.03</th><th>0.02</th><th>0.02</th></t<>	San Marino	27.00	8.00	5.00	7.00	11.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
typical 200 100 500 500 0.01 6.00 0.01 6.00 0.01 6.00 0.01 6.00 0.02 0.03	Sao Tome and Principe	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
100 100 5.00 2.00 0.20 0.22 1.50 1.50 1.50 1.50 0.20 0.22 0.23 1.50 0.25 0.	Saudi Arabia	2.00	1.00	2.00	2.00	2.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
and Montanegro 11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 0.76 1.58 1.58 1.58 0.75 0.79 1.58 1.58 1.58 0.02 0.03 close 1.00 1.00 5.00 1.00 1.00 1.00 0.02 0.22 2.19 1.20 1.58 1.58 1.58 0.02 0.02 close 1.00 1.00 5.00 1.00	Senegal	1.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
and Montenegro 1100 6.00 5.00 4.00 0.13 1.56 0.76 1.58 1.58 1.68 0.02 0.03 alles 1.00 1.00 1.00 1.00 0.10	Serbia	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
	Serbia and Montenegro	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Loone 1,00 1,00 5,00 1,00 <t< th=""><th>Seychelles</th><th>1.00</th><th>1.00</th><th>5.00</th><th>1.00</th><th>1.00</th><th>0.15</th><th>0.17</th><th>1.64</th><th>06.0</th><th>06.0</th><th>1.92</th><th>1.92</th><th>0.02</th><th>0.02</th><th>0.02</th><th>0.02</th></t<>	Seychelles	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
ope 31,00 1,00 2,00 700 0.20 0.22 2.19 1,20 1,20 2,00 200 0.20 0.22 2.19 1,20 1,20 2,00 2,00 0.02 0.13 1,56 0.76 1,58 1,58 0.50 0.02 0.03 nish 1,100 6,00 5,00 3,00 4,00 0.13 0.13 1,56 0.76 1,58 1,58 0.02 0.02 0.03 nish 1,100 6,00 5,00 13.00 24,00 0.13 0.13 1,56 0.76 1,58 1,58 0.02 0.03 0.03 Africa 1,100 1,00 1,00 1,00 1,00 0.10 0.12 1,19 1,20 1,26 2,56 2,50 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.0	Sierra Leone	1.00	1.00	5.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
tigh 1100 6.00 5.00 4.00 0.13 1.56 0.76 1.56 1.56 1.56 1.56 0.76 1.56 1.56 1.56 1.56 1.56 0.76 1.56 1.56 0.76 1.56 1.56 1.56 0.76 1.56 0.76 1.56 0.76 1.56 1.56 0.76 1.56 <th< th=""><th>Singapore</th><th>31.00</th><th>1.00</th><th>2.00</th><th>7.00</th><th>7.00</th><th>0.20</th><th>0.22</th><th>2.19</th><th>1.20</th><th></th><th>2.56</th><th>2.56</th><th>0.02</th><th>0.02</th><th>0.02</th><th>0.02</th></th<>	Singapore	31.00	1.00	2.00	7.00	7.00	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
tial 1100 6.00 5.00 3.00 4.00 0.13 1.56 0.76 1.58 1.58 0.02 0.03 ia on Islands 3100 2.00 2.00 13.00 24.00 0.20 0.22 2.19 1.20 1.56 0.56 0.02 0.02 0.19 1.20 2.56 0.56 0.02 0.02 0.17 1.64 0.30 1.50 1.00 0.00 0.00 0.00 1.00 0.00 0.00 0.00 1.00 0.00	Slovakia	11.00	00'9	2.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
on Islands 3100 2.00 13.00 24.00 0.20 0.22 2.19 1.20 1.26 2.56 2.56 0.02 0.02 sight 1.00 1.00 1.00 1.00 0.20 0.22 2.19 1.20 1.56 2.56 0.26 0.02 0.02 Africa 1.00 1.00 1.00 0.10 0.20 0.22 2.19 1.20 1.26 2.56 0.02 0.02 Sudan 6.00 2.00 6.00 6.00 6.00 0.20 0.22 2.19 1.20 1.26 1.86 2.56 0.02 0.02 Sudan 6.00 2.00 6.00 6.00 6.00 0.20 0.22 2.19 1.20 1.26 1.86 1.86 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 0.02 <t< th=""><th>Slovenia</th><th>11.00</th><th>00.9</th><th>2.00</th><th>3.00</th><th>4.00</th><th>0.19</th><th>0.13</th><th>1.56</th><th>92.0</th><th>92.0</th><th>1.58</th><th>1.58</th><th>0.02</th><th>0.03</th><th>0.02</th><th>0.02</th></t<>	Slovenia	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	92.0	1.58	1.58	0.02	0.03	0.02	0.02
tigh 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.12 2.19 1.20 1.20 2.56 2.56 2.56 0.02 0.02 Sudan 6.00 2.00 6	Solomon Islands	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
Africa 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 0.90 0.90 1.92 1.92 1.92 1.92 0.02 0.02 Sudan 6.00 2.00 6.00 6.00 6.00 6.00 6.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.02 0.22 2.19 1.20 1.56 0.76 1.58 1.58 0.02 0.02 0.00 Ida 6.00 2.00 6.00 6.00 6.00 6.00 0.20 0.22 2.19 1.20 1.58 1.58 0.02 0.02 Ida 8.00 1.00 1.00 0.00 0.20 0.22 2.19 1.20 1.50 1.00 0.00 0.20 0.22 2.19 1.20 1.20 2.56 2.56 0.02 0.00 <th>Somalia</th> <th>1.00</th> <th>1.00</th> <th>2.00</th> <th>1.00</th> <th>1.00</th> <th>0.20</th> <th>0.22</th> <th>2.19</th> <th>1.20</th> <th>1.20</th> <th>2.56</th> <th>2.56</th> <th>0.02</th> <th>0.02</th> <th>0.02</th> <th>0.02</th>	Somalia	1.00	1.00	2.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Sudan 6:00 2:00 6:00 0:20 0:21 1:20 1:20 1:56 2:56 0:56 0:02 0:02 rida 0:00 5:00 5:00 1:00 0:10 0:13 1:56 0:76 1:58 1:58 0:02 0:03 rida 6:00 2:00 5:00 6:00 6:00 6:00 0:20 0:22 2:19 1:20 1:56 1:58 0:56 0:02 0:03 ridad 0:00 2:00 6:00 6:00 6:00 0:00 0:20 0:22 2:19 1:20 1:26 2:56 2:56 0:02 0:02 ridad 0:00 1:00 1:00 1:00 1:00 0:00 0:20 0:22 2:19 1:20 1:26 2:56 0:02 0:02 0:02 ridad 0:00 1:00 0:00 0:00 0:00 0:00 0:00 0:00 0:00 0:00 0:00 0:00 0:00	South Africa	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
tka 6:00 8:00 5:00 1100 0:19 0:13 1:56 0:76 0:76 1:58 1:58 0:50 0:00 tka 6:00 2:00 6:00 6:00 6:00 6:00 0:22 2:19 1:20 1:26 2:56 2:56 0:02 0:02 feat 1:00 2:00 5:00 6:00 6:00 6:00 0:20 0:22 2:19 1:20 1:26 2:56 0:02 0:02 feat 1:00 1:00 1:00 6:00 6:00 0:20 0:22 2:19 1:20 1:20 2:56 0:02 0:02 met 2:00 1:00 1:00 0:00 0:00 0:20 0:22 2:19 1:20 1:20 2:56 0:02 0:02 rd and Jan Mayen Islands 1:00 4:00 6:00 0:00 0:00 0:13 1:26 0:76 0:76 1:20 1:20 1:20 1:20 0:20 <	South Sudan	00.9	2.00	5.00	00.9	00.9	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
6.00 2.00 6.00 6.00 6.22 2.19 1.20 1.20 2.56 2.56 0.02 0.02 6.00 2.00 6.00 6.00 6.00 6.02 6.23 2.19 1.20 1.20 2.56 2.56 0.02 0.02 1.00 1.00 1.00 0.20 0.20 0.22 2.19 1.20 1.20 2.56 2.56 0.02 0.02 2.00 1.00 1.00 0.20 0.20 0.22 2.19 1.20 1.20 2.56 2.56 0.02 0.02 0.02 0.02 0.20 <th>Spain</th> <th>27.00</th> <th>8.00</th> <th>2.00</th> <th>7.00</th> <th>11.00</th> <th>0.19</th> <th>0.13</th> <th>1.56</th> <th>0.76</th> <th>0.76</th> <th>1.58</th> <th>1.58</th> <th>0.02</th> <th>0.03</th> <th>0.02</th> <th>0.02</th>	Spain	27.00	8.00	2.00	7.00	11.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
6.00 2.00 6.00 6.00 0.20 0.22 2.19 1.20 1.20 2.56 2.56 2.56 0.02 0.02 1.00 1.00 1.00 1.00 0.20 0.22 2.19 1.20 1.20 2.56 2.56 0.02 0.02 2.00 1.00 2.00 2.00 0.20 0.20 0.22 2.19 1.20 1.20 2.56 2.56 0.02 0.02 4.00 6.00 4.00 6.00 0.00 0.10 0.12 1.20 1.20 1.26 2.56 2.56 0.02 0.02 2.100 6.00 4.00 6.00 0.10 0.11 1.64 0.90 0.90 1.92 1.58 1.58 0.02 0.03 2.100 6.00 4.00 6.00 0.10 0.11 1.64 0.90 0.90 1.28 1.58 1.58 0.02 0.03 2.100 6.00 6.00 0.10	Sri Lanka	00.9	2.00	2.00	00.9	00.9	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
1.00 1.00 5.00 1.00 0.20 0.21 1.20 1.20 1.26 2.56 2.56 2.56 0.02 0.02 2.00 1.00 2.00 2.00 0.20 0.22 2.19 1.20 1.26 2.56 2.56 0.02 0.02 2.100 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 1.58 1.58 0.02 0.03 2.100 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 2.100 6.00 4.00 6.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 2.100 6.00 9.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 2.100 6.00 9.00 0.19 0.13 1.64 0.90 0.90	Sudan	00.9	2.00	2.00	00.9	00.9	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
45 2.00 1.00 2.00 2.00 0.22 2.19 1.20 1.20 2.56 2.56 2.56 0.02 0.02 41 2.100 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 1.58 1.58 0.02 0.03 0.03 2.100 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 1.58 1.58 0.02 0.03 2.100 6.00 4.00 6.00 9.00 0.13 1.56 0.76 1.58 1.58 0.02 0.03 2.100 6.00 4.00 6.00 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 2.100 6.00 4.00 6.01 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 2.00 1.00 2.00 0.13 0.14 0.16 0.16 0.19 0.19	Sudan (former)	1.00	1.00	2.00	1.00	1.00	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
45 21.00 6.00 4.00 6.01 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 1.00 1.00 1.00 1.00 1.00 0.15 0.17 1.64 0.90 0.90 1.92 1.92 0.02 0.02 21.00 6.00 4.00 6.00 9.00 0.13 1.56 0.76 1.58 1.58 0.02 0.03 21.00 6.00 4.00 6.00 9.00 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 21.00 6.00 4.00 6.00 0.13 0.15 0.76 0.76 1.58 1.58 0.02 0.03 21.00 1.00 5.00 0.10 0.11 1.64 0.90 0.90 1.92 0.02 0.03 0.03 21.00 1.00 1.00 0.10 0.11 1.00 0.00 1.90 0.11 1.00 0.90	Suriname	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20		2.56	2.56	0.02	0.02	0.02	0.02
1.00 1.00 5.00 1.00 1.00 0.15 0.17 1.64 0.90 0.90 1.92 1.92 0.02 0.02 0.02 21.00 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 21.00 6.00 4.00 6.00 0.019 0.13 1.56 0.76 1.58 1.58 0.02 0.03 2.00 1.00 5.00 3.00 0.19 0.13 1.64 0.30 0.90 1.92 0.92 0.03 0.03 0.03 0.03 0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.01 0.02<	Svalbard and Jan Mayen Islands	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
21.00 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 21.00 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 2.00 1.00 5.00 3.00 0.15 0.17 1.64 0.90 0.90 1.92 1.92 0.02 0.03 9.00 1.00 2.00 2.00 0.10 0.11 1.09 0.60 0.60 1.28 1.28 0.01 0.01 28.00 1.00 6.00 6.00 0.20 0.20 0.21 1.20 1.26 2.56 0.02 0.02 11.00 6.00 5.00 4.00 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03	Swaziland	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
21.00 6.00 4.00 6.00 9.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03 2.00 1.00 5.00 3.00 0.15 0.17 1.64 0.90 0.90 1.92 1.92 0.02 0.02 9.00 1.00 2.00 2.00 0.10 0.11 1.09 0.60 0.60 1.28 1.28 0.01 0.01 28.00 1.00 6.00 6.00 6.00 0.20 0.21 1.20 1.20 2.56 0.02 0.02 11.00 6.00 5.00 4.00 0.13 1.56 0.76 1.58 0.76 0.03	Sweden	21.00	6.00	4.00	00.9	9.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02
2.00 1.00 5.00 3.00 0.15 0.17 1.64 0.90 0.90 1.92 1.92 0.02 0.02 9.00 1.00 2.00 2.00 0.10 0.11 1.09 0.60 0.60 1.28 1.28 0.01 0.01 28.00 1.00 2.00 6.00 6.00 0.20 0.22 2.19 1.20 1.20 2.56 0.02 0.02 11.00 6.00 5.00 4.00 0.19 0.13 1.56 0.76 1.58 1.58 0.02 0.03	Switzerland	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
9.00 1.00 1.00 2.00 0.10 0.11 1.09 0.60 0.60 1.28 1.28 1.28 0.01 0.01 28.00 1.00 2.00 6.00 6.00 6.00 0.20 0.22 2.19 1.20 1.20 2.56 2.56 0.02 0.02 11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 0.76 1.58 1.58 0.02 0.03	Syrian Arab Republic	2.00	1.00	2.00	3.00	3.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
28.00 1.00 2.00 6.00 6.00 6.00 0.20 0.22 2.19 1.20 1.20 2.56 2.56 2.56 0.02 0.02 11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 0.76 1.58 1.58 0.02 0.03	Tajikistan	9.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
11.00 6.00 5.00 3.00 4.00 0.19 0.13 1.56 0.76 0.76 1.58 1.58 0.02 0.03	Thailand	28.00	1.00	2.00	00'9	00.9	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
The second secon	The former Yugoslav Republic	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	92.0	0.76	1.58	1.58	0.02	0.03	0.02	0.02

FAO Country Name	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Horses	Asses Mules		Camels 1	Llamas	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys
Timor-Leste	26.00	1.00	2.00	00.9	00.9	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Togo	1.00	1.00	5.00	1.00	1.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Tokelau	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Tonga	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Trinidad and Tobago	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Tunisia	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Turkey	10.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Turkmenistan	12.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Turks and Caicos Islands	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Tuvalu	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Uganda	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Ukraine	11.00	00.9	5.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
United Arab Emirates	3.00	1.00	5.00	5.00	5.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
United Kingdom	21.00	00.9	4.00	00.9	9.00	0.19	0.13	1.56	0.76	92.0	1.58	1.58	0.02	0.03	0.02	0.02
United Republic of Tanzania	1.00	1.00	5.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
United States of America	48.00	1.00		10.00	19.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
United States Virgin Islands	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Uruguay	1.00	1.00	1.00	1.00	1.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
USSR	11.00	00.9	5.00	3.00	4.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Uzbekistan	11.00	1.00	1.00	2.00	2.00	0.10	0.11	1.09	09.0	09.0	1.28	1.28	0.01	0.01	0.01	0.01
Vanuatu	30.00	2.00	2.00	13.00	24.00	0.15	0.17	1.64	06.0	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Venezuela (Bolivarian Republic of)	2.00	1.00	2.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Viet Nam	24.00	1.00	2.00	5.00	5.00	0.15	0.17	1.64	06.0	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Wallis and Futuna Islands	31.00	2.00	2.00	13.00	24.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Western Sahara	1.00	1.00	5.00	2.00	2.00	0.20	0.22	2.19	1.20	1.20	2.56	2.56	0.02	0.02	0.02	0.02
Yemen	2.00	1.00	5.00	2.00	2.00	0.15	0.17	1.64	0.90	0.90	1.92	1.92	0.02	0.02	0.02	0.02
Yugoslav SFR	11.00	00.9	2.00	3.00	4.00	0.19	0.13	1.56	0.76	0.76	1.58	1.58	0.02	0.03	0.02	0.02
Zambia	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02
Zimbabwe	1.00	1.00	2.00	1.00	1.00	0.15	0.17	1.64	0.90	06.0	1.92	1.92	0.02	0.02	0.02	0.02

TABLE 3A Nitrogen excretion rate (kg N (1000 kg animal mass) $^{-1}$ day $^{-1}$), by animal category and IPCC area

IPCC Area	Cattle, Cattle,	Cattle,	Buffalo Swine,	Swine,	Swine,	Sheep	Goats	Chickens,	Chickens,	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
	dairy	non-dairy		market	breeding			Broilers	Layers							
Indian Subcontinent	0.47	0.34	0.32	0.42	0.24	1.17	1.37	1.10	0.82	0.83	0.74	0.46	0.46	0.46	0.46	0.46
Eastern Europe	0.35	0.35	0.32	0.55	0.46	6.0	1.28	1.10	0.82	0.83	0.74	0:30	0:30	0:30	0.38	0.38
Africa	9.0	0.63	0.32	1.57	0.55	1.17	1.37	1.10	0.82	0.83	0.74	0.46	0.46	0.46	0.46	0.46
Oceania	0.44	0.5	0.32	0.53	0.46	1.13	1.42	1.10	0.82	0.83	0.74	0:30	0:30	0:30	0.38	0.38
Western Europe	0.48	0.33	0.32	0.51	0.42	0.85	1.28	1.10	96:0	0.83	0.74	0.26	0.26	0.26	0.38	0.38
Latin America	0.48	98.0	0.32	1.57	0.55	1.17	1.37	1.10	0.82	0.83	0.74	0.46	0.46	0.46	0.46	0.46
Asia	0.47	0.34	0.32	0.42	0.24	1.17	1.37	1.10	0.82	0.83	0.74	0.46	0.46	0.46	0.46	0.46
Middle east	0.7	0.79	0.32	1.57	0.55	1.17	1.37	1.10	0.82	0.83	0.74	0.46	0.46	0.46	0.46	0.46
Northern America	0.44	0.31	0.32	0.42	0.24	0.42	0.45	1.10	0.83	0.83	0.74	0:30	0:30	0:30	0.38	0.38

TABLE 4A Typical animal mass (kg), by animal category and IPCC area

IDCC A 200	01##10	01#00	D. ff. olo	Curino	Carino	Chickon	Chiokong	Picks	Tirekovo
800000	dairy	non-dairy	Dallalo	market	breeding	Broilers	Cilickelis, Layers	Ducks	idineys
Indian Subcontinent	275	110	295	28	28	6.0	1.8	2.7	6.8
Eastern Europe	550	391	380	20	180	6.0	1.8	2.7	8.9
Africa	275	173	380	28	28	6.0	1.8	2.7	8.9
Oceania	200	330	380	45	180	6.0	1.8	2.7	8.9
Western Europe	009	420	380	50	198	6.0	1.8	2.7	8.9
Latin America	400	305	380	28	28	6.0	1.8	2.7	8.9
Asia	350	391	380	20	180	6.0	1.8	2.7	8.9
Middle east	275	173	380	28	28	6.0	1.8	2.7	8.9
Northern America	604	389	380	46	198	6.0	1.8	2.7	8.9

IPCC Area	Sheep	Goats	Horses	Asses	Mules	Camels	Llamas
Developed	48.50	38.50	377.00	130	130	217	217
Developing	28.00	30.00	238.00	130	130	217	217

TABLE 5A Share of nitrogen manure treated in MS lagoon, by animal category and IPCC area

	dairy non-dairy	Buffalo Swine. market		Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys Horses Asses	Horses	Asses	Mules	Camels	Llamas
Indian 0,000 Subcontinent	000′0 0	0,000	060'0	060'0	0,000	0,000	0,010	0,010	0,010	0,010	000'0	0,000	0,000	0,000	000'0
Eastern Europe 0,000	000′0 0	000'0	0,030	00000	000'0	0,000	000'0	000'0	0,000	000'0	000'0	000'0	000'0	0,000	000'0
Africa 0,000	000′0 0	000'0	000'0	000'0	000'0	000'0	000'0	000'0	000'0	000'0	000'0	000'0	000'0	000'0	000'0
Oceania 0,160	000'0 0	0000'0	0,540	0,540	0,000	0,000	0,000	0,000	0,000	000'0	000'0	000'0	0,000	0,000	0,000
Western Europe 0,000	000′0 0	0000'0	0,087	0,087	0,000	0,000	000'0	0000'0	000'0	000'0	000'0	000'0	0,000	000'0	000'0
Latin America 0,000	000′0 0	000'0	000'0	000'0	000'0	000'0	0,000	000'0	000'0	000'0	000'0	000'0	0,000	000'0	0,000
Asia 0,040	000′0 0	0000'0	0,000	0,000	0,000	0,000	0,010	0,010	0,010	0,010	000'0	000'0	0,000	0,000	0,000
Middle east 0,000	000′0 0	000'0	000'0	000'0	000'0	0,000	000'0	000'0	000'0	000'0	000'0	000'0	000'0	0,000	000'0
Northern America 0,150	000'0 0	0000'0	0,328	0,328	0,000	0,000	0,050	0,050	0,050	0,050	000'0	000'0	0,000	0,000	0,000

TABLE 6A Share of nitrogen manure treated in MS slurry, by animal category and IPCC area

IPCC Area	Cattle.	Cattle.	Buffalo Swine.		Swine.	Sheep Goats	Goats	Chickens.	Chickens. Ducks Turkeys Horses Asses Mules	Ducks	Turkeys	Horses	Asses	Mules	Camels Llamas	Llamas
Indian	0.010	0.010	0.000		0.220	0.000	0.000	0.020	0.020	0.020	0.020	0.000	0.000	0.000	0.000	0.000
Subcontinent																
Eastern Europe	0.175	0.225	0.240	0.000	0.000	0.000	0.000	0.280	0.280	0.280	0.280	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	090.0	090.0	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.010	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.357	0.252	0.200	0.000	0.000	0.000	0.000	0.130	0.130	0.130	0.130	0.000	0.000	0.000	0.000	0.000
Latin America	0.010	0.000	0.000	0.080	0.080	0.000	0.000	0.090	0.090	0.090	060.0	0.000	0.000	0.000	0.000	0.000
Asia	0.380	0.000	0.000	0.400	0.400	0.000	0.000	0.020	0.020	0.020	0.020	0.000	0.000	0.000	0.000	0.000
Middle east	0.010	0.000	0.000	0.140	0.140	0.000	0.000	0.010	0.010	0.010	0.010	0.000	0.000	0.000	0.000	0.000
Northern America	0.270	0.002	0.000	0.185	0.185	0.000	0.000	0.040	0.040	0.040	0.040	0.000	0.000	0.000	0.000	0.000

TABLE 7A Share of nitrogen manure treated in MS solid storage, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo Swine, market		Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Ducks Turkeys Horses Asses Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.000	0.000	0.000	0.160	0.160	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.600	0.440	0.000	0.420	0.420	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Africa	0.010	0.000	0.000	090.0	090.0	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0:030	0.030	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.368	0.390	0.000	0.137	0.137	0.020	0.000	0.010	0.010	0.010	0.010	0.000	0.000	0.000	0.000	0.000
Latin America	0.010	0.000	0.000	0.100	0.100	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.020	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.263	0.000	0.000	0.042	0.042	0.020	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Table 8A Share of nitrogen manure treated in MS dry lot, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo Swine, market		Swine, breeding	Sheep Goats	Goats	Chickens, Broilers	Chickens, Ducks Turkeys Horses Asses Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.000	0.040	0.040	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.010	0.000	0.870	0.870	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.090	0.000	0.150	0.150	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.790	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.410	0.410	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.000	0.460	0.410	0.540	0.540	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.000	0.010	0.000	0.690	0.690	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America 0.000	0.000	0.184	0.000	0.040	0.040	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000

TABLE 9A Share of nitrogen manure treated in MS daily spread, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys Horses Asses Mules Camels	Horses	Asses	Mules		Llamas
Indian Subcontinent	0.190	0.200	0.210	0.090	0.090	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Eastern Europe	0.025	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Africa	0.050	0.010	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Oceania	080.0	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000		0.000	0.000
Western Europe	0.070	0.018	0.000	0.020	0.020	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000		0.000	0.000
Latin America	0.620	0.000	0.000	0.020	0.020	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Asia	0.290	0.020	0.040	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Middle east	0.020	0.020	0.190	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Northern America	0.184	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000		0.000	0.000

TABLE 10A Share of nitrogen manure treated in MS digester, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo Swine, market	Swine, market	Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys Horses Asses Mules Camels	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.010	0.010	0.010	0.080	0.080	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000 0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Asia	0.020	0.000	0.000	0.070	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America 0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000 0.000	0.000	0.000

TABLE 11A Share of nitrogen manure treated in MS pit below one month, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys Horses	Horses	Asses	Mules Camels	Camels	Llamas
Indian Subcontinent	0.000	0.000	0.000	0:030	0.030	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.247	0.247	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.010	0.010	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.028	0.028	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 12A Share of nitrogen manure treated in MS pit above one month, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers		Ducks Turkeys Horses Asses Mules Camels	Horses	Asses	Mules		Llamas
Indian Subcontinent	0.010	0.010	0.000	0.220	0.220	0.000	0.000	0.020	0.020	0.020	0.020	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.175	0.225	0.240	0.000	0.000	0.000	0.000	0.280	0.280	0.280	0.280	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	090'0	090.0	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Oceania	0.010	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.357	0.252	0.200	0.000	0.000	0.000	0.000	0.130	0.130	0.130	0.130	0.000	0.000	0.000	0.000	0.000
Latin America	0.010	0.000	0.000	0.080	0.080	0.000	0.000	060.0	060.0	060.0	060.0	0.000	0.000	0.000	0.000	0.000
Asia	0.380	0.000	0.000	0.400	0.400	0.000	0.000	0.020	0.020	0.020	0.020	0.000	0.000 0.000	0.000	0.000	0.000
Middle east	0.010	0.000	0.000	0.140	0.140	0.000	0.000	0.010	0.010	0.010	0.010	0.000	0.000	0.000	0.000	0.000
Northern America	0.270	0.002	0.000	0.185	0.185	0.000	0.000	0.040	0.040	0.040	0.040	0.000	0.000	0.000 0.000	0.000	0.000

TABLE 13A Share of nitrogen manure treated in other MS, by animal category and IPCC area

IPCC Area	Cattle.	Cattle.	Buffalo	Swine.	Swine.	Sheep Goats	Goats	Chickens.	Chickens. Ducks Turkeys Horses Asses Mules Camels	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
	dairy	non-dairy		market	breeding			Broilers	Layers							
Indian Subcontinent	0.000	0.000	0.000	0.000	0.000	0.170	0.050	0.520	0.520	0.520	0.520	0.050	0.050	0.050	0.050	0.050
Eastern Europe	0.020	0.135	0.470	0.000	0.000	0.270	0.080	0.710	0.710	0.710	0.710	0.080	0.080	0.080 0.080	0.080	0.080
Africa	0.040	0.000	0.000	0.000	0.000	0.010	0.010	0.190	0.190	0.190	0.190	0.010	0.010	0.010	0.010	0.010
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.970	0.970	0.970	0.970	0.000	0.000	0.000	0.000	0.000
Western Europe	0.005	0.020	0.000	0.000	0.000	0.110	0.040	0.840	0.840	0.840	0.840	0.040	0.040	0.040	0.040	0.040
Latin America	0.000	0.010	0.010	0.000	0.000	0.000	0.010	0.490	0.490	0.490	0.490	0.010	0.010	0.010	0.010	0.010
Asia	0.000	0.000	0.000	0.000	0.000	0.170	0.050	0.520	0.520	0.520	0.520	0.050	0.050	0.050	0.050	0.050
Middle east	0.000	0.020	0.190	0.000	0.000	0.000	0.000	0.280	0.280	0.280	0.280	0.000	0.000	0.000 0.000	0.000	0.000
Northern America	0.026	0.000	0.000	0.000	0.000	0.100	0.080	0.900	0.900	0.900	0.900	080.0	0.080	0.080 0.080	0.080	0.080

TABLE 14A Fraction of nitrogen manure from MS slurry applied to soil, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 15A Fraction of nitrogen manure from MS lagoon applied to soil, by animal category and IPCC area

,)													
IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Africa	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Oceania	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Western Europe	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Latin America	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Asia	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Middle east	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000
Northern America	0.350	0.000	0.000	0.400	0.400	0.000	0.000	0.550	0.550	0.550	0.550	0.000	0.000	0.000	0.000	0.000

TABLE 16A Fraction of nitrogen manure from MS solid storage applied to soil, by animal category and IPCC area

•				,												
IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Eastern Europe	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Africa	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Oceania	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Western Europe	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Latin America	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Asia	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Middle east	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120
Northern America	0.300	0.450	0.450	0.450	0.450	0.120	0.120	0.400	0.400	0.400	0.400	0.120	0.120	0.120	0.120	0.120

TABLE 17A Fraction of nitrogen manure from MS dry lot applied to soil, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Turkeys Horses	Asses	Mules Camels	Camels	Llamas
Indian Subcontinent	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.200	0.300	0.300	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 18A Fraction of nitrogen manure from MS daily spread applied to soil, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys	Horses Asses		Mules	Camels	Llamas
Indian Subcontinent	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.070	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 19A Fraction of nitrogen manure from MS digester applied to soil, by animal category and IPCC area

			:			7	,									
IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	lurkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 20A Fraction of nitrogen manure from MS pit below and above one month applied to soil, by animal category and IPCC area

IPCC Area	Cattle, dairy	Cattle, non-dairy	Buffalo	Swine, market	Swine, breeding	Sheep	Goats	Chickens, Broilers	Chickens, Layers	Ducks	Turkeys Horses Asses	Horses	Asses	Mules Camels		Llamas
Indian Subcontinent	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 21A Fraction of nitrogen manure from other MS applied to soil, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.280	0.300	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 22A Emission factor for direct N₂O emissions (kg N₂O-N (kg N)⁻¹), by MS and IPCC area

IDCC A mag	20000	l iquid/Cluzzy	Colid Ctorage	Daylot	Daily Caroad	Digostor	0.4.0	74:0	0+hor
ILCC Area	Lagoon	Liquid/Sidiry	Solid Storage	Dryiot	Dalily Spread	nigester	LIKI	7	Jamo
Indian	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Subcontinent									
Eastern Europe	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Africa	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Oceania	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Western Europe	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Latin America	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Asia	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Middle east	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005
Northern America	0	0.005	0.005	0.02	0	0	0.002	0.002	0.005

TABLE 23A
Percent of nitrogen losses due to runoff and leaching in MS, by IPCC area

IPCC Area	FracLeachMS
Indian Subcontinent	0.10
Eastern Europe	0.10
Africa	0.10
Oceania	0.10
Western Europe	0.10
Latin America	0.10
Asia	0.10
Middle east	0.10
Northern America	0.10

TABLE 24A Emission factor for indirect N₂O emissions from atmospheric deposition (kg N₂O-N (kg NH₃-N + NO_x-N)⁻¹), by IPCC area

IPCC Area	EF4
Indian Subcontinent	0.01
Eastern Europe	0.01
Africa	0.01
Oceania	0.01
Western Europe	0.01
Latin America	0.01
Asia	0.01
Middle east	0.01
Northern America	0.01

TABLE 25A Emission factor for indirect N₂O emissions from leaching and runoff (kg N₂O–N (kg N) $^{-1}$), by IPCC area

IPCC Area	EF5
Indian Subcontinent	0.0075
Eastern Europe	0.0075
Africa	0.0075
Oceania	0.0075
Western Europe	0.0075
Latin America	0.0075
Asia	0.0075
Middle east	0.0075
Northern America	0.0075

TABLE 26A
Emission factor for direct N₂O emissions from nitrogen inputs (kg N₂O-N (kg N input)⁻¹), by IPCC area

IPCC Area	EF1
Indian Subcontinent	0.01
Eastern Europe	0.01
Africa	0.01
Oceania	0.01
Western Europe	0.01
Latin America	0.01
Asia	0.01
Middle east	0.01
Northern America	0.01

TABLE 27A

Percent of nitrogen losses due to runoff and leaching in managed soils and pasture (kg N (kg N additions or deposition, by grazing animals)⁻¹), by IPCC area

IPCC Area	Frac leach
IFCC Alea	riac leacii
Indian Subcontinent	0.30
Eastern Europe	0.30
Africa	0.30
Oceania	0.30
Western Europe	0.30
Latin America	0.30
Asia	0.30
Middle east	0.30
Northern America	0.30

TABLE 28A
Fraction of synthetic nitrogen fertilizer that volatilises (kg N volatilised (kg of N applied)⁻¹), by IPCC area

IPCC Area	Frac_GASF
Indian Subcontinent	0.10
Eastern Europe	0.10
Africa	0.10
Oceania	0.10
Western Europe	0.10
Latin America	0.10
Asia	0.10
Middle east	0.10
Northern America	0.10

TABLE 29A Water regimes (ha), by country

Afghanistan 1 0 0 Albania 1 0 0 Angola 1 0 0 Argentina 1 0 0 Australia 1 0 0 Australia 1 0 0 Bangladesh 0.22 0.7 0.08 Belize 0.1 0 0.9 Benin 0.1 0 0.9 Benin 0.5 0.46 0.04 Bolivia (Plurinational State of) 0.25 0 0.75 Brunei Darussalam 0.79 0 0.21 0 Brunei Darussalam 0.79 0 0.21 0 Burundi 0.25 0 0.75 0 Brundi 0.25 0 0.75 0 Cambodia 0.08 0.9 0.02 0 Cambodia 0.08 0.9 0.02 0 Chad 0.25 0 0.75 </th <th>Country Name</th> <th>irrigated share</th> <th>rainfed share</th> <th>upland share</th>	Country Name	irrigated share	rainfed share	upland share
Angola	Afghanistan	1	0	0
Angola 1 0 0 Argentina 1 0 0 Australia 1 0 0 Bangladesh 0.22 0.7 0.08 Belize 0.1 0 0.9 Benin 0.1 0 0.9 Bhutan 0.5 0.46 0.04 Bolivia (Plurinational State of) 0.25 0 0.75 Brazil 0.19 0.06 0.75 Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burundi 0.25 0 0.75 Cambodia 0.89 0 0.11 Burundi 0.25 0 0.75 Cameron 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameron 0.25 0 0.75 Chad 0.25 0 0.75 Chad 0.25 0 0.75 <th>Albania</th> <th>1</th> <th>0</th> <th>0</th>	Albania	1	0	0
Argentina 1 0 0 0 Australia 1 0 0 0 Bangladesh 0.22 0.7 0.08 Belize 0.1 0 0.9 Benin 0.1 0 0.9 Bhutan 0.5 0.46 0.04 Bolivia (Plurinational State of) Brazil 0.19 0.06 0.75 Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burkina Faso 0.89 0 0.11 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Taiwan 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Cote d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic 0.98 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 Et Salvador 0.1 0.9 Ethiopia 0.5 0.5 France 1 0 0.5	Algeria	1	0	0
Australia 1 0 0 0 Azerbaijan 1 0 0 0 Bangladesh 0.22 0.7 0.08 Belize 0.1 0 0.9 Benin 0.1 0 0.9 Bhutan 0.5 0.46 0.04 Bolivia (Plurinational State of) Brazil 0.19 0.06 0.75 Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burkina Faso 0.89 0 0.11 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Taiwan 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 1 0.00 Democratic Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 Et Salvador 0.1 0 0.9 Ethiopia 0.5 0.5 France 1 0 0 0.5	Angola	1	0	0
Name	Argentina	1	0	0
Bangladesh 0.22 0.7 0.08 Belize 0.1 0 0.9 Benin 0.1 0 0.9 Bhutan 0.5 0.46 0.04 Bolivia (Plurinational State of) 0.25 0 0.75 Brazil 0.19 0.06 0.75 Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chide 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0 Costa Rica 0.	Australia	1	0	0
Belize 0.1 0 0.9 Benin 0.1 0 0.9 Bhutan 0.5 0.46 0.04 Bolivia (Plurinational State of) 0.25 0 0.75 Brazil 0.19 0.06 0.75 Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Cameroon 0.25 0 0.75 Chad 0.25 0 0.75 Chad 0.25 0 0.75 Chad 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 0 0 0 Colombia 0.67 0.1 <th>Azerbaijan</th> <th>1</th> <th>0</th> <th>0</th>	Azerbaijan	1	0	0
Benin 0.1 0 0.9 Bhutan 0.5 0.46 0.04 Bolivia (Plurinational State of) 0.25 0 0.75 Brazil 0.19 0.06 0.75 Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burkina Faso 0.89 0 0.11 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Cameroon 0.25 0 0.75 Cameroon 0.25 0 0.75 Chad 0.25 0 0.75 Chad 0.25 0 0.75 Chad 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Taiwan 1 0 0 Colombia 0.67 0.1	Bangladesh	0.22	0.7	0.08
Bhutan 0.5 0.46 0.04	Belize	0.1	0	0.9
Bolivia (Plurinational State of) 0.25 0 0.75	Benin	0.1	0	0.9
Plurinational State of	Bhutan	0.5	0.46	0.04
Brazil 0.19 0.06 0.75 Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burkina Faso 0.89 0 0.11 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chad 0.25 0 0.75 Chide 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Costa Rica 0.1 0 0 Costa Rica 0.1 0 0 Cote d'Ivoire 0.67 0.2 0.13 Republic of Korea 0 <th>Bolivia</th> <th>0.25</th> <th>0</th> <th>0.75</th>	Bolivia	0.25	0	0.75
Brunei Darussalam 0.79 0 0.21 Bulgaria 1 0 0 Burkina Faso 0.89 0 0.11 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chad 0.25 0 0.75 Chide 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0 Cote d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea				
Bulgaria 1 0 0 Burkina Faso 0.89 0 0.11 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chad 0.25 0 0.75 Chile 0.79 0 0.21 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0			0.06	
Burkina Faso 0.89 0 0.11 Burundi 0.25 0 0.75 Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chad 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.05 0.05 0.9 Democratic Republic of Longo 0.98 0 0.02 E	Brunei Darussalam	0.79	0	0.21
Burundi 0.25 0 0.75		1	0	0
Cambodia 0.08 0.9 0.02 Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chad 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Compose 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of Longo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.1 0 0 <	Burkina Faso	0.89	0	0.11
Cameroon 0.25 0 0.75 Central African Republic 0.25 0 0.75 Chad 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of Longo 0.98 0 0.02 Ecuador 0.4 0.5 0.1 <tr< th=""><th>Burundi</th><th>0.25</th><th>0</th><th>0.75</th></tr<>	Burundi	0.25	0	0.75
Central African Republic 0.25 0 0.75 Chad 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, Hong Kong SAR 1 0 0 China, Taiwan Province of 1 0 0 Coloma Colombia 0.67 0.1 0.23 Comoros 1 0 0 0 Congo 0.25 0 0.75 0 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.5 0 0.5 Fiji 0.5 0 0.5	Cambodia	0.08	0.9	0.02
Chad 0.25 0 0.75 Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, mainland 0.93 0.05 0.02 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of Korea 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.5 Fiji 0.5 0 0.5 France <	Cameroon	0.25	0	0.75
Chile 0.79 0 0.21 China, Hong Kong SAR 1 0 0 China, mainland 0.93 0.05 0.02 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Republic of Korea 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1	Central African Republic	0.25	0	0.75
China, Hong Kong SAR 1 0 0 China, mainland 0.93 0.05 0.02 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Chad	0.25	0	0.75
China, mainland 0.93 0.05 0.02 China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Chile	0.79	0	0.21
China, Taiwan Province of 1 0 0 Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	China, Hong Kong SAR	1	0	0
Province of Colombia 0.67 0.1 0.23 Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	China, mainland	0.93	0.05	0.02
Comoros 1 0 0 Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	•	1	0	0
Congo 0.25 0 0.75 Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Colombia	0.67	0.1	0.23
Costa Rica 0.1 0 0.9 Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Comoros	1	0	0
Côte d'Ivoire 0.06 0.07 0.87 Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.9 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Congo	0.25	0	0.75
Cuba 1 0 0 Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.09 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Costa Rica	0.1	0	0.9
Democratic People's Republic of Korea 0.67 0.2 0.13 Democratic Republic of the Congo 0.05 0.05 0.05 Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Côte d'Ivoire	0.06	0.07	0.87
Democratic Republic of the Congo	Cuba	1	0	0
of the Congo Dominican Republic 0.98 0 0.02 Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	•	0.67	0.2	0.13
Ecuador 0.4 0.5 0.1 Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	-	0.05	0.05	0.9
Egypt 1 0 0 El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Dominican Republic	0.98	0	0.02
El Salvador 0.1 0 0.9 Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Ecuador	0.4	0.5	0.1
Ethiopia 0.5 0 0.5 Fiji 0.5 0 0.5 France 1 0 0	Egypt	1	0	0
Fiji 0.5 0 0.5 France 1 0 0	El Salvador	0.1	0	0.9
France 1 0 0	Ethiopia	0.5	0	0.5
	Fiji	0.5	0	0.5
French Guiana 0.95 0 0.05	France	1	0	0
	French Guiana	0.95	0	0.05
Gabon 0.25 0 0.75	Gabon	0.25	0	0.75

Country Name	irrigated share	rainfed share	upland share
Gambia	0.9	0	0.1
Ghana	0.24	0	0.76
Greece	1	0	0
Guatemala	0.1	0	0.9
Guinea	0.08	0.45	0.47
Guinea-Bissau	0.25	0	0.75
Guyana	0.95	0	0.05
Haiti	0.4	0	0.6
Honduras	0.1	0	0.9
Hungary	1	0	0
India	0.53	0.32	0.15
Indonesia	0.72	0.17	0.11
Islamic Republic of Iran	1	0	0
Iraq	1	0	0
Italy	1	0	0
Jamaica	0.4	0	0.6
Japan	0.99	0	0.01
Kazakhstan	1	0	0
Kenya	0.25	0	0.75
Kyrgyzstan	1	0	0
Lao People's Democratic Republic	0.02	0.61	0.37
Liberia	0	0.06	0.94
Madagascar	0.1	0.76	0.14
Malawi	0.25	0	0.75
Malaysia	0.66	0.22	0.12
Mali	0.25	0	0.75
Mauritania	1	0	0
Mauritius	0.1	0.76	0.14
Mexico	0.41	0	0.59
Micronesia	0.72	0.17	0.11
Morocco	1	0	0
Mozambique Myanmar	0.25 0.18	0.76	0.75
Nepal	0.18	0.76	0.03
Nicaragua	0.23	0.74	0.03
Niger	0.35	0	0.65
Nigeria	0.16	0.33	0.51
Pacific Islands Trust	0.72	0.17	0.11
Territory			
Pakistan	1	0	0
Panama	0.05	0	0.95
Papua New Guinea	0.72	0.17	0.11
Paraguay	0.5	0	0.5
Peru	0.84	0	0.16

Country Name	irrigated share	rainfed share	upland share
Philippines	0.61	0.37	0.02
Portugal	1	0	0
Puerto Rico	0.75	0	0.25
Republic of Korea	1	0	0
Réunion	0.1	0.76	0.14
Romania	1	0	0
Russian Federation	1	0	0
Rwanda	0.25	0	0.75
Saudi Arabia	1	0	0
Senegal	0.25	0	0.75
Sierra Leone	0.01	0.32	0.67
Solomon Islands	0.72	0.17	0.11
Somalia	0.5	0	0.5
South Africa	1	0	0
Spain	1	0	0
Sri Lanka	0.37	0.56	0.07
South Sudan	0.5	0	0.5
Sudan	0.5	0	0.5
Sudan (former)	0.5	0	0.5
Suriname	1	0	0
Swaziland	0.25	0	0.75
Syrian Arab Republic	1	0	0

Country Name	irrigated share	rainfed share	upland share
Tajikistan	1	0	0
Thailand	0.07	0.92	0.01
The former Yugoslav Republic of Macedonia	1	0	0
Timor-Leste	0.72	0.17	0.11
Togo	0.04	0	0.96
Trinidad and Tobago	0.45	0	0.55
Turkey	1	0	0
Turkmenistan	1	0	0
Uganda	0.25	0	0.75
Ukraine	1	0	0
United Republic of Tanzania	0.03	0.75	0.22
United States of America	1	0	0
Uruguay	1	0	0
USSR	1	0	0
Uzbekistan	1	0	0
Venezuela	0.9	0	0.1
Viet Nam	0.53	0.39	0.08
Yugoslav SFR	1	0	0
Zambia	0.25	0	0.75
Zimbabwe	0.25	0	0.75

TABLE 30A Seasonal emission factor of methane(g m-2), by country

Country Name	EF
Afghanistan	10
Albania	20
Algeria	20
Angola	20
Argentina	20
Australia	22.5
Azerbaijan	20
Bangladesh	10
Belize	20
Benin	20
Bhutan	10
Bolivia (Plurinational State of)	20
Brazil	20
Brunei Darussalam	15.7
Bulgaria	20
Burkina Faso	20
Burundi	20
Cambodia	15.7
Cameroon	20
Central African Republic	20

Country Name	EF
Chile	20
China, Hong Kong SAR	13
China, mainland	13
China, Taiwan Province of	13
Colombia	20
Comoros	20
Congo	20
Costa Rica	20
Côte d'Ivoire	20
Cuba	20
Democratic People's Republic of Korea	15.7
Democratic Republic of the Congo	20
Dominican Republic	20
Ecuador	20
Egypt	20
El Salvador	20
Ethiopia	20
Fiji	20
France	36

Country Name	EF
French Guiana	20
Gabon	20
Gambia	20
Ghana	20
Greece	36
Guatemala	20
Guinea	20
Guinea-Bissau	20
Guyana	20
Haiti	20
Honduras	20
Hungary	20
India	10
Indonesia	18
Iran (Islamic Republic of)	15.7
Iraq	20
Italy	36
Jamaica	20
Japan	15
Kazakhstan	15.7
Kenya	20
Kyrgyzstan	15.7
Lao People's Democratic Republic	15.7
Liberia	20
Madagascar	20
Malawi	20
Malaysia	15.7
Mali	20
Mauritania	20
Mauritius	20
Mexico	20
Micronesia	20
Morocco	20
Mozambique	20
Myanmar	15.7
Nepal	10
Nicaragua	20
Niger	20
Nigeria	20
Pacific Islands Trust Territory	20
Pakistan	10
Panama	20
Papua New Guinea	20
Paraguay	20

Carratura Nama	rr.
Country Name	EF
Peru	20
Philippines	27.5
Portugal	36
Puerto Rico	20
Republic of Korea	15
Réunion	20
Romania	20
Russian Federation	20
Rwanda	20
Saudi Arabia	20
Senegal	20
Sierra Leone	20
Solomon Islands	20
Somalia	20
South Africa	20
Spain	36
Sri Lanka	10
South Sudan	20
Sudan	20
Sudan (former)	20
Suriname	20
Swaziland	20
Syrian Arab Republic	20
Tajikistan	15.7
Thailand	16
The former Yugoslav Republic of Macedonia	20
Timor-Leste	15.7
Togo	20
Trinidad and Tobago	20
Turkey	20
Turkmenistan	15.7
Uganda	20
Ukraine	20
United Republic of Tanzania	20
United States of America	25
Uruguay	20
USSR	20
Uzbekistan	15.7
Venezuela (Bolivarian Republic of)	20
Viet Nam	15.7
Yugoslav SFR	20
Zambia	20
Zimbabwe	20

TABLE 31A Amount of nitrogen from bedding (kg N animal $^{-1}$), by animal category and IPCC area

IPCC Area	Cattle.	Cattle.	Buffalo	Swine.	Swine.	Sheep Goats	Goats	Chickens.	Chickens. Ducks Turkeys Horses Asses Mules Camels Llamas	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
	dairy	non-dairy		market	breeding			Broilers	Layers							
Indian Subcontinent	2.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Africa	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Northern America	7.000	4.000	0.000	0.800	5.500	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000 0.000	0.000	0.000

TABLE 32A Fraction of nitrogen manure lost in MS lagoon, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Turkeys Horses Asses		Mules	Camels	Llamas
Indian Subcontinent	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Africa	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Oceania	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Western Europe	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Latin America	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Asia	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Middle east	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000
Northern America	0.770	0.000	0.000	0.780	0.780	0.000	0.000	0.770	0.770	0.770	0.770	0.000	0.000	0.000	0.000	0.000

TABLE 33A Fraction of nitrogen manure lost in MS slurry, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.400	0.000	0.000	0.480	0.480	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 34A Fraction of nitrogen manure lost in MS solid storage, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses	Asses		Mules Camels	Llamas
Indian Subcontinent	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Eastern Europe	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Africa	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Oceania	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Western Europe	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Latin America	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Asia	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Middle east	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150
Northern America	0.400	0.500	0.500	0.500	0.500	0.150	0.150	0.500	0.500	0.500	0.500	0.150	0.150	0.150	0.150	0.150

TABLE 35A Fraction of nitrogen manure lost in MS dry lot, by animal category and IPCC area

	IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep Goats	Goats	Chickens. Broilers	Chickens. Ducks Turkeys Horses Asses Mules Camels Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
ope 0.300 0.400 0.000 0	Indian Subcontinent	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.300 0.400 0.400 0.000 <th< th=""><th>Eastern Europe</th><th>0.300</th><th>0.400</th><th>0.400</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>l</th><th>0.000</th><th>l .</th><th>0.000 0.000</th><th>0.000</th><th>0.000</th></th<>	Eastern Europe	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	l	0.000	l .	0.000 0.000	0.000	0.000
ope 0.300 0.400 0.400 0.000 0	Africa	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000
ope 0.300 0.400 0.400 0.000 0	Oceania	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000	0.000	0.000	0.000	0.000
ia 0.300 0.400 0.400 0.000 0.	Western Europe	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
0.300 0.400 0.400 0.000 <th< th=""><th>Latin America</th><th>0.300</th><th>0.400</th><th>0.400</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th></th><th>0.000</th><th></th><th>0.000 0.000</th><th>0.000</th><th>0.000</th></th<>	Latin America	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000		0.000 0.000	0.000	0.000
0.300 0.400 0.400 0.000 <th< th=""><th>Asia</th><th>0.300</th><th>0.400</th><th>0.400</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000</th><th>0.000 0.000</th><th>0.000</th><th>0.000</th></th<>	Asia	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000
0.300 0.400 0.400 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000	Middle east	0.300	0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000	0.000 0.000 0.000	0.000	0.000	0.000
	Northern America		0.400	0.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000 0.000 0.000	0.000	0.000	0.000

TABLE 36A Fraction of nitrogen manure lost in MS daily spread, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses Asses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.220	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 37A Fraction of nitrogen manure lost in MS pit below and above one month, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo Swine. market		Swine. breeding	Sheep Goats		Chickens. Broilers	Chickens. Ducks Turkeys Horses Asses Mules Camels Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000 0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.250	0.250	0.000 0.000	0.000	0.000	0.000	0.000	0.000 0.000		0.000 0.000 0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000 0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.250	0.250	0.000 0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000 0.000 0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.250	0.250	0.000 0.000	0.000	0.000	0.000	0.000	0.000 0.000		0.000 0.000 0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000 0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.250	0.250	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000 0.000	0.000	0.000
Northern America 0.000	0.000	0.000	0.000	0.250	0.250	0.000 0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000 0.000 0.000	0.000	0.000	0.000

TABLE 38A Fraction of nitrogen manure, urine and dung deposited on pastures that volatilises (kg N volatilised (kg of N applied or deposited)⁻¹), by animal category and IPCC area

IPCC Area	Frac GASM
Indian Subcontinent	0.20
Eastern Europe	0.20
Africa	0.20
Oceania	0.20
Western Europe	0.20
Latin America	0.20
Asia	0.20
Middle east	0.20
Northern America	0.20

TABLE 39A Fraction of managed manure used for feed, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 40A Fraction of managed manure used for fuel, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys Horses Asses Mules Camels	Horses	Asses	Mules	Camels	Llamas
Indian Subcontinent	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000		0.000 0.000	0.000	0.000

TABLE 41A Fraction of managed manure used for construction, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses	Asses	Mules Camels	Camels	Llamas
Indian Subcontinent	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America 0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

TABLE 42A Fraction of nitrogen manure deposited on pastures, ranges and paddocks, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Turkeys	Horses	Asses	Mules Camels	Camels	Llamas
Indian Subcontinent	0.270	0.220	0.190	0.030	0:030	0.830	0.950	0.440	0.440	0.440	0.440	0.950	0.950	0.950	0.950	0.950
Eastern Europe	0.180	0.200	0.290	0.057	0.057	0.730	0.920	0.010	0.010	0.010	0.010	0.920	0.920	0.920	0.920	0.920
Africa	0.830	0.950	0.000	0.000	0.000	0.990	0.990	0.810	0.810	0.810	0.810	0.990	066.0	0.990	0.990	0.990
Oceania	092'0	0.910	0.000	0.280	0.280	1.000	1.000	0.030	0:030	0.030	0.030	1.000	1.000	1.000	1.000	1.000
Western Europe	0.200	0.320	0.000	0.030	0.030	0.870	0.960	0.020	0.020	0.020	0.020	096.0	096.0	0.960	0.960	0.960
Latin America	0.360	066.0	0.990	0.400	0.400	1.000	0.990	0.420	0.420	0.420	0.420	066.0	0.990	0.990	0.990	0.990
Asia	0.200	0.500	0.500	0.000	0.000	0.830	0.950	0.450	0.450	0.450	0.450	0.950	0.950	0.950	0.950	0.950
Middle east	0.800	0.790	0.200	0.000	0.000	1.000	1.000	0.710	0.710	0.710	0.710	1.000	1.000	1.000	1.000	1.000
Northern America	0.108	0.815	0.000	0.000	0.000	0.880	0.920	0.010	0.010	0.010	0.010	0.920	0.920	0.920	0.920	0.920

TABLE 43A Share of manure of dung and urine excreted on fields and burned for fuel, by animal category and IPCC area

IPCC Area	Cattle. dairy	Cattle. non-dairy	Buffalo	Swine. market	Swine. breeding	Sheep	Goats	Chickens. Broilers	Chickens. Layers	Ducks	Ducks Turkeys Horses Asses	Horses	Asses	Mules Camels	Camels	Llamas
Indian Subcontinent	0.510	0.530	0.550	0.000	0.000	0.000	0.000	0.010	0.010	0.010	0.010	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Africa	090.0	0.030	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Oceania	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Western Europe	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Latin America	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Asia	0.070	0.020	0.050	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Middle east	0.170	0.170	0.420	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Northern America 0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000 0.000	0.000	0.000	0.000

TABLE 44A Emission factor for N₂O emissions from urine and dung deposited on pastures (kg N₂O–N (kg N input)⁻¹), by animal category and IPCC area

IPCC Area	Cattle.	Cattle.	Buffalo	Swine.	Swine.	Sheep	Goats	Chickens.	Chickens.	Ducks	Turkeys	Horses	Asses	Mules	Camels	Llamas
	dairy	non-dairy		market	breeding			Broilers	Layers							
Indian Subcontinent	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Eastern Europe	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Africa	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Oceania	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Western Europe	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Latin America	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Asia	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Middle east	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000
Northern America	0.02	0.02	0.02	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.02	0.000	0.000	0.000	0.000	0.000

TABLE 45A Dry matter fraction of harvested product, by crop and IPCC area

IPCC Area	Barley	Beans, dry	Maize	Millet	Oats	Potatoes	Rice, paddy	Rye	Sorghum	Soybeans	Wheat
Indian Subcontinent	0.89	6.0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Eastern Europe	0.89	6.0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Africa	0.89	6:0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Oceania	0.89	6.0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Western Europe	0.89	6:0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Latin America	0.89	6.0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Asia	0.89	6:0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Middle east	0.89	6:0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89
Northern America	0.89	6.0	0.87	6.0	0.89	0.22	0.89	0.88	0.89	0.91	0.89

TABLE 46A Slope element, by crop and IPCC area

IPCC Area	Barley	Beans, dry	Maize	Millet	Oats	Potatoes	Rice, paddy	Rye	Sorghum	Soybeans
Indian Subcontinent	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Eastern Europe	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Africa	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Oceania	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Western Europe	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Latin America	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Asia	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Middle east	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93
Northern America	0.98	0.36	1.03	1.43	0.91	0.1	0.95	1.09	0.88	0.93

TABLE 47A Intercept element, by crop and IPCC area

IPCC Area	Barley	Beans, dry	Maize	Millet	Oats	Potatoes	Rice, paddy	Rye	Sorghum	Soybeans	Wheat
Indian Subcontinent	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Eastern Europe	0.59	89.0	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Africa	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Oceania	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Western Europe	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Latin America	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Asia	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Middle east	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52
Northern America	0.59	0.68	0.61	0.14	0.89	1.06	2.46	0.88	1.33	1.35	0.52

TABLE 48A Ratio of below-ground residues to above-ground biomass, by crop and IPCC area

IPCC Area	Barley	Beans, dry	Maize	Millet	Oats	Potatoes	Rice, paddy	Rye	Sorghum	Soybeans	Wheat
Indian Subcontinent	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Eastern Europe	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Africa	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Oceania	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Western Europe	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Latin America	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Asia	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Middle east	0.22	0	0.22	0	0.25	0.2	0.16	0	0	0.19	0.24
Northern America	000	c	0.22	c	0.25	0.0	0.16	c	c	01.0	0.24

TABLE 49A Combustion factor, by crop and IPCC area

IPCC Area	Barley	Beans, dry	Maize	Millet	Oats	Potatoes	Rice, paddy	Rye	Sorghum	Soybeans	Wheat
Indian Subcontinent	8.0	0.8	8.0	8.0	8.0	8.0	0.8	8.0	0.8	0.8	6.0
Eastern Europe	8.0	0.8	8.0	8.0	8.0	0.8	0.8	8.0	0.8	0.8	6.0
Africa	8.0	0.8	8.0	8.0	8.0	8.0	0.8	8.0	0.8	0.8	6.0
Oceania	8.0	0.8	8.0	8.0	8.0	8.0	0.8	8.0	0.8	0.8	6.0
Western Europe	8.0	0.8	8.0	8.0	8.0	0.8	0.8	0.8	0.8	0.8	6.0
Latin America	8.0	0.8	8.0	8.0	8.0	0.8	0.8	0.8	0.8	0.8	6.0
Asia	8.0	0.8	8.0	8.0	8.0	0.8	0.8	0.8	0.8	0.8	6.0
Middle east	8.0	0.8	8.0	8.0	8.0	8.0	0.8	0.8	0.8	0.8	6.0
Northern America	8.0	0.8	8.0	0.8	8.0	0.8	0.8	0.8	0.8	0.8	6.0

TABLE 50A Nitrogen content of above-ground residues, by crop and IPCC area

IPCC Area	Barley	Beans, dry	Maize	Millet	Oats	Potatoes	Rice, paddy	Rye	Sorghum	Soybeans	Wheat
Indian Subcontinent	0.007	0.01	90000	0.007	0.007	0.019	0.007	0.005	0.007	0.008	900.0
Eastern Europe	0.007	0.01	90000	0.007	0.007	0.019	0.007	0.005	0.007	0.008	0.006
Africa	0.007	0.01	90000	0.007	0.007	0.019	0.007	0.005	0.007	0.008	0.006
Oceania	0.007	0.01	90000	0.007	0.007	0.019	0.007	0.005	0.007	0.008	90000
Western Europe	0.007	0.01	90000	0.007	0.007	0.019	0.007	0.005	0.007	0.008	0.006
Latin America	0.007	0.01	0.006	0.007	0.007	0.019	0.007	0.005	0.007	0.008	0.006
Asia	0.007	0.01	900.0	0.007	0.007	0.019	0.007	0.005	0.007	0.008	90000
Middle east	0.007	0.01	90000	0.007	0.007	0.019	0.007	0.005	0.007	0.008	0.006
Northern America	0.007	0.01	900'0	0.007	0.007	0.019	0.007	0.005	0.007	0.008	0.006

TABLE 51A Ratio of below-ground residues to harvested yield, by crop and IPCC area

IPCC Area	Barley	Beans, dry	Maize	Millet	Oats	Potatoes	Rice, paddy	Rye	Sorghum	Soybeans	Wheat
Indian Subcontinent	0.014	0.01	0.007	0	0.008	0.014	0	0.011	900.0	0.008	600.0
Eastern Europe	0.014	0.01	0.007	0	0.008	0.014	0	0.011	900.0	0.008	600.0
Africa	0.014	0.01	0.007	0	0.008	0.014	0	0.011	900.0	0.008	600.0
Oceania	0.014	0.01	0.007	0	0.008	0.014	0	0.011	0.006	0.008	0.009
Western Europe	0.014	0.01	0.007	0	0.008	0.014	0	0.011	0.006	0.008	600.0
Latin America	0.014	0.01	0.007	0	0.008	0.014	0	0.011	90000	0.008	0.009
Asia	0.014	0.01	0.007	0	0.008	0.014	0	0.011	0.006	0.008	0.009
Middle east	0.014	0.01	0.007	0	0.008	0.014	0	0.011	900.0	0.008	0.009
Northern America	0.014	0.01	0.007	0	0.008	0.014	0	0.011	90000	0.008	0.009

TABLE 52A Fuel biomass consumption value (t d.m ha⁻¹), by crop

Item	Value
Surgarcane	6.5
Maize	10
Rice paddy	5.5
Wheat	4

TABLE 53A Emission factors for burning agricultural residues (g $\,\mathrm{kg}^{-1}\,\mathrm{d.m.}$ burnt)

Parameter Code	Value	Parameter Description
Gef(N ₂ O)	0.07	Emission factor (g kg ⁻¹ d.m. burned)
Gef(CH₄)	2.7	Emission factor (g kg ⁻¹ d.m. burned)

TABLE 54A Emission factors and net calorific value, by energy carrier

Element	Gas-Diesel oil (including gas-diesel oil in fisheries)	Motor gasoline	Natural Gas	LPG	"Residual Fuel Oil (including residual fuel oil in fisheries)"	Coal (Other bituminous coal)	Electricity
Net calorific value [GJ/t]	43.0	44.3		47.3	40.4	26.7	
Emission factor CO ₂ [kg/TJ]	74100	69300	64200	63100	77400	94600	Depending on the national energy mix for electricity generation
Emission factor CH ₄ [kg/TJ]	4.15	80	10	5	10	300	1.2
Emission factor N ₂ O [kg/TJ]	28.6	2.0	0.6	0.1	0.6	1.5	0.01

TABLE 55A Average carbon stock in forest living biomass (t ha⁻¹), by region and year

Years	1990	2000	2005	2010
Regions-Subregions				
Eastern Africa	57.6	58.2	58.5	58.9
Southern Africa	57.6	58.2	58.5	58.9
Northern Africa	21.7	22.1	22.2	22.2
Western Africa	115.4	116.2	116.6	116.9
Central Africa	115.4	116.2	116.6	116.9
Eastern Asia	31.5	33.9	34.5	34.4
Southern Asia	89.5	91.4	88.7	85.6
South-Eastern Asia	89.5	91.4	88.7	85.6
Western Asia	36.4	37.9	38.7	39.8
Central Asia	36.4	37.9	38.7	39.8
Caribbean	65.5	72.4	74.4	74.4
Central America	88.6	89.6	89.9	90.4
Northern America	51.9	53.3	54.1	55
South America	116.5	117.5	117.8	118.2
Australia and New Zealand	54.7	54.5	54.4	54.8
Melanesia	54.7	54.5	54.4	54.8
Micronesia	54.7	54.5	54.4	54.8
Polynesia	54.7	54.5	54.4	54.8
Northern Europe	53.7	58.5	61.2	63.9
Southern Europe	53.7	58.5	61.2	63.9
Eastern Europe	42.7	43.3	43.9	44.8
Western Europe	53.7	58.5	61.2	63.9

TABLE 56A Total carbon stock in forest living biomass (M t), by country

Years	1990	2000	2005	2010
Regions-Subregions				
,	00.0	00.0	20.0	20.0
Afghanistan	38.3	38.3	38.3	38.3
Albania	49.2	49.3	48.3	48.8
Algeria	78	74	72	70
American Samoa	2.02	1.98	1.96	1.94
Andorra				
Angola	4.573.00	4.479.00	4.432.00	4.385.00
Anguilla				
Antigua and Barbuda				
Argentina	3.414.00	3.236.00	3.143.00	3.062.00
Armenia	16.52	14.54	13.54	12.55
Aruba				
Australia	6.724.00	6.702.00	6.641.00	6.606.00
Austria	339	375	399	393
Azerbaijan	54.44	54.44	54.44	54.44
Bahamas				
Bahrain				
Bangladesh	83	82	81	81
Barbados				
Belarus	385.6	481.6	540.4	610.7
Belgium	50.35	60.8	63.08	64.37
Belize	195	183	177	171
Benin	331.59	291.29	276.93	262.56
Bermuda				
Bhutan	296	313	324	336
Bolivia (plurinational state of)	4.877.00	4.666.00	4.561.00	4.442.00
Bosnia and Herzegovina	95.85	117.88	117.88	117.88
Botswana	680	663	655	646
Brazil	68.119.00	65.304.00	63.679.00	62.607.00
British Virgin Islands				
Brunei Darussalam	80.82	76.21	73.9	71.59
Bulgaria	126.6	161.2	181.9	202.1
Burkina Faso	355	323	308	292
Burundi	25.2	19.2	17.6	16.5
Cambodia	609	537	495	464
Cameroon	3.292.00	2.993.00	2.844.00	2.696.00
Canada	14.283.00	14.317.00	14.021.00	13.909.00
Cape Verde	3.3	4.7	4.8	4.9
Cayman Islands				
Central African Republic	2.936.00	2.898.00	2.880.00	2.861.00
Chad	722	677	655	635
Chile	1.293.60	1.328.40	1.338.30	1.348.50
China	4.414.40	5.295.00	5.801.90	6.202.90
Colombia	7.032.00	6.919.00	6.862.00	6.805.00
Comoros	1.65	1.16	0.72	0.35
Cook Islands	38.3	38.3	38.3	38.3
Costa Rica	49.2	49.3	48.3	48.8
Croatia	78	74	72	70
Cuba	2.02	1.98	1.96	1.94
Cyprus				
Czech Republic	4.573.00	4.479.00	4.432.00	4.385.00

Years	1990	2000	2005	2010
Regions-Subregions				
Côte d'Ivoire	1.811.00	1 022 00	1.047.00	1.042.00
Democratic People's Republic of Korea	239	1.832.00	1.847.00	1.842.00
• •	20.433.00	20.036.00	19.838.00	19.639.00
Democratic Republic of the Congo Denmark	20.433.00	25.5	36.2	36.9
Djibouti	0.23	0.23	0.23	0.23
Dominica	0.25	0.23	0.23	0.25
Dominican Republic	113.5	113.5	113.5	113.5
Ecuador				
Egypt	4.38	5.87	6.67	6.96
El Salvador				
Equatorial Guinea	231.9	217.3	210	202.8
Eritrea				
Estonia	168.5283333	168.33	167.28	165.04
Ethiopia	289	254	236	219
Faroe Islands				
Falkland Islands (Malvinas)	-	-	-	-
Fiji				
Finland	720.8	802.4	832.4	832.4
France	965	1.049.00	1.165.00	1.208.00
French Guiana	1.672.00	1.657.00	1.654.00	1.651.00
French Polynesia	20.51	20.51	20.51	20.51
Gabon	2.710.00	2.710.00	2.710.00	2.710.00
Gambia	29.09	30.32	30.94	31.56
Georgia	191.64	202.64	207.45	212.25
Germany	981	1.193.00	1.283.00	1.405.00
Ghana	564	465.3	423.1	380.9
Gibraltar	-	-	-	-
Greece	67	73	76	80
Greenland				
Grenada	1.32	1.32	0.84	1.08
Guadeloupe	13.3	12.8	12.6	12.4
Guam	1.8	1.8	1.8	1.8
Guatemala Guinea	365.2 687	323.6 653	302.9 636	281.3 619
Guinea-Bissau	105	101	99	96
Guyana	1.629.00	1.629.00	1.629.00	1.629.00
Haiti	6.29	5.9	5.69	5.47
Holy See	-	-	-	-
Honduras	517	407	368	330
Hungary	117	130	136	142
Iceland	0.11	0.17	0.22	0.27
India	2.223.00	2.377.00	2.615.00	2.800.00
Indonesia	16.335.00	15.182.00	14.299.00	13.017.00
Iran (Islamic Republic of)	248.6	249.1	253.8	258
	2-0.0	273.1	200.0	230
Iraq	45.0	40.0	40.0	00.0
Ireland	15.8	18.2	19.9	22.6
Isle of Man				
Israel	4.8	4.5	4.6	4.7
Italy	375.3	466.6	512.3	557.9
Jamaica	48.41	48.06	47.83	47.65
Japan	1.159.00	1.381.00	1.526.00	1.722.33

Years	1990	2000	2005	2010
Regions-Subregions				
Jordan	2.36	2.36	2.36	2.36
Kazakhstan	137.42	136.61	136.79	136.79
Kenya	525.3	502.7	488.7	476.2
Kiribati				
Kuwait				
Kyrgyzstan	26.9	33.7	37	56
Lao People's Democratic Republic	1.186.00	1.133.00	1.106.00	1.074.00
Latvia	193.48	234.23	244.11	271.56
Lebanon	1.65	1.66	1.66	1.73
Lesotho	2.11	2.21	2.27	2.32
Liberia	666.02	625.48	605.21	584.94
Libya	6.05	6.05	6.05	6.05
Liechtenstein	0.48	0.51	0.51	0.51
Lithuania	134.1	145.6	150.8	152.5
Luxembourg	7.35	9.36	9.36	9.36
Madagascar	1.778.00	1.691.00	1.663.00	1.626.00
Malawi	173	159	151	144
Malaysia	2.822.00	3.558.00	3.361.00	3.212.00
Maldives				
Mali	317.42	299.58	290.66	281.74
Malta	0.06	0.06	0.06	0.06
Marshall Islands	2.31	2.31	2.31	2.31
Martinique	8.4	8.4	8.4	8.4
Mauritania	12.5	9.5	8	7.3
Mauritius	2.53	2.52	2.27	2.27
Mayotte				
Mexico	2.186.00	2.111.00	2.076.00	2.043.00
Micronesia (Federated States of)	20.2	20.29	20.33	20.38
Monaco Mongolia	671	626	605	583
Montenegro	33.3	33.3	33.3	33.3
Montserrat	33.3	33.3	33.3	33.3
Morocco	189.88	211.97	224.19	222.78
Mozambique	1.878.00	1.782.00	1.733.00	1.692.00
Myanmar	2.040.00	1.814.00	1.734.00	1.653.00
Namibia	252.77	231.71	220.99	210.23
Nauru	-	-	-	_
Nepal	602	520	484	484
Netherlands	20.5	24	25.6	27.7
Netherlands Antilles				
New Caledonia	60.49	60.49	60.49	60.49
New Zealand	1.176.00	1.234.00	1.263.00	1.292.00
Nicaragua	506.3	427.8	388.5	349.3
Niger	60	41	38	37
Nigeria	2.016.00	1.550.00	1.317.00	1.085.00
	2.010.00	1.330.00	1.317.00	1.000.00
Niue				
Norfolk Island				
Northern Mariana Islands	3.37	3.21	3.12	3.04
Norway	280	323	360	395
Occupied Palestinian Territory				
Oman				

Regions-Subregions Pakista 330 271 243 213 268 10.66 20.00 20.00 20.00 20.00 20.00 20.00 10.00	Years	1990	2000	2005	2010
Palau 330 271 243 213 Palau 10.09 10.47 10.66 10.68 Panama 428.6 380.7 374 387.4 Papua New Guinea 2.536.80 2.422.50 2.365.40 2.306.20 Paraguay Per 8.832.00 8.713.00 8.654.00 8.559.00 Philippines 641.39 655.35 660.12 663.42 Polad Poland 691 807 887 968 Poland 102.00 102.00 102 102 102 Poland 10.20 10.20 102<	Regions-Subregions				
Palau 10.09 10.47 10.66 10.66 Panama 428.6 380.7 374 367.4 Papua New Guinea 2.586.80 2.42.50 2.365.40 2.306.20 Paraguay Peru 8.832.00 8.713.00 8.654.00 8.559.00 Philippines 641.33 655.35 660.12 682.00 Poland 691 807 887 988 Poland 102.00 102.00 102	•	330	271	243	213
Panama 428.6 380.7 374 367.4 Papua New Guinea 2.536.80 2.422.50 2.366.40 2.306.20 Peru 8.832.00 8.713.00 8.654.00 8.555.00 Philippines 641.39 655.35 660.12 663.42 Potudal 691 807 887 988 Portugal 102.00 102.00 102 102 Querto Rico 14.24 23.47 25.76 28.05 Gater 1 2 24 268 Republic of Korea 109 181 224 268 Republic of Moldova 22.27 26.09 275 28.9 Republic of Moldova 22.27 26.09 275 28.9 Republic of Moldova 32.504.00 32.15700 32.210.00 32.500.00 Republic of Moldova 22.27 26.09 275 28.9 Republic of Moldova 32.500.00 32.15700 32.150.00 32.150.00 32.150.00 32.150.00 <th></th> <th></th> <th></th> <th></th> <th>-</th>					-
Papus New Guinea 2.536.80 2.422.50 2.365.40 2.306.20 Paraguay Peru 8.832.00 8.713.00 8.654.00 8.559.00 Philippines 641.39 655.35 660.12 663.42 Poland 691 807 887 988 Portugal 102.00 102.00 102 102 Puerto Rico 14.24 23.47 25.76 28.05 Catar Republic of Korea 109 181 . 22.4 . 28.05 Republic of Moldova 22.27 26.09 . 25.5 . <th>Panama</th> <th></th> <th></th> <th></th> <th>367.4</th>	Panama				367.4
Paraguay	Papua New Guinea				2.306.20
Peru 8.832.00 8.713.00 8.654.00 8.559.00 Philippines 641.39 655.35 660.12 663.42 Poland 691 807 887 988 Portugal 102.00 102.00 102 102 Puerto Rico 14.24 23.47 25.76 28.05 Catar 1 2 2 25.76 28.05 Republic of Korea 109 181 224 268 Republic of Moldova 22.27 26.09 22.5 28.9 Romania 599.72 599.25 60.13 618.05 Resian Federation 32.504.00 32.150.00 32.210.00 32.504.00 32.150.00 32.210.00 32.504.00 32.150.00 32.210.00 32.504.00 32.150.00 32.210.00 32.504.00 32.150.00 32.210.00 32.504.00 32.150.00 32.210.00 32.504.00 32.150.00 32.210.00 32.504.00 32.150.00 32.150.00 32.210.00 32.504.00 32.150.00 <th< th=""><th>•</th><th></th><th></th><th></th><th></th></th<>	•				
Poland Family F		8.832.00	8.713.00	8.654.00	8.559.00
Portugal 102.00 102.00 102 102 Puerto Rico 14.24 23.47 25.76 28.05 Qatar -	Philippines	641.39	655.35	660.12	663.42
Puerto Rico 14.24 23.47 25.76 28.05 Qatar - - - - - Republic of Moldova 22.27 26.09 27.5 28.9 Romania 599.72 599.25 601.13 618.05 Russian Federation 32.504.00 32.157.00 32.210.00 32.500.00 Rwanda 35.2 17.58 34.55 39.48 Réunion 6 6 5.9 6 Saint Lucia 5 5.9 6 Saint Lucia 5 5.9 6 Saint Lucia 5 5.9 6 Saint Mincent and the Grenadines 5 5.9 6 Sam Mario - - - - - Sand Tome and Principe 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81 3.81	Poland	691	807	887	968
Page	Portugal	102.00	102.00	102	102
Republic of Korea 109	Puerto Rico	14.24	23.47	25.76	28.05
Republic of Moldova 22.27 26.09 27.5 28.9	Qatar	-	-	-	-
Romania 599.72 599.25 601.13 618.05 Russian Federation 32.504.00 32.157.00 32.210.00 32.500.00 Rwanda 35.2 17.58 34.55 39.48 Révinion 6 6 5.9 6 Saint Kitts and Nevis Saint Pierre and Miquelon Saint Pierre and Miquelon Saint Pierre and Miquelon Saint Marino -	Republic of Korea	109	181	224	268
Russian Federation 32.504.00 32.157.00 32.210.00 32.500.00 Rwanda 35.2 17.58 34.55 39.48 Réunion 6 6 5.9 6 Saint Kitts and Nevis Saint Lucia Saint Pierre and Miquelon Saint Vincent and the Grenadines Samoa Samoa Sam Marino - - - - Sa o Tome and Principe 3.81 3.81 3.81 3.81 Saudi Arabia 5.93 5.93 5.93 5.93 Seychelles 3.57 3.57 3.48 340 Seychelles 3.57 3.57 3.48 340 Signapore 247.13 231.6 223.83 216.07 Singapore 116 140.7 159.2 178.3 Slovakia 162.7 189.8 202.4 211.2 Slovania 483 438 416 184.2 181.9 Somalia 483 438 415 394	Republic of Moldova	22.27	26.09	27.5	28.9
Réunion 35.2 17.58 34.55 39.48 Réunion 6 6 5.9 6 Saint Kitts and Nevis Saint Lucia Saint Vincent and the Grenadines Samoa Samoa Sam Marino - <th< th=""><th>Romania</th><th>599.72</th><th>599.25</th><th>601.13</th><th>618.05</th></th<>	Romania	599.72	599.25	601.13	618.05
Réunion 6 6 5.9 6 Saint Kitts and Nevis Saint Lucia Saint Pierre and Miquelon Samoa Samoa	Russian Federation	32.504.00	32.157.00	32.210.00	32.500.00
Saint Kitts and Nevis Saint Pierre and Miquelon Saint Pierre and Miquelon Saint Vincent and the Grenadines Samoa Sam Marino -	Rwanda	35.2	17.58	34.55	39.48
Saint Lucia Saint Pierre and Miquelon Saint Vincent and the Grenadines Samoa San Marino -	Réunion	6	6	5.9	6
Saint Pierre and Miquelon Samoa	Saint Kitts and Nevis				
Saint Vincent and the Grenadines Samoa San Marino -	Saint Lucia				
Samoa - <th>Saint Pierre and Miquelon</th> <th></th> <th></th> <th></th> <th></th>	Saint Pierre and Miquelon				
San Marino - - - - Sao Tome and Principe 3.81 3.81 3.81 3.81 Saudi Arabia 5.93 5.93 5.93 5.93 Senegal 3.77 357 348 340 Seychelles 3.57 3.57 3.57 3.57 Sierra Leone 247.13 231.6 223.83 216.07 Singapore 3.57 189.8 202.4 211.2 Slovakia 162.7 189.8 202.4 211.2 Slovania 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.3	Saint Vincent and the Grenadines				
Sao Tome and Principe 3.81 3.81 3.81 3.81 Saudi Arabia 5.93 5.93 5.93 5.93 Senegal 3.77 357 348 340 Seychelles 3.57 3.57 3.57 3.57 Sierra Leone 247.13 231.6 223.83 216.07 Singapore 3.57 189.8 202.4 211.2 Slovakia 162.7 189.8 202.4 211.2 Slovenia 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Swaziland 22.6 22.3 22.1	Samoa				
Saudi Arabia 5.93 5.93 5.93 5.93 Senegal 377 357 348 340 Seychelles 3.57 3.57 3.57 3.57 Sierra Leone 247.13 231.6 223.83 216.07 Singapore Slovakia 162.7 189.8 202.4 211.2 Slovania 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394.3 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 66.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 <th>San Marino</th> <th>-</th> <th>-</th> <th>-</th> <th>-</th>	San Marino	-	-	-	-
Senegal 377 357 348 340 Seychelles 3.57 3.57 3.57 3.57 Sierra Leone 247.13 231.6 223.83 216.07 Singapore Slovakia 162.7 189.8 202.4 211.2 Slovenia 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.164.50 Swaziland 2.6 22.3 22.1 22 28 28 28 28	Sao Tome and Principe	3.81	3.81	3.81	3.81
Seychelles 3.57 3.57 3.57 3.57 Sierra Leone 247.13 231.6 223.83 216.07 Singapore Slovakia 162.7 189.8 202.4 211.2 Slovenia 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 Swaziland 22.6 22.3 22.1 22 Sweden 1.178.10 1.182.70 1.219.00 1.255.30 Switzerland 3.05 2.8 2.8 2.8	Saudi Arabia	5.93	5.93	5.93	5.93
Sierra Leone 247.13 231.6 223.83 216.07 Singapore Slovakia 162.7 189.8 202.4 211.2 Slovenia 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.164.50 Swaziland 22.6 22.3 22.1 22 Sweden 1.178.10 1.182.70 1.219.00 1.255.30 Switzerland 126 135 139 143 Syrian Arab Republic 60.1 62.1 60.4	Senegal	377	357	348	340
Singapore Slovakia 162.7 189.8 202.4 211.2 Slovenia 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.50 Swaziland 22.6 22.3 22.1 22 Sweden 1.178.10 1.182.70 1.219.00 1.255.30 Switzerland 126 135 139 143 Syrian Arab Republic 60.1 62.1 60.4 60.4 of Macedonia 60.1 62.1 <th< th=""><th>Seychelles</th><th>3.57</th><th>3.57</th><th>3.57</th><th>3.57</th></th<>	Seychelles	3.57	3.57	3.57	3.57
Slovakia 162.7 189.8 202.4 211.2 Slovenia 116 140.7 159.2 178.3 Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.164.50 Swaziland 22.6 22.3 22.1 22 Sweden 1.178.10 1.182.70 1.219.00 1.255.30 Switzerland 126 135 139 143 Syrian Arab Republic 3.05 2.8 2.8 2.8 Thailand 908 882 877 881 The former Yugoslav Republic 60.1 </th <th>Sierra Leone</th> <th>247.13</th> <th>231.6</th> <th>223.83</th> <th>216.07</th>	Sierra Leone	247.13	231.6	223.83	216.07
Slovenia					
Solomon Islands 191 186.4 184.2 181.9 Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 <th></th> <th></th> <th></th> <th></th> <th></th>					
Somalia 483 438 415 394 South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 <					
South Africa 806.9 806.9 806.9 806.9 Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30					
Spain 289.15 396.13 399.56 421.84 Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.168.30 3.164.50 Svalbard and Jan Mayen Islands -					
Sri Lanka 89.63 74.12 65.94 60.85 Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.168.30 3.164.50 Svalbard and Jan Mayen Islands -					
Sudan (former) 1.520.00 1.403.00 1.398.00 1.392.00 Suriname 3.168.30 3.168.30 3.168.30 3.164.50 Svalbard and Jan Mayen Islands - <th< th=""><th><u> </u></th><th></th><th></th><th></th><th></th></th<>	<u> </u>				
Suriname 3.168.30					
Svalbard and Jan Mayen Islands - <th< th=""><th></th><th></th><th></th><th></th><th></th></th<>					
Swaziland 22.6 22.3 22.1 22 Sweden 1.178.10 1.182.70 1.219.00 1.255.30 Switzerland 126 135 139 143 Syrian Arab Republic Tajikistan 3.05 2.8 2.8 2.8 Thailand 908 882 877 881 The former Yugoslav Republic of Macedonia 60.1 62.1 60.4 60.4 Timor-Leste Togo 7 -		3.100.30	3.100.30	3.100.30	3.104.50
Sweden 1.178.10 1.182.70 1.219.00 1.255.30 Switzerland 126 135 139 143 Syrian Arab Republic 3.05 2.8 2.8 2.8 Tajikistan 3.05 2.8 2.8 2.8 Thailand 908 882 877 881 The former Yugoslav Republic of Macedonia 60.1 62.1 60.4 60.4 Timor-Leste Togo Tokelau -	<u> </u>	22.6	22.3	22 1	22
Switzerland 126 135 139 143 Syrian Arab Republic 3.05 2.8 2.8 2.8 Thailand 908 882 877 881 The former Yugoslav Republic of Macedonia 60.1 62.1 60.4 60.4 Timor-Leste Togo Tokelau - - - - - - Tonga 1.03 1.03 1.03 1.03 1.03 Trinidad and Tobago 20.54 19.75 19.62 19.17					
Syrian Arab Republic Tajikistan 3.05 2.8 2.8 2.8 2.8					
Tajikistan 3.05 2.8 2.8 2.8 Thailand 908 882 877 881 The former Yugoslav Republic of Macedonia 60.1 62.1 60.4 60.4 Timor-Leste Togo 7 -					
Thailand 908 882 877 881 The former Yugoslav Republic of Macedonia 60.1 62.1 60.4 60.4 Timor-Leste Togo -		3.05	2.8	2.8	2.8
of Macedonia Timor-Leste Togo Tokelau -	<u> </u>		882	877	881
of Macedonia Timor-Leste Togo Tokelau -					
Togo Tokelau -					
Tokelau - </th <th>Timor-Leste</th> <th></th> <th></th> <th></th> <th></th>	Timor-Leste				
Tonga 1.03 1.03 1.03 1.03 Trinidad and Tobago 20.54 19.75 19.62 19.17	Togo				
Trinidad and Tobago 20.54 19.75 19.62 19.17	Tokelau	-	-	-	-
	Tonga	1.03	1.03	1.03	1.03
Tunisia 5.8 7.5 8.2 9.3	Trinidad and Tobago	20.54	19.75	19.62	19.17
	Tunisia	5.8	7.5	8.2	9.3

Years	1990	2000	2005	2010
Regions-Subregions				
Turkey	686.14	742.62	782.23	821.85
Turkmenistan	11.3	11.3	11.7	11.7
Turks and Caicos Islands				
Tuvalu				
Uganda	171.3	139.7	123.9	108.8
Ukraine	499	662	711.5	761
United Arab Emirates	12.24	15.49	15.56	15.85
United Kingdom	120	119	128	136
United Republic of Tanzania	2.505.00	2.262.00	2.139.00	2.019.00
United States Virgin Islands	0.6	0.57	0.56	0.54
United States of America	16.950.00	17.998.00	18.632.00	19.308.00
Uruguay				
Uzbekistan	7.8	14	17.5	19.3
Vanuatu				
Venezuela (Bolivarian Republic of)				
Viet Nam	777	927	960	992
Wallis and Futuna Islands				
Western Sahara	32.65	32.65	32.65	32.65
Yemen	5.16	5.16	5.16	5.16
Zambia	2.578.68	2.497.32	2.456.64	2.415.96
Zimbabwe	697.23	594.27	542.8	491.51

TABLE 57A Emission factor for N_2O emissions from cropland and grassland organic soils, by climatic class $\,$

Climatic Class	Climatic Class Name	EF: N ₂ O-N (Kg ha ⁻¹ yr ⁻¹)
1	WarmTemperate Moist	8
2	WarmTemperate Dry	8
3	CoolTemperate Moist	8
4	Cool Temperate Dry	8
5	Polar Moist	8
6	Polar Dry	8
7	Boreal Moist	8
8	Boreal Dry	8
9	Tropical Montane	16
10	Tropical Wet	16
11	Tropical Moist	16
12	Tropical Dry	16

TABLE 58A Emission factor for C emissions from cropland organic soils, by climatic class

Climatic Class	Climatic Class Name	EF: C (tonnes ha ⁻¹ yr ⁻¹)
1	Warm Temperate Moist	10
2	Warm Temperate Dry	10
3	CoolTemperate Moist	5
4	Cool Temperate Dry	5
5	Polar Moist	5
6	Polar Dry	5
7	Boreal Moist	5
8	Boreal Dry	5
9	Tropical Montane	20
10	Tropical Wet	20
11	Tropical Moist	20
12	Tropical Dry	20

TABLE 59A Emission factor for C emissions from grassland organic soils, by climatic class

Climatic Class	Climatic Class Name	EF: C (tonnes ha ⁻¹ yr ⁻¹)
1	WarmTemperate Moist	2.5
2	Warm Temperate Dry	2.5
3	CoolTemperate Moist	0.25
4	Cool Temperate Dry	0.25
5	Polar Moist	0.25
6	Polar Dry	0.25
7	Boreal Moist	0.25
8	Boreal Dry	0.25
9	Tropical Montane	5
10	Tropical Wet	5
11	Tropical Moist	5
12	Tropical Dry	5

TABLE 60A Correspondence between IPCC and FAOSTAT Nomenclatures of Animals categories

	IPCC		FAOSTAT Emissions database	
Livestock population	Main categories	Animal / Subcategories	Animal category	Livestock category
	Mature	High-producing dairy cows that have calved at least once and are used principally for milk production		Cattle*
	dairy cow	Low-producing dairy cows that have calved at least once and are used principally for milk production		
		Cows used principally for producing meat	Cattle, dairy	
	Other mature cattle	Cows used for more than one production purpose: milk, meat, draft		
Cattle	mature cattle	Bulls used principally for breeding purposes		
		Bullocks used principally for draft power		
		Calves pre-weaning		
	Crowing	Replacement dairy heifers	Cattle	
	Growing cattle	Growing / fattening cattle post-weaning	Cattle, non-dairy	
		Feedlot-fed cattle on diets containing > 90 % concentrates		
		High-producing dairy buffalo that have calved at least once and are used principally for milk production	Buffaloes	Buffaloes*
d	Mature dairy buffalo	Low-producing dairy buffalo that have calved at least once and are used principally for milk production		
		Buffalos used to produce offspring for meat		
Buffalo	Mature	Buffalos used for more than one production purpose: milk, meat, draft		
	non-dairy buffalo	Bulls used principally for breeding purposes		
	banaio	Bullocks used principally for draft power		
		Calves pre-weaning		
	Carrier a	Replacement dairy heifers		
	Growing buffalo	Growing / fattening buffalo post-weaning		
		Feedlot-fed buffalo on diets containing > 90 % concentrates		
	N/atoma access	Breeding ewes for production of offspring and wool production	Sheep	Sheep*
Sheep Other mat sheep (>1 y	Mature ewes	Milking ewes where commercial milk production is the primary purpose		
	Other mature sheep (>1 year)	No further sub-categorisation recommended		
		Intact male		
	Growing lambs	Castrates		
		Females		
	Mature goat	Milking goat where commercial milk production is the primary purpose	. Goats Go	Goats*
	Other mature goat (>1 year)	Intact males		
		Castrates		
	Growing goat	Females		
		Goats in gestation		

	IPCC		FAOSTAT Emissions database	
Livestock population	Main categories	Animal / Subcategories	Animal category	Livestock category
Swine	Mature swine (breeding)	Sows in gestation Sows which have farrowed and are nursing young Boars that are used for breeding purposes	Swine, breeding	Pigs
	Growing swine	Nursery Gilts that will be used for breeding purposes Growing boars that will be used for		
	Market growing swine	breeding purposes Finishing	Swine, market	
		Broiler chickens grown for producing meat	Chickens, broilers	
Chickens	Chickens	Layer chickens for producing eggs, where manure is managed in dry systems (e.g., high-rise houses)	Chickens, layers	Chickens*
		Layer chickens for producing eggs, where manure is managed in wet systems (e.g., lagoons)		
		Chickens under free-range conditions for egg or meat production		
Turkeys	Turkeys	Breeding turkeys in confinement systems Turkeys grown for producing meat in confinement systems Turkeys under free-range conditions for	Turkeys	Turkeys
		meat production Breeding ducks		
Ducks	Ducks	Ducks grown for producing meat	Ducks	Ducks
Camels	Camels	Camels	Camels	Camels
and llamas	Llamas, Alpacas	Llamas, Alpacas	Llamas –	Camelids, other
Mules	Mules and Asses	Mules and Asses	Mules	Mules
and Asses			Asses	Asses
Horses	Horses	Horses	Horses	Horses
Others	Rabbits	Rabbits Geese	_	Rabbits and hares Geese and guinea
	Geese		_	fowls
	Fur bearing animals	Fur bearing animals	-	Rodents, other
	Ostrich	Ostrich		Animals live NES
	Deer	Deer	-	(not else specified)

^{*} FAOSTAT database includes data subset of cattle, buffalos, sheep, goat and camels related to the milk animals, a subset of chickens related to laying animals for Supply Utilization Account/Food Balance Sheet purpose, although they are not considered as categories of the FAOSTAT commodity list.

Annex 4

OVERVIEW OF EXISTING REPORTING REQUIREMENTS FOR MITIGATION IN THE AGRICULTURE, FORESTRY AND OTHER LAND USE SECTOR UNDER THE UNFCCC

UN Member States meeting at the United Nations Conference on Environment and Development (UNCED, Rio de Janeiro, 1992), agreed to sign a Declaration on sustainable development as an instrument to ensure a healthy and productive life for human beings, in harmony with nature, for current and future generations. Sustainable Development commitments of UN Member States include the need to ensure that activities within their jurisdiction or control do not cause damage to the environment, both within their territories as well as of other States or areas beyond the limits of their national jurisdiction.

For these reasons, UNCED signatories also signed the United Nations Framework Convention on Climate Change (UNFCCC)¹⁰, with the commitment to *stabilize greenhouse gas* (*GHG*) concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system, within a time-frame sufficient to allow ecosystems to adapt naturally to climate change, to ensure that food production is not threatened, and to enable economic development to proceed in a sustainable manner (UNFCCC, Art. 2) To achieve these global sustainable development goals in the Agriculture, Forestry and Other Land Use sector (AFOLU), Parties to the UNFCCC shall promote sustainable management, and promote and cooperate in the conservation and enhancement, as appropriate, of sinks and reservoirs, including biomass, forests, as well as other terrestrial and coastal ecosystems.

At the same time, sustainable rural development and eradication of hunger, as specified in both UNCED and UNFCCC core principles, are at the basis of FAO mission¹¹. In particular, FAO generates knowledge and provides assistance to its member countries towards appropriate adaptation and mitigation responses in the agriculture, fisheries and forestry sectors, ¹² to ensure sustainability of production systems and maintain food security under climate change.

Parties to the UNFCCC have committed to implement specific actions and programs to mitigate climate change. In particular, UNFCCC Art. 4 states:

- "All Parties, taking into account their common but differentiated responsibilities and their specific national and regional development priorities, objectives and circumstances, shall:
- (a) Develop, periodically update, publish and make available to the Conference of the Parties... ...national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol, using comparable methodologies to be agreed upon by the Conference of the Parties;
- (b) Formulate, implement, publish and regularly update national and, where appropriate, regional programmes containing measures to mitigate climate change by addressing anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol...
- (c) Communicate to the Conference of the Parties information related to implementation..."

⁹ http://www.un.org/documents/ga/conf151/aconf15126-1annex1.htm

 $^{^{10}\} http://unfccc.int/files/essential_background/background_publications_htmlpdf/application/pdf/conveng.pdf$

¹¹ http://www.fao.org/docrep/x3551e/x3551e02.htm

¹² Climate Change and Food Security: A Framework document – FAO 2008; available at: http://www.fao.org/forestry/15538-079b31d45081f-e9c3dbc6ff34de4807e4.pdf

From Art. 4 of the Convention, different reporting requirements have emerged for developed and developing countries (UNFCCC Annex I and non-Annex I Parties, respectively)¹³.

This note provides information on UNFCCC reporting requirements associated to mitigation of climate change in the AFOLU sector for developed and developing countries. Within the AFOLU sector, mitigation consists of actions taken to reduce GHG emissions and enhance carbon sinks and reservoirs, compared to a business-as-usual scenario/ reference level. Parties need to report these actions periodically to the UNFCCC, as specified in a series of decisions agreed upon at the annual Conference of Parties (COP) and summarized in Tab. 1 in the Appendix. Specifically:

Annex I Parties Report:

- National Communications (NC), containing information, from the last submitted GHG Inventory, on national GHG emissions/removals, climate-related policies and measures, GHG projections, vulnerability and adaptation to climate change, financial assistance and technology transfer to non-Annex I Parties, and actions to raise public awareness on climate change;
- National GHG Inventories (NGHGI), containing information on GHG emissions and removals, such as activity data, emission factors, and methodologies used to estimate these emissions. A NGHGI is composed of two distinct documents: the Common Reporting Format (CRF) tables, which contain a time-series of GHG emission estimates (from 1990 till the year x-2; where x is the submission year of the NGHGI); and the National Inventory Report (NIR), which includes all information on background data and methods used, and the data analysis and institutional arrangements underlying the preparation of the NGHGI.
- Biennial Reports (BRs), which outline progress in achieving net emissions reductions and provision of financial, technological, and capacity-building support to non-Annex I Parties for dealing with climate change.

National communications are to be submitted by Annex I Parties every 4 years ¹⁴ (decisions 8/CP.1, 11/CP.4). They are prepared and reported periodically by Annex I Parties based on agreed reporting guidelines (decision 4/CP.5)¹⁵, and, for the GHG estimates, on methodology developed by the Intergovernmental Panel on Climate Change (IPCC), as adopted for the NGHGI by the COP. Submissions by Annex I Parties can be found here: http://unfccc.int/national_reports/annex_i_natcom/submitted_natcom/items/4903.php

National GHG inventories are submitted (3/CP.5) by Annex I Parties annually. They are prepared on the basis of reporting guidelines¹⁶ agreed by the COP (24/CP.19) and on methodologies developed by IPCC¹⁷. National inventory arrangements should be in place to ensure that a NGHGI is fully compliant with reporting requirements and is submitted on time. Submissions by Annex I Parties can be found here: http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/8108.php

¹³ Annex I Parties include the industrialized countries that were members of the OECD in 1992, plus countries with economies in transition (EIT); Non-Annex I Parties are mostly developing countries.

¹⁴ The exact timing depends on specific decisions taken by the Conference of the Parties (COP), which establishes the due date for each subsequent submission; this could vary between 3-5 years.

¹⁵ Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, Part II: UNFCCC reporting guidelines on national communications (pages 80-100), Annex to Decision 4/CP.5 at http://unfccc.int/resource/docs/cop5/07.pdf

Annex I Parties use the Guidelines for the preparation of national communications by Parties included in Annex I to the Convention, Part I: UNFCCC reporting guidelines on annual greenhouse gas inventories (Annex I to Decision 24/CP.19) at http://unfccc.int/resource/docs/2013/cop19/eng/10a03.pdf#page=2

Annex I Parties will use the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (at http://www.ipcc-nggip.iges.or.jp/pub-lic/ 2006gl/index.html), may also use the 2013 Supplement to the 2006 Guidelines for National Greenhouse Gas Inventories: Wetlands (Wetlands Supplement) (at http://www.ipcc-nggip.iges.or.jp/public/wetlands/index.html) and for the KP reporting they will use the 2013 Revised Supplementary Methods and Good Practice Guidance Arising from the Kyoto Protocol (KP Supplement) (at http://www.ipcc-nggip.iges.or.jp/public/kpsg/index.html)

Biennial Reports are to be submitted (2/CP.17) by developed country Parties every 2 years; the first biennial report (BR1) is due by 1 January 2014¹⁸. They are prepared on the basis of agreed reporting guidelines (2/CP.17 Annex I) and on methodologies developed by IPCC, as per NGHGI. Submissions by Parties can be found here: http://unfccc.int/national_reports/biennial_reports_and_iar/submitted_biennial_reports/items/7550.php

Each report is subject to a review¹⁹ process²⁰, assisted by the UNFCCC Secretariat and implemented by experts selected from the UNFCCC Roster of Experts (RoE).

Non-Annex I Parties Report:

- National Communications (NC), containing information on national circumstances, national GHG emissions/ removals²¹, steps taken or envisaged to implement the Convention, and any other information considered relevant to the achievement of the objective of the Convention including, if feasible, material relevant to calculations of global emissions and emission trends;
- Biennial Update Reports (BURs), containing updated information on national circumstances and institutional arrangements for reporting on a continuous basis²², national GHG emissions/removals information²³, including a national inventory report, and information on mitigation actions²⁴, effects, needs, and support received.

National communications may be submitted (decision 10/CP.2) by non-Annex I Parties every 4 years following decisions for each submission taken by the Conference of the Parties (COP). They are prepared and reported periodically by non-Annex I Parties based on agreed reporting guidelines (decision 17/CP.8)²⁵ based on methodologies developed by the IPCC²⁶ and adopted by the COP. Submissions by non-Annex I can be found here: http://unfccc.int/national_reports/non-annex_i_natcom/items/2979.php

²² This includes, for REDD+ activities, the national forest monitoring system, including for providing information on how the safeguards are being addressed and respected (decision 1/CP.16).

¹⁸ UNFCCC Biennial Reporting Guidelines for Developed Country Parties (Decision 2/CP.17) (at http://unfccc.int/resource/docs/2011/cop17/eng/09a01.pdf)

¹⁹ The review process is ruled by the Annex to Decision 13/CP.20 (Guidelines for the technical review of information reported under the Convention related to greenhouse gas inventories, biennial reports and national communications by Parties included in Annex I to the Convention) at http://unfccc.int/files/meetings/lima_dec_2014/decisions/application/pdf/auv_cop20_aireviewgdls.pdf

The objectives of the review of information reported under the Convention related to GHG inventories, BRs and NCs and pursuant to relevant decisions of the COP are the following:

a. To provide, in a facilitative, non-confrontational, open and transparent manner, a thorough, objective and comprehensive technical review of all aspects of the implementation of the Convention by individual Annex I Parties and Annex I Parties as a whole;

b. To promote the provision of consistent, transparent, comparable, accurate and complete information by Annex I Parties;

c. To assist Annex I Parties in improving their reporting of information contained in GHG inventories, BRs and NCs and pursuant to other relevant decisions of the COP and the implementation of their commitments under the Convention;

d. To ensure that the COP has accurate, consistent and relevant information in order to review the implementation of the Convention.

 $^{^{21}}$ For the years 1994 (1st NC), and 2000 (2nd NC).

²³ The decision text has not fixed the starting year nor the time-series of GHG estimates to be reported in the BUR. Anyhow the pace of the time series will be biennial from 2014 onwards.

²⁴ i.e. Nationally Appropriate Mitigation Actions (NAMAs), REDD+ activities.

²⁵ Guidelines for the preparation of national communications from Parties not included in Annex I to the Convention (decision 17/CP.8) at http://unfccc.int/resource/docs/cop8/07a02.pdf.

²⁶ Currently for non-Annex I Parties, Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (at http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html) have been adopted and 2000 IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (at http://www.ipcc-nggip.iges.or.jp/public/gp/english/index.html) and 2003 IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry (at http://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf/gpglulucf.html) have been encouraged to be used (see Decision 17/CP.8). Note that for the LULUCF (Land Use, Land-Use Change & Forestry) sector methodologies provided in the 2003 GPG for LULUCF replace those provided in the Revised 1996 IPCC Guidelines. Although, non-Annex I Parties may use the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (at http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html) and any further IPCC supplement to these Guidelines as adopted under the UNFCCC.

Biennial Update Reports are to be submitted (2/CP.17) by non-Annex I Parties every 2 years, and are prepared on the basis of agreed reporting guidelines (decision 2/CP.17)²⁷ based on methodologies developed by the IPCC²⁸ and adopted by the COP. Least developed country Parties and small island developing States may submit biennial update reports at their discretion. The first biennial report (BUR1) is due by December 2014 and it is expected to contain information on current levels and trends of GHG emissions and removals within their territories.

The Biennial Update Reports will be subject²⁹ to a technical assessment³⁰ as part of the International Consultation and Analysis process, which is aimed at increasing the transparency of mitigation actions and their effects.

For activities (*commonly referred ad REDD*+) listed in Decision 1/CP.16, paragraph 70³¹:

• Information, on the Forest Reference Emissions Levels and/or Forest Reference Levels, is prepared on the basis of agreed reporting guidelines (Decision 12/CP.17)³² and methodologies developed by the IPCC³³. The information is subject to assessment³⁴;

²⁷ UNFCCC biennial update reporting guidelines for Parties not included in Annex I to the Convention (Decision 2/CP.17) can be found at http://unfccc.int/resource/docs/2011/cop17/eng/09a01.pdf.

Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (at http://www.ipcc-nggip.iges.or.jp/public/gl/invs1.html), 2000 IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (at http://www.ipcc-nggip.iges.or.jp/public/gp/english/index.html) and 2003 IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry (at http://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf.html) have to be used for reporting (see Annex III to Decision 2/CP.17). Note that for the LULUCF (Land Use, Land-Use Change & Forestry) sector methodologies provided in the 2003 GPG for LULUCF replace those provided in the Revised 1996 IPCC Guidelines. Although, non-Annex I Parties may use the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (at http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html) and any further IPCC supplement to these Guidelines as adopted under the LINECCC

²⁹ Decision 2/CP.17 (Outcome of the work of the Ad Hoc Working Group on Long-term Cooperative Action under the Convention) at http://unfccc.int/resource/docs/2011/cop17/eng/09a01.pdf#page=4

³⁰ Decision 20/CP.19 (Composition, modalities and procedures of the team of technical experts under international consultation and analysis) at http://unfccc.int/resource/docs/2013/cop19/eng/10a02.pdff.

³¹ Developing country Parties are encouraged to contribute to mitigation actions in the forest sector by undertaking the following activities, as deemed appropriate by each Party and in accordance with their respective capabilities and national circumstances:

a. Reducing emissions from deforestation;

b. Reducing emissions from forest degradation;

c. Conservation of forest carbon stocks;

d. Sustainable management of forests;

e. Enhancement of forest carbon stocks.

 $^{^{32} \} Guidelines \ for \ submissions \ of \ information \ on \ reference \ levels; \ at \ http://unfccc.int/resource/docs/2011/cop17/eng/09a02.pdf\#page=16.$

Methodologies of the most recent IPCC Guidelines, which have been either adopted or encouraged to be used by non-Annex I Parties, should be applied, as appropriate. Currently:

for compiling the NC of non-Annex I Parties, the use of the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories
have been adopted and the 2000 IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories
and the 2003 IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry has been encouraged (see Decision 17/CP.8).
 Note that for the LULUCF (Land Use, Land-Use Change & Forestry), the sector methodologies provided in the 2003 GPG for LULUCF
replace those provided in the Revised 1996 IPCC Guidelines;

for compiling the BUR of non-Annex I Parties, the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories, the 2000 IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories and the 2003 IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry have been adopted (see Annex III to Decision 2/CP.17). Note that for the LULUCF (Land Use, Land-Use Change & Forestry), the sector methodologies provided in the 2003 GPG for LULUCF replace those provided in the Revised 1996 IPCC Guidelines.

Although, non-Annex I Parties may use the 2006 IPCC Guidelines for National Greenhouse Gas Inventories and any further IPCC supplement to these Guidelines as adopted under the UNFCCC.

³⁴ Annex to Decision 13/CP.19 (Guidelines and procedures for the technical assessment of submissions from Parties on proposed forest reference emission levels and/or forest reference levels) at http://unfccc.int/resource/docs/2013/cop19/eng/10a01.pdf.

- Information on safeguard is to be reported³⁵;
- For receiving payments for results-based actions, information on forest-related emissions by sources and removals by sinks resulting from the implementation of the activities is prepared on the basis of agreed reporting guidelines (Decision 14/CP.19)³⁶ and methodologies developed by the IPCC³⁷ and is to be reported, in an annex to the BUR. The information is assessed³⁸, as part of the ICA process.

35 Annex to Decision 12/CP.17 (Guidance on systems for providing information on how safeguards are addressed and respected) at http://unfccc.int/resource/docs/2011/cop17/eng/09a02.pdf#page=16.

³⁷ See footnote 35.

³⁶ Guidelines for elements to be included in the technical annex referred to in paragraph 7 of decision 14/CP.19 as contained in the Annex to Decision 14/CP.19 (*Modalities for measuring, reporting and verifying*) at http://unfccc.int/resource/docs/2013/cop19/eng/10a01.pdf#page=39

³⁸ Information on results of activities should be reported accordingly with the Guidelines for elements to be included in the technical annex referred to in paragraph 7 of Decision 14/CP.19 as provided in the Annex to Decision 14/CP.19 (Modalities for measuring, reporting and verifying) and their assessment occurs as part of the ICA process taking into account also provisions of Decision 14/CP.19 (Modalities for measuring, reporting and verifying) http://unfccc.int/resource/docs/2013/cop19/eng/10a01.pdf

References

- **Alexandratos, N. & J. Bruinsma.** 2012. *World agriculture towards* 2030/2050: the 2012 revision. ESA Working paper No. 12-03. Rome, FAO. Available at http://www.fao.org/fileadmin/templates/esa/Global_persepctives/world_ag_2030_50_2012_rev.pdf. Accessed on day month year.
- EC-JR. 2010. European Soil Portal. http://eusoils.jrc.ec.europa.eu/projects/RenewableEnergy.
- Eggleston H.S., Buendia L., Miwa K., Ngara T. & Tanabe K. (Eds). 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, prepared by the National Greenhouse Gas Inventories Programme, IGES, Hayama, Japan. IPCC Publication. Available at http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html.
- **European Commission, Joint Research Centre (EC-JRC).** 2003. *Global Land Cover 2000 database*. Available at http://bioval.jrc.ec.europa.eu/products/glc2000/glc2000.php
- **European Commission, Joint Research Centre (EC-JRC).** 2010. European Soil Portal. http://eusoils.jrc.ec.europa.eu/projects/RenewableEnergy
- **FAO/IIASA/ISRIC/ISSCAS/JRC.** 2012. Harmonized World Soil Database (Version 1.2). FAO, Rome, Italy and IIASA, Laxenburg, Austria. Available at http://www.fao.org/soils-portal/soil-survey/soil-maps-and-databases/harmonized-world-soil-database-v12/en/.
- **FAO.** 2005. *Global Forest Resources Assessment 2005*, FAO Forestry Paper 147, FAO Publication, Rome. Available at http://www.fao.org/forestry/fra/fra2005/en/
- **FAO.** 2010. *Global Forest Resources Assessment 2010*, FAO Forestry Paper 163, FAO Publication, Rome. Available at http://www.fao.org/forestry/fra/fra2010/en/.
- **FAO.** 2010. "FRA 2010 Terms and Definitions", *Global Forest Resources Assessment 2010* website. FAO Publication, Rome. Available at http://www.fao.org/docrep/014/am665e/am665e00.pdf
- **FAO.** 2012. Global Ecological Zones for FAO Forest Reporting: 2010 Update. Forest Resources Assessment Working Paper 179. FAO, Rome.
- **FAO.** 2012. "Glossary". Methods & Standards, FAOSTAT database website. Available http://faostat3.fao.org/faostat-gateway/go/to/mes/glossary/*/E.
- **FAO.** 2012. Classification and Standards section, Statistics Division, Economic and Social Development Department website. Available at http://www.fao.org/waicent/faoinfo/economic/faodef/faodefe.htm.
- **FAO.** 2012. Data Structure, Concepts and Definitions common to FAOSTAT and CountrySTAT framework, Statistics Division, Economic and Social Development Department, website. Available at https://dl.dropboxusercontent.com/u/37798211/concepts_definitions.pdf
- **FAO.** 2012. Statistics Division, Economic and Social Development Department web site. Agri-environmental methodology. Available on FAOSTAT Fertilizers statistics methodology at http://www.fao.org/economic/ess/ess-agri/ess-resource-meth/en/
- FAO. AQUASTAT Database. Available at http://www.fao.org/nr/water/aquastat/main/index.stm
- **Giglio, L., Randerson, J.T., & van der Werf, G. R.** 2013. Analysis of daily, monthly, and annual burned area using the fourth generation Global Fire Emissions Database (GFED4). *Journal of Geophysical Research: Biogeosciences*, 118: 1-12.
- Giri, C., Zhu, Z., Reed, B. 2005. A comparative analysis of the Global Land Cover 2000 and MODIS land cover datasets. *Remote Sensing of Environment* 94: 123–132. Available at http://www.sciencedirect.com/science/article/pii/S0034425704002895.
- **Global Forest Observation Initiative.** 2014. *GFOI portal. Methods & Guidance Documentation*. Available at http://gfoi.org/methods-guidance-documentation
- **Hansen, M. C., DeFries, R. S., Townshend, J. R. G., & Sohlberg, R.** 2000. Global land cover classification at 1km spatial resolution using a classification tree approach. *International Journal of Remote Sensing*, 21: 1331–1364.
- **IEA.** 2013. Statistics. Available at http://www.iea.org/statistics/
- **IEA.** 2013. *Statistics, Balance Definitions, Oil Products*. Available at http://www.iea.org/statistics/resources/balancedefinitions/#fueloil.

- **IIASA/FAO.** 2012. *Global Agro-Ecological Zones* (GAEZ v3.0), IIASA, Laxenburg, Austria, and FAO, Rome. Available at http://www.fao.org/nr/gaez
- **IPCC.** 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Glossary, Japan. Available at http://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf_files/Glossary_Acronyms_BasicInfo/Glossary.pdf
- **IPCC.** 2002. Background Papers, IPCC Expert Meetings on Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC-NGGIP, pp. 399-417.
- **IPCC.** 2000. Good practice guidance and uncertainty management in national greenhouse gas inventories. In: J. Penman *et al.* (Eds.), IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Japan.
- **IPCC.** 1996. Climate Change 1995 The Science of Climate Change: Contribution of Working Group I to the Second Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.
- IPCC. 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories. OECD, Paris, France.
- **OECD/IEA.** 2005. *Energy Statistics Manual*. Prepared by the Energy Statistics Division (ESD) of the International Energy Agency (IEA) in co-operation with the Statistical Office of the European Communities (Eurostat). Paris, France.
- **OECD/IEA.** 2012. *CO*₂ *emissions from fuel combustion Highlights 2012 edition*. Paris, France. Available at http://www.iea.org/publications/freepublications/publication/name,4010,en.html
- **Stout B.** 1990. *Handbook of Energy for World Agriculture*. Elsevier Science Publishers LTD New York, USA **United Nations.** 2014. UN Data portal. Available at http://data.un.org/
- **UNSD.** 2011. *International Recommendations for Energy Statistics (IRES) Draft version*. Prepared by the United Nations Statistics Division, New York, USA. Available at http://unstats.un.org/unsd/energy/ires
- **UNSD.** 2011. Energy Balances and Electricity Profiles Concepts and definitions Glossary. Available at http://unstats.un.org/unsd/energy/balance/concepts.htm
- Wint, W. & Robinson, T. 2007. Gridded livestock of the world. FAO Publication, FAO, Rome.
- You, L., Wood, S., Sebastian, K. 2008. Comparing and Synthesizing Different Global Agricultural Land Datasets for Crop Allocation Modeling. ISPRS Proceedings of the 21st Congress, Beijing. Available at http://www.isprs.org/proceedings/XXXVII/congress/7_pdf/9_ThS-17/06.pdf.

This Manual seeks to guide the staff of national statistical offices and environmental ministries and agencies in compiling statistics related to GHG emissions and removals. In particular, the Manual provides information on accessing and using the FAOSTAT Emissions database. In addition, the manual sets out a step-by-step approach on estimating GHG emissions with the 2006 IPCC Guidelines for National GHG Emission Inventories.

The FAOSTAT Emissions database provides Member Countries with the opportunity to identify the official agricultural and forestry statistics (data gap filling) required for GHG emission estimations, and to perform GHG data analysis linked to data gap filling and Quality Assurance/Quality Control processes.

Global Strategy To Improve Agricultural & Rural Statistics

Statistics Division (ESS)

Food and Agriculture Organization of the United Nations Viale delle Terme di Caracalla, 00153 Rome, Italy

ESS-Global-Strategy@fao.org

www.fao.org/economic/ess/globalstrategy

